

NON-PERTURBATIVE APPROACH TO THE BOURGAIN–SPENCER CONJECTURE IN STOCHASTIC HOMOGENIZATION

MITIA DUERINCKX

ABSTRACT. In the context of stochastic homogenization, the Bourgain–Spencer conjecture states that the ensemble-averaged solution of a divergence-form linear elliptic equation with random coefficients admits an intrinsic description in terms of higher-order homogenized equations with an accuracy four times better than the almost sure solution itself. While previous rigorous results were restricted to a perturbative regime with small ellipticity ratio, we prove the first half of this conjecture for the first time in a non-perturbative setting. Our approach involves the construction of a new corrector theory in stochastic homogenization: while only a bounded number of correctors can be constructed as stationary L^2 random fields, we show that twice as many stationary correctors can be defined in a Schwartz-like distributional sense on the probability space.

CONTENTS

1. Introduction	1
2. Main results	6
3. Preliminary: higher-order homogenization	9
4. Functional setup	12
5. Weak corrector theory	19
6. Effective description of ensemble averages	29
Appendix A. Intrinsic description of weak fluctuations	34
Acknowledgements	38
References	38

1. INTRODUCTION

Let \mathbf{a} be a stationary and ergodic random coefficient field on \mathbb{R}^d with values in $\mathbb{R}^{d \times d}$, with the following boundedness and ellipticity properties, for some $\lambda > 0$,

$$|\mathbf{a}(x)e| \leq |e|, \quad e \cdot \mathbf{a}(x)e \geq \lambda|e|^2, \quad \text{for all } x, e \in \mathbb{R}^d, \quad (1.1)$$

and denote by (Ω, \mathbb{P}) the underlying probability space. In the sequel, we further assume that \mathbf{a} satisfies some strong mixing condition, and our main results focus for simplicity on the Gaussian setting, see Section 2.1. Given a deterministic vector field $f \in C_c^\infty(\mathbb{R}^d)^d$, we consider the random family $(\nabla u_{\varepsilon, f})_{\varepsilon > 0}$ of unique almost sure gradient solutions in $L^2(\mathbb{R}^d)^d$ of the rescaled Poisson problems

$$-\nabla \cdot \mathbf{a}\left(\frac{\cdot}{\varepsilon}\right) \nabla u_{\varepsilon, f} = \nabla \cdot f, \quad \text{in } \mathbb{R}^d, \quad (1.2)$$

where the rescaled coefficients $\mathbf{a}\left(\frac{\cdot}{\varepsilon}\right)$ vary on a microscopic scale $\varepsilon > 0$. While stochastic homogenization theory has been focussing on the intrinsic description of spatial oscillations

and random fluctuations of the random solution $\nabla u_{\varepsilon, f}$ in the macroscopic limit $\varepsilon \downarrow 0$, the present work is rather concerned with the effective description of the ensemble-averaged solution $\mathbb{E}[\nabla u_{\varepsilon, f}]$.

Before describing the Bourgain–Spencer conjecture and our results, in order to contextualize them properly, we start with a review of the state of the art in higher-order stochastic homogenization. First, the qualitative theory [29, 24] states the almost sure weak convergence of fields $\nabla u_{\varepsilon, f} \rightharpoonup \nabla \bar{u}_f^1$ and fluxes $\mathbf{a}(\frac{\cdot}{\varepsilon})\nabla u_{\varepsilon, f} \rightharpoonup \bar{\mathbf{a}}^1\nabla \bar{u}_f^1$ in $L^2(\mathbb{R}^d)^d$, where $\nabla \bar{u}_f^1$ is the unique gradient solution in $L^2(\mathbb{R}^d)^d$ of the (deterministic) homogenized equation

$$-\nabla \cdot \bar{\mathbf{a}}^1 \nabla \bar{u}_f^1 = \nabla \cdot f, \quad \text{in } \mathbb{R}^d, \quad (1.3)$$

where the effective coefficients $\bar{\mathbf{a}}^1 \in \mathbb{R}^{d \times d}$ are given in each direction e_i , $1 \leq i \leq d$, by

$$\bar{\mathbf{a}}^1 e_i = \mathbb{E}[\mathbf{a}(\nabla \varphi_i^1 + e_i)],$$

in terms of the so-called corrector gradient $\nabla \varphi_i^1$ in the direction e_i , which is defined as the unique almost sure gradient solution in $L_{\text{loc}}^2(\mathbb{R}^d)^d$ of

$$-\nabla \cdot \mathbf{a}(\nabla \varphi_i^1 + e_i) = 0, \quad \text{in } \mathbb{R}^d, \quad (1.4)$$

such that $\nabla \varphi_i^1$ is a stationary field with vanishing expectation and with bounded second moment. This convergence result for fields and fluxes is understood as follows: the heterogeneous field-flux constitutive law $\nabla w_\varepsilon \mapsto \mathbf{a}(\frac{\cdot}{\varepsilon})\nabla w_\varepsilon$ is replaced on large scales by the effective law $\nabla \bar{w} \mapsto \bar{\mathbf{a}}^1 \nabla \bar{w}$.

As the coefficient field $\mathbf{a}(\frac{\cdot}{\varepsilon})$ is oscillating on scale $O(\varepsilon)$, fields and fluxes display oscillations on the same scale, which is why their convergence is only weak in $L^2(\mathbb{R}^d)^d$. Homogenization theory has aimed at describing these oscillations intrinsically by means of two-scale expansions. The key ingredient is the corrector $\varphi^1 = \{\varphi_i^1\}_{1 \leq i \leq d}$, which is defined in (1.4) up to an additive random constant, and which is almost surely sublinear at infinity as a standard consequence of the ergodic theorem. This corrector is interpreted as correcting Euclidean coordinates $\{x \mapsto x_i\}_{1 \leq i \leq d}$ into \mathbf{a} -harmonic ones $\{x \mapsto x_i + \varphi_i^1(x)\}_{1 \leq i \leq d}$. The following so-called two-scale expansion

$$\mathcal{F}_\varepsilon^1[\bar{u}_f^1] := \bar{u}_f^1 + \varepsilon \varphi_i^1(\frac{\cdot}{\varepsilon}) \nabla_i \bar{u}_f^1 \quad (1.5)$$

is then viewed as an intrinsic Taylor expansion of the limiting profile \bar{u}_f^1 in \mathbf{a} -harmonic coordinates. (Throughout, we use Einstein's summation convention on repeated indices.) The classical corrector result [29, 24] ensures that this expansion indeed captures leading-order oscillations in the sense that almost surely

$$\nabla u_{\varepsilon, f} - \nabla \mathcal{F}_\varepsilon^1[\bar{u}_f^1] \rightarrow 0, \quad \text{strongly in } L^2(\mathbb{R}^d)^d.$$

In the last decade, many works have managed to further establish sharp convergence rates, which mainly requires to investigate quantitatively the sublinearity of the corrector φ^1 ; see [17, 16, 4, 3]. In dimension $d > 2$, the corrector can be chosen itself as a stationary field with bounded moments: it is then uniquely defined up to an additive *deterministic* constant, which is fixed for instance by choosing $\mathbb{E}[\varphi^1] = 0$. In dimension $d \leq 2$, the corrector cannot be chosen stationary as it has some nontrivial (sublinear) growth at

infinity. Based on optimal corrector estimates, it is deduced that for all $q < \infty$,

$$\mathbb{E} \left[\|\nabla u_{\varepsilon, f} - \nabla \mathcal{F}_\varepsilon^1[\bar{u}_f^1]\|_{L^2(\mathbb{R}^d)}^q \right]^{\frac{1}{q}} \lesssim_{f, q} \varepsilon \times \begin{cases} 1 & : d > 2, \\ |\log \varepsilon|^{\frac{1}{2}} & : d = 2, \\ \varepsilon^{-\frac{1}{2}} & : d = 1. \end{cases}$$

Next, we recall how this intrinsic description of oscillations is pursued to higher orders. While the first corrector φ^1 is defined to correct Euclidean coordinates into \mathbf{a} -harmonic ones, higher-order correctors are defined iteratively to correct higher-order polynomials and make them adapted to the heterogeneous elliptic operator $\nabla \cdot \mathbf{a} \nabla$. More precisely, assuming that correctors $\varphi^1, \dots, \varphi^{n-1}$ are defined as stationary fields, say with vanishing expectation, the next-order corrector φ^n is uniquely defined up to an additive random constant by the following properties:

- for any n th-order polynomial \bar{q} , the corrected polynomial $\mathcal{F}^n[\bar{q}] := \sum_{k=0}^n \varphi_{i_1 \dots i_k}^k \nabla_{i_1 \dots i_k}^k \bar{q}$ captures oscillations of the operator in the sense that $\nabla \cdot \mathbf{a} \nabla \mathcal{F}^n[\bar{q}]$ is deterministic;
- the gradient $\nabla \varphi^n$ is a stationary field with vanishing expectation and with bounded second moment.

In fact, this definition implies $\nabla \cdot \mathbf{a} \nabla \mathcal{F}^n[\bar{q}] = \nabla \cdot (\sum_{k=1}^{n-1} \bar{\mathbf{a}}^k \nabla^{k-1}) \nabla \bar{q}$ for some higher-order effective tensors $\bar{\mathbf{a}}^1, \dots, \bar{\mathbf{a}}^{n-1}$. As recalled in Section 3.2, higher-order correctors are alternatively defined by a hierarchy of cell problems, and higher-order tensors are given by

$$\bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k = \mathbb{E}[\mathbf{a}(\nabla \varphi_{i_1 \dots i_k}^k + \varphi_{i_1 \dots i_{k-1}}^{k-1} e_{i_k})]. \quad (1.6)$$

The latter define corrections to the effective field-flux constitutive law: while the heterogeneous law $\nabla w_\varepsilon \mapsto \mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla w_\varepsilon$ is replaced on large scales by the effective law $\nabla \bar{w} \mapsto \bar{\mathbf{a}}^1 \nabla \bar{w}$ at leading order, dispersive corrections must be included when looking for finer accuracy,

$$\nabla \bar{w} \mapsto \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{w} := \sum_{k=1}^n \varepsilon^{k-1} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla_{i_1 \dots i_{k-1}}^{k-1} \bar{w}.$$

To order n , the homogenized equation (1.3) is formally replaced by

$$-\nabla \cdot \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{U}_{\varepsilon, f}^n = \nabla \cdot f, \quad \text{in } \mathbb{R}^d. \quad (1.7)$$

This equation is however ill-posed in general for $n \geq 2$, as the symbol may be indefinite, and we use a suitable proxy $\nabla \bar{\mathcal{U}}_\varepsilon^n[f]$ for $\nabla \bar{U}_{\varepsilon, f}^n$, see Definition 3.3. In these terms, the two-scale expansion (1.5) is replaced by its higher-order version

$$\mathcal{F}_\varepsilon^n[\bar{\mathcal{U}}_\varepsilon^n[f]] := \sum_{k=0}^n \varepsilon^k \varphi_{i_1 \dots i_k}^k(\frac{\cdot}{\varepsilon}) \nabla_{i_1 \dots i_k}^k \bar{\mathcal{U}}_\varepsilon^n[f]. \quad (1.8)$$

In case of a periodic coefficient field \mathbf{a} , all correctors can be constructed as bounded periodic fields, and the approximation $\nabla u_{\varepsilon, f} - \nabla \mathcal{F}_\varepsilon^n[\bar{\mathcal{U}}_\varepsilon^n[f]] = O(\varepsilon^n)$ can be justified to all orders. In contrast, a key specificity of the random setting is that only a finite number of correctors can be defined as well-behaved stationary objects. More precisely, as shown in [19, 7, 14], the corrector φ^n can be chosen stationary with bounded moments only for $n < \ell := \lfloor \frac{d}{2} \rfloor$. Although higher-order correctors could still be defined, their lack of stationarity and their spatial growth would make them unsuitable for error estimates. The accuracy of the two-scale expansion (1.8) is then shown to saturate at order $O(\varepsilon^{d/2})$: for all $q < \infty$,

$$\mathbb{E} \left[\|\nabla u_{\varepsilon, f} - \nabla \mathcal{F}_\varepsilon^\ell[\bar{\mathcal{U}}_\varepsilon^\ell[f]]\|_{L^2(\mathbb{R}^d)}^q \right]^{\frac{1}{q}} \lesssim_{f, q} \varepsilon^{\frac{d}{2}} \times \begin{cases} 1 & : d \text{ odd}, \\ |\log \varepsilon|^{\frac{1}{2}} & : d \text{ even}, \end{cases} \quad (1.9)$$

and this accuracy is optimal as random fluctuations are known to become dominant at that order $O(\varepsilon^{d/2})$, see [13, 14].

In this context, the present work is devoted to the effective description of the ensemble-averaged field $\mathbb{E}[\nabla u_{\varepsilon,f}]$ and flux $\mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon})\nabla u_{\varepsilon,f}]$ with optimal accuracy. Although relevant for applications, this question has only been addressed very recently by Sigal [31]. A naïve estimate follows by taking the expectation in (1.9), which gives

$$\|\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{U}_{\varepsilon}^{\ell}[f]\|_{L^2(\mathbb{R}^d)} \lesssim_f \varepsilon^{\frac{d}{2}} \times \begin{cases} 1 & : d \text{ odd,} \\ |\log \varepsilon|^{\frac{1}{2}} & : d \text{ even.} \end{cases} \quad (1.10)$$

While the limitation to accuracy $O(\varepsilon^{d/2})$ in (1.9) is related to random fluctuations, the optimality argument fails for (1.10), thus indicating that this effective description of the ensemble-averaged field might be pursued to higher orders. Following preliminary calculations by Sigal [31] and Spencer, a striking work by Bourgain [8] and its key refinement by Kim and Lemm [23] have suggested that the expansion might be pursued in fact almost to order $O(\varepsilon^{2d})$, see [11]. More precisely, the following so-called Bourgain–Spencer conjecture was formulated. (The case of dimension $d = 1$ is naturally omitted as a direct computation simply yields $\mathbb{E}[\nabla u_{\varepsilon,f}] = \nabla \bar{u}_f^1$ in that case.)

Conjecture 1 (Bourgain, Spencer). *Let $d > 1$. There exists a collection $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq 2d}$ of constant tensors such that for all $\eta > 0$,*

$$\|\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{U}_{\varepsilon}^{2d}[f]\|_{L^2(\mathbb{R}^d)} \lesssim_{f,\eta} \varepsilon^{2d-\eta},$$

where $\nabla \bar{U}_{\varepsilon}^{2d}[f]$ is a proxy for the solution of the higher-order homogenized equation (1.7) with coefficients $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq 2d}$. \diamond

Bourgain’s work [8] and its refinement by Kim and Lemm [23] state that this conjecture holds to order $O(\varepsilon^{2d-\eta})$ provided that the random coefficient field \mathbf{a} has small enough ellipticity ratio $|\mathbf{a} - \text{Id}| \ll \eta$. This perturbative result was obtained in the model framework of discrete elliptic equations with iid coefficients, but can be extended to the present continuum setting. Note that this result was originally expressed in terms of harmonic analysis: as explained in [11], there exists a uniformly elliptic matrix-valued kernel \mathcal{B} such that the ensemble-averaged field satisfies

$$-\nabla \cdot \mathcal{B}(\varepsilon \nabla) \mathbb{E}[\nabla u_{\varepsilon,f}] = \nabla \cdot f,$$

and Sigal proposed to investigate the C^α regularity of \mathcal{B} at the origin, which turns out to be indeed equivalent to an effective description to order $O(\varepsilon^\alpha)$ as in the above conjecture.

Beyond the perturbative results by Bourgain [8] and by Kim and Lemm [23], the conjecture concerns the corresponding conclusion in a non-perturbative regime. While their arguments proceed by a fine analysis of perturbative expansions at small ellipticity ratio, we propose a novel argument based on non-perturbative quantitative homogenization techniques. Our main result proves the validity of the first half of the conjecture, that is, the accuracy of an effective description almost to order $O(\varepsilon^d)$, see Theorem 2 below. Our argument splits into two parts:

(i) *Weak corrector theory:*

Although higher-order correctors φ^n with $n \geq \ell = \lceil \frac{d}{2} \rceil$ cannot be constructed as stationary fields with bounded moments, we show that stationary correctors can be

constructed up to order $n < d$ in a Schwartz-like distributional sense on the probability space. More precisely, for $n < d$, weak-type expressions of the form $\mathbb{E}[X\varphi^n(x)]$ can be meaningfully defined for all “smooth and local” test random variables X , in a way that is compatible with corrector equations and stationarity. At order $n = d$, only the corrector gradient $\nabla\varphi^d$ can be defined as a stationary object in the same weak sense. Denoting by $\mathcal{S}(\Omega)$ the space of suitable test random variables, correctors $\{\varphi^n\}_{1 \leq n \leq d}$ are thus defined as (smooth) maps $\mathbb{R}^d \rightarrow \mathcal{S}'(\Omega)$ with values in the space of linear functionals on $\mathcal{S}(\Omega)$. The Gelfand triple $\mathcal{S}(\Omega) \subset L^2(\Omega) \subset \mathcal{S}'(\Omega)$ is viewed as a stochastic version of the usual triple $\mathcal{S}(\mathbb{R}^d) \subset L^2(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$ of Schwartz test functions and distributions. We refer to Section 2.3 for details.

(ii) *Weak two-scale expansions:*

In terms of these weak correctors $\{\varphi^n\}_{1 \leq n \leq d}$, we may construct the corresponding two-scale expansion $\mathcal{F}_\varepsilon^d[\bar{\mathcal{U}}_\varepsilon^d[f]]$ as in (1.8). The difficulty is that $\nabla\mathcal{F}_\varepsilon^d[\bar{\mathcal{U}}_\varepsilon^d[f]]$ is only defined in the weak sense of $\mathcal{S}'(\Omega; L^2(\mathbb{R}^d))$ and can in particular not be used in any energy estimate as is usually done in the classical theory. By duality, we however manage to prove the accuracy of this two-scale expansion in a suitable weak sense, and we deduce the validity of Conjecture 1 almost to order $O(\varepsilon^d)$. We refer to Section 2.2 for details.

Such a description of the field $\nabla u_{\varepsilon, f}$ in a distributional sense on the probability space was originally inspired by our work with Shirley [15] on random Schrödinger operators, and it constitutes a new type of results in the field. Note that spaces of Schwartz-like distributions on the probability space first appeared in the works of Kondratiev and Hida in the context of SPDEs, see e.g. [21] and references therein, but our definition is quite different and is closer to [2], see Section 4.4.

Notation.

- We denote by $C \geq 1$ any constant that only depends on the dimension d , on the ellipticity constant λ in (1.1), and on $\|a_0\|_{W^{1,\infty}}$ and $\int_{\mathbb{R}^d} [C_0]_\infty$ in (2.1) and (2.3) below. We use the notation \lesssim (resp. \gtrsim) for $\leq C \times$ (resp. $\geq \frac{1}{C} \times$) up to such a multiplicative constant C . We write \simeq when both \lesssim and \gtrsim hold. The notation \ll stands for $\leq \frac{1}{C} \times$ for some large enough constant C . We add subscripts to C , \lesssim , \gtrsim , \simeq , \ll to indicate dependence on other parameters.
- For a matrix field $H = \{H_{ij}\}_{1 \leq i, j \leq d}$, we define its divergence $\nabla \cdot H$ as the vector field given by $(\nabla \cdot H)_i := \nabla_j H_{ij}$, where we use Einstein’s summation convention on repeated indices. For a vector field $h = \{h_i\}_{1 \leq i \leq d}$, we define its curl as the skew-symmetric matrix field given by $(\nabla \times h)_{ij} := \nabla_i h_j - \nabla_j h_i$.
- The ball centered at x of radius r in \mathbb{R}^d is denoted by $B_r(x)$, and we simply write $B(x) := B_1(x)$, $B_r := B_r(0)$, and $B := B_1(0)$. We also write $Q_r := [-\frac{r}{2}, \frac{r}{2}]^d$ for the cube centered at the origin with side length r .
- For $r \in \mathbb{R}$ we denote by $\lceil r \rceil$ the smallest integer $\geq r$, and we recall the notation $\ell = \lceil \frac{d}{2} \rceil$. For $r, s \in \mathbb{R}$ we write $r \vee s := \max\{r, s\}$ and $r \wedge s := \min\{r, s\}$. For $x \in \mathbb{R}^d$ we set $\langle x \rangle := (1 + |x|^2)^{1/2}$, and similarly $\langle \nabla \rangle := (1 - \Delta)^{1/2}$.
- For a function g and for $1 \leq p < \infty$, we write $[g]_p(x) := (f_{B(x)} |g|^p)^{1/p}$ for the local moving L^p average, and similarly $[g]_\infty(x) := \sup_{B(x)} |g|$. For averages at the scale ε , we write $[g]_{p;\varepsilon}(x) := (f_{B_\varepsilon(x)} |g|^p)^{1/p}$.

2. MAIN RESULTS

This section is devoted to the statement and discussion of our main results. For the reader's convenience, detailed statements are postponed to the proof sections.

2.1. Assumptions. We focus on the model Gaussian setting for the random coefficient field \mathbf{a} , in which case Malliavin calculus is available and substantially simplifies the analysis. More precisely, we set

$$\mathbf{a}(x) := a_0(G(x)), \quad (2.1)$$

where $a_0 \in C_b^1(\mathbb{R}^\kappa)^{d \times d}$ is such that the ellipticity and boundedness assumptions (1.1) are pointwise satisfied, and where $G : \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^\kappa$ is an \mathbb{R}^κ -valued centered stationary Gaussian random field on \mathbb{R}^d with covariance function \mathcal{C} , constructed on a probability space (Ω, \mathbb{P}) .¹ Note that we write $G(x) = G(x, \cdot) : \Omega \rightarrow \mathbb{R}^\kappa$ for the evaluation at x . In addition, we assume that G has integrable correlations in the following sense: starting from the representation

$$G = \mathcal{C}_0 * \xi, \quad (2.2)$$

where ξ is a κ -dimensional Gaussian white noise on \mathbb{R}^d and where the kernel $\mathcal{C}_0 : \mathbb{R}^d \rightarrow \mathbb{R}^{\kappa \times \kappa}$ satisfies $\mathcal{C}_0 * \mathcal{C}_0 = \mathcal{C}$, we assume that \mathcal{C}_0 satisfies the integrability condition

$$\int_{\mathbb{R}^d} [\mathcal{C}_0]_\infty < \infty. \quad (2.3)$$

In particular, this entails that the covariance function \mathcal{C} satisfies the same integrability condition $\int_{\mathbb{R}^d} [\mathcal{C}]_\infty < \infty$. Moreover, \mathcal{C} is continuous, so that G and \mathbf{a} are stochastically continuous and jointly measurable on $\mathbb{R}^d \times \Omega$.

Restricting to this Gaussian setting is essential as our approach relies on Malliavin calculus techniques. Our analysis is easily repeated *mutatis mutandis* in the Poisson setting and in the iid discrete setting, using corresponding stochastic calculus tools that are available in those cases, e.g. [30, 9]. The general case of an α -mixing coefficient field is more involved and is postponed to future work. Regarding the integrability condition (2.3), it could be easily relaxed: the case of non-integrable correlations could be treated similarly but would yield different scalings; see also [16, 12, 10].

Without loss of generality, we can assume that the probability space (Ω, \mathbb{P}) is endowed with the σ -algebra generated by the underlying white noise ξ in (2.2). This countably generated σ -algebra coincides with the one generated by the Gaussian field G under assumptions (2.2)–(2.3). We then consider the following model subspace of “smooth and local” random variables,

$$\mathcal{S}(\Omega) := \left\{ g \left(\int_{\mathbb{R}^d} h_1 \cdot \xi, \dots, \int_{\mathbb{R}^d} h_n \cdot \xi \right) : n \in \mathbb{N}, g \in C_b^\infty(\mathbb{R}^n), \right. \\ \left. h_1, \dots, h_n \in C_c^\infty(\mathbb{R}^d)^\kappa \right\}, \quad (2.4)$$

and the choice of the σ -algebra ensures that this subspace is dense in $L^q(\Omega)$ for all $q < \infty$. We denote by $\mathcal{S}'(\Omega)$ the space of continuous linear functionals on $\mathcal{S}(\Omega)$, viewed as a space

¹By centered stationary Gaussian random field, we mean the following: $G = \{G(x)\}_{x \in \mathbb{R}^d}$ is a collection of \mathbb{R}^κ -valued Gaussian random variables with expectation $\mathbb{E}[G(x)] = 0$ and with covariance of the form $\mathbb{E}[G(x) \otimes G(y)] = \mathcal{C}(x - y)$, where \mathcal{C} belongs to $L^\infty(\mathbb{R}^d)^{\kappa \times \kappa}$ and satisfies $\mathcal{C}(-x) = \mathcal{C}(x)$.

of Schwartz-like distributions on the probability space; see Section 4.4. For $X \in \mathcal{S}'(\Omega)$ and $Y \in \mathcal{S}(\Omega)$, we use the abusive notation $\mathbb{E}[XY] = \langle X, Y \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)}$ for the duality product.

2.2. First half of the Bourgain–Spencer conjecture. The following main result justifies the effective description of the ensemble-averaged field and flux almost to order $O(\varepsilon^d)$, thus proving the first half of Conjecture 1. The non-perturbative extension to higher orders remains an open question. We refer to Theorem 6.1 for a more detailed statement.

Theorem 2 (Effective description of ensemble averages). *Let $d > 1$ and let the higher-order effective tensors $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq d}$ be defined in (2.6) below. For all $f \in C_c^\infty(\mathbb{R}^d)^d$, denoting by $\nabla \bar{U}_\varepsilon^d[f]$ a proxy for the solution of the associated higher-order homogenized equation (1.7) with $n = d$, see Definition 3.3, there holds for all $p > \frac{d}{\eta}$ and $0 < \eta \ll 1$,*

$$\begin{aligned} & \left\| \left[\mathbb{E}[\nabla u_{\varepsilon, f}] - \nabla \bar{U}_\varepsilon^d[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} \\ & + \left\| \left[\mathbb{E}[\mathbf{a}(\cdot/\varepsilon) \nabla u_{\varepsilon, f}] - \bar{\mathcal{A}}_\varepsilon^d \nabla \bar{U}_\varepsilon^d[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} \lesssim_{f, p, \eta} \varepsilon^{d-\eta}. \quad \diamond \end{aligned}$$

While the Bourgain–Spencer conjecture is concerned with ensemble averages, and while our proof is based on a two-scale expansion for $\nabla u_{\varepsilon, f}$ in the weak sense of $\mathcal{S}'(\Omega; L^2(\mathbb{R}^d))$, a natural side question concerns the intrinsic description of fluctuations $\nabla u_{\varepsilon, f} - \mathbb{E}[\nabla u_{\varepsilon, f}]$ in this weak sense; this is briefly addressed in Appendix A.

2.3. Weak corrector theory. While only the first $\ell = \lceil \frac{d}{2} \rceil$ correctors can be constructed as stationary fields with bounded moments, see Theorem 3.2 below, we show that twice as many stationary correctors can be constructed in the distributional sense on the probability space, which we expect to be optimal. We refer to Theorem 5.1 for a more detailed statement, where correctors are further controlled in some dual Malliavin–Sobolev spaces.

Theorem 3 (Weak correctors). *Let $d > 1$. Higher-order weak correctors $\{\varphi^n\}_{1 \leq n \leq d}$ are uniquely defined in (a suitable subspace of) $C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$ such that the standard cell problems (see (3.3)–(3.6)) are satisfied in a corresponding weak sense, such that φ^n is stationary with $\mathbb{E}[\varphi^n] = 0$ for all $n < d$, and such that $\nabla \varphi^d$ is stationary with $\mathbb{E}[\nabla \varphi^d] = 0$ and with the anchoring condition $\int_B \varphi^d = 0$. In addition, the following estimates hold:*

(i) Weak sublinearity of φ^d : for all $X \in \mathcal{S}(\Omega)$ and $\eta > 0$,

$$\left| \mathbb{E}[X \varphi^d(x)] \right| \lesssim_{X, \eta} \langle x \rangle^\eta.$$

(ii) Weak fluctuation scaling: for all $n < d$, $X \in \mathcal{S}(\Omega)$, $g \in C_c^\infty(\mathbb{R}^d)$, and $p < \frac{d}{n}$,

$$\left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \nabla \varphi^{n+1} \right] \right| + \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \varphi^n \right] \right| \lesssim_{X, p} \|g\|_{L^p(\mathbb{R}^d)}. \quad \diamond$$

Remarks 2.1. A few comments are in order.

(a) *Criticality at order d :*

Rather than constructing the corrector φ^n itself, the proof of Theorem 3 can be viewed as constructing its Malliavin derivative $D\varphi^n$ and monitoring its decay: item (ii) is formally understood as $D_z \nabla \varphi^n(x) = O(\langle x - z \rangle^{n-d-1})$ in a weak sense. The limitation is manifest: for $n > d$, no decay is left and the Malliavin derivative could at best be defined up to $(n - d)$ th-order \mathbf{a} -harmonic polynomials. Both the lack of uniqueness and the growth thus make it unclear how the theory could be extended to higher order.

Criticality at order d is new in stochastic homogenization, but the same holds in the following simple deterministic example: given a deterministic coefficient field of the form $\mathbf{a} = \text{Id} + \mathbf{c}$ with \mathbf{c} supported in the unit ball B , corresponding correctors $\{\psi^n\}_{1 \leq n \leq d}$ can be constructed as unique decaying solutions of cell problems (3.3)–(3.5) with $|\nabla \psi^n(x)| \lesssim \langle x \rangle^{n-d-1}$, while for $n > d$ correctors would similarly be defined only up to $(n-d)$ th-order harmonic polynomials.

(b) *Explicit 1D case:*

In dimension $d = 1$, the first corrector is explicit and its analysis provides an instructive illustration of the above result. Choosing it with anchoring $\phi^1(0) = 0$, it takes the form

$$\nabla \phi^1(x) := \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]^{-1} \left(\frac{1}{\mathbf{a}} - \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]\right), \quad \phi^1(x) := \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]^{-1} \int_0^x \left(\frac{1}{\mathbf{a}} - \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]\right).$$

Comparing $\frac{1}{\mathbf{a}} - \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]$ to a white noise, ϕ^1 is compared to a Brownian motion. The central limit theorem then yields $\|\phi^1(x)\|_{L^q(\Omega)} \simeq_q \langle x \rangle^{1/2}$ for all $q < \infty$, which entails that no random constant can be added to ϕ^1 to make it stationary. Formally, a stationary version of ϕ^1 would be given by

$$\varphi^1(x) := \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]^{-1} \int_{-\infty}^x \left(\frac{1}{\mathbf{a}} - \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]\right),$$

but this integral makes no pointwise sense. Such an object can however be defined as follows: assuming for simplicity that the covariance function \mathcal{C} is compactly supported in B , so that \mathbf{a} has a unit range of dependence, we can formally “compute” conditional expectations from the above formula: for all $R > 0$,

$$\mathbb{E}\left[\varphi^1(x) \mid \mathbf{a}|_{[-R,R]}\right] := \mathbb{E}\left[\frac{1}{\mathbf{a}}\right]^{-1} \int_{-(R+1)}^x \mathbb{E}\left[\frac{1}{\mathbf{a}} - \mathbb{E}\left[\frac{1}{\mathbf{a}}\right] \mid \mathbf{a}|_{[-R,R]}\right], \quad (2.5)$$

where the right-hand side now makes perfect sense as a bounded random variable. In other words, this stationary corrector φ^1 is only defined via its conditional expectations with respect to the coefficient field \mathbf{a} on bounded sets. This gives sense to weak-type expressions of the form $\mathbb{E}[X\varphi^1(x)]$ for all local test random variables X . Note that this explicit 1D argument makes no use of Malliavin calculus.

(c) *Improving on the topology:*

As stated in Theorem 5.1, our analysis provides a control on correctors in some dual Malliavin–Sobolev spaces, which quantifies how smooth and local the test random variable X should be for $\mathbb{E}[X\varphi^n]$ to be well-defined. However, we do not know whether this control is optimal, and in particular whether Malliavin calculus could be avoided. At order $\ell = \lceil \frac{d}{2} \rceil$, our analysis yields the following improved estimate: if the covariance function \mathcal{C} is compactly supported, then the stationary corrector φ^ℓ is uniquely defined in $C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$ with $\mathbb{E}[\varphi^\ell] = 0$, and it satisfies for all $x \in \mathbb{R}^d$, $R > 0$, and $q < \infty$,

$$\|\mathbb{E}[\varphi^\ell(x) \mid \mathbf{a}|_{B_R}]\|_{L^q(\Omega)} \lesssim_{\mathcal{C},q} \begin{cases} \langle R \rangle^{\frac{1}{2}} & : d \text{ odd,} \\ \log(2+R)^{\frac{1}{2}} & : d \text{ even.} \end{cases}$$

This generalizes the above explicit 1D construction (2.5), and we do not know whether such estimates extend to higher orders $n > \ell$.

(d) *Evaluating higher-order effective tensors:*

In view of the definition of weak correctors in Theorem 3, formula (1.6) now makes sense and defines higher-order effective tensors for all $1 \leq n \leq d$,

$$\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n e_{i_n} = \mathbb{E}[\mathbf{a}(\nabla \varphi_{i_1 \dots i_n}^n + \varphi_{i_1 \dots i_{n-1}}^{n-1} e_{i_n})]. \quad (2.6)$$

This formula can be substantially simplified in view of the following observation: if \mathbf{a} is pointwise symmetric, then we have for all $1 \leq n \leq d$,

$$\nabla \cdot \bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n \nabla \nabla_{i_1 \dots i_{n-1}}^{n-1} = \begin{cases} \nabla \cdot \bar{\mathbf{b}}_{i_1 \dots i_{n-1}}^n \nabla \nabla_{i_1 \dots i_{n-1}}^{n-1} & : \text{ if } n \text{ is odd,} \\ 0 & : \text{ if } n \text{ is even,} \end{cases}$$

where we have defined for all $n = 2m + 1$ with $m < \ell = \lfloor \frac{d}{2} \rfloor$,

$$e_j \cdot \bar{\mathbf{b}}_{i_1 \dots i_{n-1}}^n e_{i_n} := (-1)^{m+1} \mathbb{E} \left[\nabla \varphi_{j i_n \dots i_{m+2}}^{m+1} \cdot \mathbf{a} \nabla \varphi_{i_1 \dots i_{m+1}}^{m+1} - \varphi_{j i_n \dots i_{m+3}}^m e_{i_{m+2}} \cdot \mathbf{a} \varphi_{i_1 \dots i_m}^m e_{i_{m+1}} \right].$$

(This identity states that the tensors $\bar{\mathbf{a}}^n$ and $\bar{\mathbf{b}}^n$ coincide at least after symmetrizing their indices, which is what matters when considering associated differential operators.)

This shows that $\bar{\mathbf{a}}^n$ can be computed in terms of correctors of order $\leq \lfloor \frac{n}{2} \rfloor$ only, so that the definition of $\{\bar{\mathbf{a}}^k\}_{1 \leq k \leq d}$ only requires the use of standard “strong” correctors $\{\varphi^k\}_{1 \leq k \leq \ell}$. This identity was first noticed in our recent work [14, Remark 2.5], and also independently in [1, Theorem 3.5]. \diamond

3. PRELIMINARY: HIGHER-ORDER HOMOGENIZATION

In this section, we recall some tools and notation from the higher-order quantitative theory of stochastic homogenization. We start with large-scale regularity theory for linear elliptic equations with random coefficients [6, 3, 18]; for the purpose of this work, we focus on annealed L^p regularity as introduced in [14]. Next, we recall the higher-order intrinsic description of spatial oscillations by means of two-scale expansions [19, 7, 14].

3.1. Annealed regularity theory. Given $h \in L^2(\mathbb{R}^d; L^2(\Omega)^d)$, we consider the unique gradient solution $\nabla w_h \in L^2(\mathbb{R}^d; L^2(\Omega)^d)$ of the linear elliptic equation

$$-\nabla \cdot \mathbf{a} \nabla w_h = \nabla \cdot h, \quad \text{in } \mathbb{R}^d, \quad (3.1)$$

and we denote by \mathcal{H} the associated solution operator, or heterogeneous Helmholtz projection on $L^2(\mathbb{R}^d; L^2(\Omega)^d)$,

$$\mathcal{H}h := \nabla w_h, \quad \mathcal{H} = \nabla(-\nabla \cdot \mathbf{a} \nabla)^{-1} \nabla. \quad (3.2)$$

Due to the random character of the coefficient field \mathbf{a} , aside from the energy inequality and from Meyers’ perturbative improvements, maximal L^p regularity cannot hold in a deterministic form. Appealing to homogenization, however, the heterogeneous elliptic operator can be replaced on large scales by a homogeneous one, cf. (1.3), for which the standard constant-coefficient regularity theory is available. For this reason, the solution to (3.1) is expected to enjoy improved regularity properties on large scales. In this spirit, a quenched large-scale regularity theory was first outlined by Armstrong and Smart [6], and further developed in [5, 4, 3, 18]. The following annealed regularity result is a useful variant of this theory and was first established with Otto in [14, Section 6]: it states that maximal L^p regularity holds as in the constant-coefficient setting after taking ensemble averages (up to a tiny loss in stochastic integrability).

Theorem 3.1 (Annealed L^p regularity; [14]). *With the above notation, there holds for all $h \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^d)$, $1 < p, q < \infty$, and $\delta > 0$,*

$$\|[\mathcal{H}h]_2\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{p,q,\delta} \| [h]_2 \|_{L^p(\mathbb{R}^d; L^{q+\delta}(\Omega))}. \quad \diamond$$

3.2. Higher-order description of oscillations. Correctors are defined to correct Euclidean polynomials and make them adapted to the heterogeneous elliptic operator, and they constitute the key ingredient to describe spatial oscillations for solutions of the heterogeneous Poisson problem (1.2). The following standard result recalls the hierarchy of corrector equations. Item (i) gives sharp corrector estimates [19, 7, 14], and item (ii) states that higher-order correctors have less integrable correlations [26, 19] so that their large-scale averages have worse fluctuation scaling [14, proof of Lemma 3.3].

Theorem 3.2 (Higher-order correctors; [19, 7, 14]). *Let $d \geq 1$ and recall $\ell = \lceil \frac{d}{2} \rceil$. Higher-order correctors $\{\varphi^n\}_{0 \leq n \leq \ell}$, flux correctors $\{\sigma^n\}_{0 \leq n \leq \ell}$, fluxes $\{q^n\}_{1 \leq n \leq \ell}$, and effective tensors $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq \ell}$ are uniquely well-defined iteratively as follows:*

- $\varphi^0 := 1$ and for all $1 \leq n \leq \ell$ we define $\varphi^n := (\varphi_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\varphi_{i_1 \dots i_n}^n$ is the unique weak solution in $H_{\text{loc}}^1(\mathbb{R}^d; L^2(\Omega))$ of

$$-\nabla \cdot \mathbf{a} \nabla \varphi_{i_1 \dots i_n}^n = \nabla \cdot ((\mathbf{a} \varphi_{i_1 \dots i_{n-1}}^{n-1} - \sigma_{i_1 \dots i_{n-1}}^{n-1}) e_{i_n}), \quad \text{in } \mathbb{R}^d, \quad (3.3)$$

such that φ^n is stationary with $\mathbb{E}[\varphi^n] = 0$ for all $n < \ell$, and such that $\nabla \varphi^\ell$ is stationary with $\mathbb{E}[\nabla \varphi^\ell] = 0$ and with anchoring $\int_B \varphi^\ell = 0$.

- $\sigma^0 := 0$ and for all $1 \leq n \leq \ell$ we define $\sigma^n := (\sigma_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\sigma_{i_1 \dots i_n}^n$ is the unique weak solution in $H_{\text{loc}}^1(\mathbb{R}^d; L^2(\Omega)^{d \times d})$, with skew-symmetric matrix values, of

$$-\Delta \sigma_{i_1 \dots i_n}^n = \nabla \times q_{i_1 \dots i_n}^n, \quad \nabla \cdot \sigma_{i_1 \dots i_n}^n = q_{i_1 \dots i_n}^n, \quad \text{in } \mathbb{R}^d, \quad (3.4)$$

such that σ^n is stationary with $\mathbb{E}[\sigma^n] = 0$ for all $n < \ell$, and such that $\nabla \sigma^\ell$ is stationary with $\mathbb{E}[\nabla \sigma^\ell] = 0$ and with anchoring $\int_B \sigma^\ell = 0$.

- For all $1 \leq n \leq \ell$ we define $q^n := (q_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ with $q_{i_1 \dots i_n}^n$ given by

$$q_{i_1 \dots i_n}^n := \mathbf{a} \nabla \varphi_{i_1 \dots i_n}^n + (\mathbf{a} \varphi_{i_1 \dots i_{n-1}}^{n-1} - \sigma_{i_1 \dots i_{n-1}}^{n-1}) e_{i_n} - \bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n e_{i_n}. \quad (3.5)$$

- For all $1 \leq n \leq \ell$ we define $\bar{\mathbf{a}}^n := (\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n)_{1 \leq i_1, \dots, i_{n-1} \leq d}$ with $\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n$ the matrix given by

$$\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n e_{i_n} := \mathbb{E}[\mathbf{a} (\nabla \varphi_{i_1 \dots i_n}^n + \varphi_{i_1 \dots i_{n-1}}^{n-1} e_{i_n})], \quad (3.6)$$

which precisely ensures $\mathbb{E}[q^n] = 0$ in (3.5).

In addition, the following estimates hold:

- (i) Corrector estimates: for all $0 \leq n < \ell$ and $q < \infty$,

$$\mathbb{E}[\|\nabla \varphi^{n+1}\|_2^q]^{\frac{1}{q}} + \mathbb{E}[\|\nabla \sigma^{n+1}\|_2^q]^{\frac{1}{q}} + \mathbb{E}[\|\varphi^n\|_2^q]^{\frac{1}{q}} + \mathbb{E}[\|\sigma^n\|_2^q]^{\frac{1}{q}} \lesssim_q 1,$$

and at critical order $n = \ell$, for all $q < \infty$,

$$\mathbb{E}[\|\varphi^\ell\|_2^q(x)]^{\frac{1}{q}} + \mathbb{E}[\|\sigma^\ell\|_2^q(x)]^{\frac{1}{q}} \lesssim_q \begin{cases} \langle x \rangle^{\frac{1}{2}} & : d \text{ odd,} \\ \log(2 + |x|)^{\frac{1}{2}} & : d \text{ even.} \end{cases}$$

- (ii) Fluctuation scaling: for all $0 \leq n < \ell$, $g \in C_c^\infty(\mathbb{R}^d)$, and $q < \infty$,

$$\mathbb{E} \left[\left| \int_{\mathbb{R}^d} g(\nabla \varphi^{n+1}, \nabla \sigma^{n+1}) \right|^q \right]^{\frac{1}{q}} + \mathbb{E} \left[\left| \int_{\mathbb{R}^d} g(\varphi^n, \sigma^n) \right|^q \right]^{\frac{1}{q}} \lesssim_q \| [g]_2 \|_{L^{\frac{2d}{d+2n}}(\mathbb{R}^d)}. \quad \diamond$$

As recalled in the introduction, with these correctors at hand, we define higher-order two-scale expansions as

$$\mathcal{F}_\varepsilon^n[\bar{w}] := \sum_{k=0}^n \varepsilon^k \varphi_{i_1 \dots i_k}^k \left(\frac{\cdot}{\varepsilon}\right) \nabla_{i_1 \dots i_k}^k \bar{w}. \quad (3.7)$$

Homogenization theory states that the heterogeneous constitutive law $\nabla w_\varepsilon \mapsto \mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla w_\varepsilon$ is replaced on large scales by the effective law $\nabla \bar{w} \mapsto \bar{\mathbf{a}}^1 \nabla \bar{w}$, and dispersive corrections must be added when looking for finer accuracy,

$$\nabla \bar{w} \mapsto \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{w} := \sum_{k=1}^n \varepsilon^{k-1} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^{k-1} \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \bar{w}, \quad (3.8)$$

in terms of the higher-order effective tensors $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq \ell}$. The first-order homogenized equation (1.3) is then formally replaced by the following corrected version,

$$-\nabla \cdot \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{U}_{\varepsilon, f}^n = \nabla \cdot f, \quad \text{in } \mathbb{R}^d. \quad (3.9)$$

This corrected equation is however ill-posed in general for $n \geq 2$, as the symbol may be indefinite, and we shall use the following well-defined proxy.

Definition 3.3 (Higher-order homogenized equations). *For all $1 \leq n \leq \ell$, the n th-order homogenized solution operator is defined as follows for all $f \in C_c^\infty(\mathbb{R}^d)^d$,*

$$\nabla \bar{U}_\varepsilon^n[f] := \sum_{k=1}^n \varepsilon^{k-1} \nabla \tilde{u}_f^k,$$

where $\nabla \tilde{u}_f^1 := \nabla \bar{u}_f^1$ is the unique gradient solution in $L^2(\mathbb{R}^d)^d$ of the first-order homogenized equation,

$$-\nabla \cdot \bar{\mathbf{a}}^1 \nabla \bar{u}_f^1 = \nabla \cdot f, \quad \text{in } \mathbb{R}^d,$$

and where for $2 \leq n \leq \ell$ we inductively define the n th-order correction $\nabla \tilde{u}_f^n$ as the unique gradient solution in $L^2(\mathbb{R}^d)^d$ of

$$-\nabla \cdot \bar{\mathbf{a}}^1 \nabla \tilde{u}_f^n = \nabla \cdot \sum_{k=2}^n \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^{n+1-k}, \quad \text{in } \mathbb{R}^d. \quad \diamond$$

This function $\nabla \bar{U}_\varepsilon^n[f]$ is indeed viewed as a proxy for a solution $\nabla \bar{U}_{\varepsilon, f}^n$ of the n th-order homogenized equation (3.9) since the following identity holds,

$$-\nabla \cdot \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{U}_\varepsilon^n[f] = \nabla \cdot f - \nabla \cdot \left(\sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \right),$$

where the second right-hand side term is an error term of order $O(\varepsilon^n)$. With these definitions, using the algebraic structure of corrector equations together with corrector estimates, the accuracy of the higher-order two-scale expansion is easily justified, cf. [19, Theorem 1.6] and [14, Proposition 2.7]. Due to the maximal number $\ell = \lceil \frac{d}{2} \rceil$ of well-behaved correctors, the accuracy saturates at the optimal order $O(\varepsilon^{d/2})$. Optimality is confirmed as random fluctuations become dominant at that order, see [13, 14].

Proposition 3.4 ([19, 14]). *For all $1 \leq n \leq \ell$ and $f \in C_c^\infty(\mathbb{R}^d)^d$, the n th-order two-scale expansion error satisfies*

$$\begin{aligned} -\nabla \cdot \mathbf{a}(\cdot) \nabla (u_{\varepsilon, f} - \mathcal{F}_\varepsilon^n[\bar{\mathcal{U}}_\varepsilon^n[f]]) &= \nabla \cdot \left(\sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \right) \\ &\quad + \nabla \cdot \left(\varepsilon^n (\mathbf{a} \varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n) (\cdot) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_\varepsilon^n[f] \right), \end{aligned} \quad (3.10)$$

which entails for all $1 < p, q < \infty$,

$$\| [\nabla u_{\varepsilon, f} - \nabla \mathcal{F}_\varepsilon^\ell[\bar{\mathcal{U}}_\varepsilon^\ell[f]]] \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{f, p, q} \varepsilon^{\frac{d}{2}} \times \begin{cases} 1 & : d \text{ odd}, \\ |\log \varepsilon|^{\frac{1}{2}} & : d \text{ even}. \end{cases} \quad \diamond$$

4. FUNCTIONAL SETUP

In this section, we introduce some functional tools and notation. We start with a reminder on stationarity and on the associated stationary differential calculus on the probability space, as is commonly used in stochastic homogenization [29, 22]. Next, we turn to a brief reminder on Malliavin calculus, and we introduce a new scale of “refined” Malliavin–Sobolev spaces. While stationary calculus is generated by spatial translations of stationary fields, Malliavin calculus is generated by changes in the underlying white noise.

4.1. Stationary calculus. In the present Gaussian setting (2.1)–(2.2), recall that the probability space (Ω, \mathbb{P}) is chosen to be generated by the underlying white noise ξ . Translations $x \mapsto \xi(\cdot + x)$ then induce an action $T = \{T_x\}_{x \in \mathbb{R}^d}$ of the additive group $(\mathbb{R}^d, +)$ on the space of random variables. In particular, for $X \in \mathcal{S}(\Omega)$, say $X = g(\int_{\mathbb{R}^d} h_1 \cdot \xi, \dots, \int_{\mathbb{R}^d} h_n \cdot \xi)$, we define

$$T_x X = g\left(\int_{\mathbb{R}^d} h_1 \cdot \xi(\cdot + x), \dots, \int_{\mathbb{R}^d} h_n \cdot \xi(\cdot + x)\right).$$

As the law of $\xi(\cdot + x)$ does not depend on the shift $x \in \mathbb{R}^d$, the map T_x is an isometry on $L^q(\Omega)$ for all q . Also note that, for any random variable X , the map $(x, \omega) \mapsto (T_x X)(\omega)$ is stochastically continuous and jointly measurable on $\mathbb{R}^d \times \Omega$.

In this setting, stationarity is defined as follows: a random field $\psi \in L_{\text{loc}}^1(\mathbb{R}^d; L^1(\Omega))$ is stationary if there exists a random variable $X \in L^1(\Omega)$ such that $\psi(x, \omega) = (T_x X)(\omega)$. In other words, a random field is stationary if it is covariant under spatial translations: spatial translations of the field amount to spatial translations of the underlying Gaussian field. For all q , this provides a canonical isomorphism between random variables in $L^q(\Omega)$ and stationary random fields in $L_{\text{loc}}^q(\mathbb{R}^d; L^q(\Omega))$: for $X \in L^q(\Omega)$ we define its stationary extension $X^\sharp(x, \omega) = (T_x X)(\omega)$, and for a stationary field $\psi \in L_{\text{loc}}^q(\mathbb{R}^d; L^q(\Omega))$ we define $\psi^\flat \in L^q(\Omega)$ such that $\psi(x, \omega) = (T_x \psi^\flat)(\omega)$. Clearly, $(\psi^\flat)^\sharp = \psi$ and $(X^\sharp)^\flat = X$.

A differential calculus on $L^q(\Omega)$ is naturally developed via this stationarity. Indeed, for all $q < \infty$, the action $T = \{T_x\}_{x \in \mathbb{R}^d}$ is easily checked to constitute a $(d$ -parameter) C_0 -group of isometries on $L^q(\Omega)$, and we then define the *stationary gradient* ∇^{st} as the generator of this group, which is a densely defined skew-adjoint operator on $L^q(\Omega)$. Its domain is denoted by $W^{1,q}(\Omega)$ and obviously contains $\mathcal{S}(\Omega)$. For all $s \geq 0$, we similarly define the higher-order Sobolev space $W^{s,q}(\Omega)$ as the domain of $(-\nabla^{\text{st}} \cdot \nabla^{\text{st}})^{s/2}$ in $L^q(\Omega)$. For $q = 2$, we write $H^s(\Omega) = W^{s,2}(\Omega)$.

This stationary gradient can be reinterpreted via the isomorphism between random variables and stationary random fields: as the space of stationary fields in $L^q_{\text{loc}}(\mathbb{R}^d; L^q(\Omega))$ is identified with $L^q(\Omega)$, the spatial gradient ∇ on locally L^q integrable functions turns into a densely defined linear operator on $L^q(\Omega)$, which is easily checked to coincide with the stationary gradient ∇^{st} . More precisely, the space $W^{1,q}(\Omega)$ coincides with the space of stationary fields in $W^{1,q}_{\text{loc}}(\mathbb{R}^d; L^q(\Omega))$, and there holds for all $X \in W^{1,q}(\Omega)$,

$$(\nabla^{\text{st}} X)^{\sharp} = \nabla X^{\sharp}.$$

In the context of corrector equations, cf. Theorem 3.2, we often consider linear elliptic equations of the following form, given a stationary field $\psi \in L^2_{\text{loc}}(\mathbb{R}^d; L^2(\Omega)^d)$,

$$-\nabla \cdot \mathbf{a} \nabla w_{\psi} = \nabla \cdot \psi, \quad \text{in } \mathbb{R}^d.$$

The gradient solution ∇w_{ψ} can be uniquely chosen as a stationary field in $L^2_{\text{loc}}(\mathbb{R}^d; L^2(\Omega)^d)$ with $\mathbb{E}[\nabla w_{\psi}] = 0$ (see e.g. [29, proof of Theorem 2]), but w_{ψ} itself cannot be chosen stationary in general due to the absence of a Poincaré inequality in this stationary setting — which is in turn the very reason why there exists only a finite number of stationary correctors. We shall abusively use the same notation as in (3.2),

$$\mathcal{H}\psi = \nabla w_{\psi}, \tag{4.1}$$

for the solution operator, or heterogeneous Helmholtz projection, in this stationary setting. In terms of stationary calculus, the above equation is equivalently written as follows,²

$$\nabla w_{\psi} = X_{\psi}^{\sharp}, \quad -\nabla^{\text{st}} \cdot (\mathbf{a}^{\flat} X_{\psi}) = \nabla^{\text{st}} \cdot \psi^{\flat}, \quad \nabla^{\text{st}} \times X_{\psi} = 0, \quad \mathbb{E}[X_{\psi}] = 0,$$

and we set $\mathcal{H}^{\text{st}} \psi^{\flat} := (\mathcal{H}\psi)^{\flat} = X_{\psi}$ on $L^2(\Omega)^d$. In parallel with the annealed L^p regularity in Theorem 3.1, we prove a corresponding regularity result for this stationary Helmholtz projection in $L^q(\Omega)$ up to a tiny loss of stochastic integrability.

Corollary 4.1 (Stationary L^q regularity). *With the above notation, there holds for all $X \in L^{\infty}(\Omega)$, $1 < q < \infty$, and $\delta > 0$,*

$$\|[\mathcal{H}X^{\sharp}]_2^{\flat}\|_{L^q(\Omega)} \lesssim_{q,\delta} \| [X^{\sharp}]_2^{\flat} \|_{L^{q+\delta}(\Omega)},$$

where we recall the notation $[X^{\sharp}]_2^{\flat} = (f_B |X^{\sharp}|^2)^{1/2}$ for the local quadratic average. \diamond

Proof. By a duality argument, it suffices to consider the case $2 \leq q < \infty$. Appealing to quenched large-scale regularity in form of [7, Lemma 16], and recalling that in the present Gaussian setting with integrable correlations the so-called minimal radius r_* in [7] satisfies $\mathbb{E}[(r_*)^{dq}] \leq (Cq)^q$ for all $q < \infty$, see [18, Theorem 3], we obtain for all $2 \leq q < d$ and $\alpha > 1$, setting $q_* := \frac{dq}{d-q}$,

$$\|[\mathcal{H}X^{\sharp}]_2^{\flat}\|_{L^{\frac{q_*}{\alpha}}(\Omega)} \lesssim q_*^3 \left(\frac{q_*}{\alpha-1}\right)^{\frac{1}{2}} \left(\| [X^{\sharp}]_2^{\flat} \|_{L^{q_*}(\Omega)} + \|[\mathcal{H}X^{\sharp}]_2^{\flat}\|_{L^q(\Omega)} \right).$$

Iterating this inequality on $\mathcal{H}X^{\sharp}$ and starting with the energy estimate

$$\|[\mathcal{H}X^{\sharp}]_2^{\flat}\|_{L^2(\Omega)} = \|\mathcal{H}^{\text{st}} X\|_{L^2(\Omega)} \lesssim \|X\|_{L^2(\Omega)} = \| [X^{\sharp}]_2^{\flat} \|_{L^2(\Omega)},$$

the conclusion follows. \square

²These equations are understood in the following weak sense: there holds $\mathbb{E}[\nabla^{\text{st}} Y \cdot (\mathbf{a}^{\flat} X_{\psi} + \psi^{\flat})] = 0$ for all $Y \in H^1(\Omega)$, and the random vector X_{ψ} belongs to the closure of the space of stationary gradients, that is, $\mathbb{E}[X_{\psi}] = 0$ and $\mathbb{E}[(\nabla_j^{\text{st}} Y)(X_{\psi})_k] = \mathbb{E}[(\nabla_k^{\text{st}} Y)(X_{\psi})_j]$ for all j, k and all $Y \in H^1(\Omega)$.

4.2. Malliavin calculus. We recall some classical notation and tools from Malliavin calculus; we refer to [28, 27] for details. The underlying white noise ξ in (2.2) is viewed as a random Schwartz distribution: by definition, the collection of linear random variables

$$\xi(h) := \int_{\mathbb{R}^d} h \cdot \xi, \quad h \in C_c^\infty(\mathbb{R}^d)^\kappa,$$

are jointly Gaussian random variables with expectation $\mathbb{E}[\xi(h)] = 0$ and with covariance

$$\langle \xi(h'), \xi(h) \rangle_{L^2(\Omega)} = \mathbb{E}[\xi(h')\xi(h)] = \int_{\mathbb{R}^d} h' \cdot h = \langle h', h \rangle_{L^2(\mathbb{R}^d)}.$$

By density, we may define $\xi(h) \in L^2(\Omega)$ for all $h \in L^2(\mathbb{R}^d)^\kappa$, which provides an embedding $L^2(\mathbb{R}^d)^\kappa \hookrightarrow L^2(\Omega)$. Similarly as $C_c^\infty(\mathbb{R}^d)^\kappa$ is viewed as a model dense subspace of $L^2(\mathbb{R}^d)^\kappa$, we recall the following model subspace of “smooth and local” random variables, cf. (2.4),

$$\mathcal{S}(\Omega) := \left\{ g(\xi(h_1), \dots, \xi(h_n)) : n \in \mathbb{N}, g \in C_b^\infty(\mathbb{R}^n), h_1, \dots, h_n \in C_c^\infty(\mathbb{R}^d)^\kappa \right\}.$$

For all $q < \infty$, as this subspace $\mathcal{S}(\Omega)$ is dense in $L^q(\Omega)$, it allows to define operators and prove properties on this simpler subspace before extending them to $L^q(\Omega)$.

For a random variable $X \in \mathcal{S}(\Omega)$, say $X = g(\xi(h_1), \dots, \xi(h_n))$, we define its Malliavin derivative $DX \in L^q(\Omega; L^2(\mathbb{R}^d)^\kappa)$ as

$$DX := \sum_{j=1}^n h_j (\partial_j g)(\xi(h_1), \dots, \xi(h_n)), \quad (4.2)$$

and we use the short-hand notation $D_x X := (DX)(x) \in L^q(\Omega)$. Similarly, for all $k \geq 1$, the k th Malliavin derivative $D^k X \in L^q(\Omega; (L^2(\mathbb{R}^d)^\kappa)^{\otimes k})$ is given by

$$D^k X := \sum_{1 \leq j_1, \dots, j_k \leq n} (h_{j_1} \otimes \dots \otimes h_{j_k}) (\partial_{j_1 \dots j_k}^k g)(\xi(h_1), \dots, \xi(h_n)).$$

This operator $D^k : \mathcal{S}(\Omega) \subset L^q(\Omega) \rightarrow L^q(\Omega; (L^2(\mathbb{R}^d)^\kappa)^{\otimes k})$ is easily checked to be closable on $L^q(\Omega)$ for any $q < \infty$, and we still denote by D^k its closure. We also set

$$\|X\|_{\mathbb{D}^{k,q}(\Omega)}^q := \|X\|_{L^q(\Omega)}^q + \sum_{j=1}^k \|D^j X\|_{L^q(\Omega; \mathfrak{H}^{\otimes j})}^q, \quad (4.3)$$

and we define the Malliavin–Sobolev space $\mathbb{D}^{k,q}(\Omega)$ as the closure of $\mathcal{S}(\Omega) \subset L^q(\Omega)$ for this norm. This is also sometimes referred to as the Watanabe–Sobolev space.

Next, we define the divergence operator D^* as the adjoint of the Malliavin derivative D , and we construct the corresponding Ornstein–Uhlenbeck operator

$$\mathcal{L} := D^* D.$$

This operator $\mathcal{L} : \mathcal{S}(\Omega) \subset L^q(\Omega) \rightarrow L^q(\Omega)$ is well-defined, symmetric, and closable. Its closure is still denoted by \mathcal{L} and is a non-negative operator with domain $\mathbb{D}^{2,q}(\Omega)$. For $q = 2$, it is also self-adjoint. In the sequel, we will often let the operator D act on functions $w \in L_{\text{loc}}^1(\mathbb{R}^d; \mathbb{D}^{1,2}(\Omega))$ by freezing the space variable, that is, $(D_z w)(x) := D_z(w(x))$, and similarly for \mathcal{L} . The following lemma collects some useful properties; although standard, a short proof is included for the reader’s convenience.

Lemma 4.2.

- (i) The Ornstein–Uhlenbeck operator \mathcal{L} commutes with stationarity, in the sense that $\mathcal{L}T_x X = T_x \mathcal{L}X$ for all $x \in \mathbb{R}^d$ and $X \in \mathcal{S}(\Omega)$.
- (ii) For all q , the operator $(1 + \mathcal{L})^{-1}$ is a contraction on $L^q(\Omega)$. Moreover, for all $w \in C_c^\infty(\mathbb{R}^d; \mathcal{R}(\Omega))$ and all p, q , we have $\|[(1 + \mathcal{L})^{-1}w]_p\|_{L^q(\Omega)} \leq \|w\|_{L^q(\Omega)}$.
- (iii) For all $X, Y \in \mathbb{D}^{1,2}(\Omega)$, the following Helffer–Sjöstrand identity holds

$$\text{Cov}[Y; X] = \mathbb{E} \left[\langle DY, (1 + \mathcal{L})^{-1}DX \rangle_{L^2(\mathbb{R}^d)} \right],$$

which implies in particular Poincaré’s inequality

$$\text{Var}[X] \leq \|DX\|_{L^2(\Omega; L^2(\mathbb{R}^d))}^2. \quad \diamond$$

Proof. We consider the Ornstein–Uhlenbeck semigroup $\{e^{-t\mathcal{L}}\}_{t \geq 0}$ and we recall Mehler’s formula: choosing an iid copy ξ' of the white noise ξ , and denoting by $\mathbb{E}'[\cdot]$ the expectation with respect to ξ' , there holds for all $t \geq 0$ and $X \in \mathcal{S}(\Omega)$,

$$e^{-t\mathcal{L}}X = \mathbb{E}'[X_t], \quad (4.4)$$

where X_t is obtained by replacing ξ with its interpolation $\xi_t := e^{-t}\xi + \sqrt{1 - e^{-2t}}\xi'$. This formula is easily justified by explicitly computing the generator of both sides on $\mathcal{S}(\Omega)$; see e.g. [27, Theorem 2.8.2]. Based on this formula, there obviously holds $e^{-t\mathcal{L}}T_x X = T_x e^{-t\mathcal{L}}X$, which proves (i).

We turn to the proof of (ii). Writing $(1 + \mathcal{L})^{-1}X = \int_0^\infty e^{-t}e^{-t\mathcal{L}}X dt$, and using that ξ_t has the same law as ξ , which entails

$$\|e^{-t\mathcal{L}}X\|_{L^q(\Omega)} = \|\mathbb{E}'[X_t]\|_{L^q(\Omega)} \leq \|X\|_{L^q(\Omega)}, \quad (4.5)$$

we find

$$\|(1 + \mathcal{L})^{-1}X\|_{L^q(\Omega)} \leq \int_0^\infty e^{-t} \|\mathbb{E}'[X_t]\|_{L^q(\Omega)} dt \leq \|X\|_{L^q(\Omega)}.$$

The same computation holds when inserting local averages, and the conclusion (ii) follows.

It remains to prove (iii), and we start with Poincaré’s inequality. As Mehler’s formula (4.4) yields $e^{-t\mathcal{L}}X \rightarrow \mathbb{E}[X]$ in $L^2(\Omega)$ as $t \uparrow \infty$, we can write

$$\text{Var}[X] = - \int_0^\infty \partial_t \mathbb{E}[(e^{-t\mathcal{L}}X)^2] dt = 2 \int_0^\infty \mathbb{E}[(e^{-t\mathcal{L}}X)\mathcal{L}(e^{-t\mathcal{L}}X)] dt,$$

and thus, integrating by parts with $\mathcal{L} = D^*D$,

$$\text{Var}[X] = 2 \int_0^\infty \|D(e^{-t\mathcal{L}}X)\|_{L^2(\mathbb{R}^d; L^2(\Omega))}^2 dt.$$

Noting that Mehler’s formula (4.4) yields

$$De^{-t\mathcal{L}}X = e^{-t}e^{-t\mathcal{L}}DX, \quad (4.6)$$

and using (4.5) again, Poincaré’s inequality follows,

$$\text{Var}[X] = 2 \int_0^\infty e^{-2t} \|e^{-t\mathcal{L}}DX\|_{L^2(\mathbb{R}^d; L^2(\Omega))}^2 dt \leq \|DX\|_{L^2(\mathbb{R}^d; L^2(\Omega))}^2.$$

We now turn to the Helffer–Sjöstrand identity. Let $X, Y \in \mathcal{S}(\Omega)$ with $\mathbb{E}[X] = \mathbb{E}[Y] = 0$. Poincaré’s inequality together with the Lax–Milgram theorem ensures that there exists $Z \in \mathbb{D}^{2,2}(\Omega)$ with $X = \mathcal{L}Z$. Integrating by parts with $\mathcal{L} = D^*D$, we then find

$$\text{Cov}[Y; X] = \mathbb{E}[YX] = \mathbb{E}[Y(\mathcal{L}Z)] = \mathbb{E}[\langle DY, DZ \rangle_{L^2(\mathbb{R}^d)}].$$

Noting that identity (4.6) entails $D\mathcal{L}Z = (1 + \mathcal{L})DZ$, we deduce $DZ = (1 + \mathcal{L})^{-1}DX$, and the conclusion follows. \square

4.3. “Refined” Malliavin–Sobolev spaces. We define the following “refined” version of Malliavin–Sobolev spaces, which play a key role in the sequel of this work and are a variant of spaces first introduced in [2]: for $1 \leq p, q < \infty$ and integer $k \geq 0$, we set

$$\|X\|_{\mathbb{M}_{p,q}^k(\Omega)} := \|[X^\sharp]_2\|_{L^q(\Omega)} + \sum_{j=1}^k \|[D_0^j X^\sharp]_2\|_{L^p((\mathbb{R}^d)^j; L^q(\Omega))}, \quad (4.7)$$

and we define the refined Malliavin–Sobolev space $\mathbb{M}_{p,q}^k(\Omega)$ as the closure of $\mathcal{S}(\Omega)$ for this norm, which is easily checked to define a separable Banach space. By stationarity, note that the norm $\|[D_x^j X^\sharp]_2\|_{L^p((\mathbb{R}^d)^j; L^q(\Omega))}$ is independent of $x \in \mathbb{R}^d$ (see the proof below), and we have simply chosen $x = 0$ in the definition of the norm. We refer to Remark 4.5 for the motivation of this precise definition and for the comparison with the slightly different choice in [2]. We start with a few general properties.

Lemma 4.3.

- (i) For $p = q = 2$, the Banach spaces $\mathbb{M}_{2,2}^k(\Omega)$ and $\mathbb{D}^{k,2}(\Omega)$ are isomorphic.
- (ii) The Banach space $\mathbb{M}_{p,q}^k(\Omega)$ embeds into $\mathbb{M}_{r,s}^l(\Omega)$ whenever $k \geq l$, $p \leq r$, $q \geq s$.
- (iii) For all $X, Y \in \mathcal{S}(\Omega)$ and $1 \leq p, q \leq \infty$,

$$\text{Cov}[Y; X] \leq \|[D_0 Y^\sharp]_2\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} \|[D_0 X^\sharp]_2\|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

- (iv) The following Poincaré inequality holds for all $X \in \mathcal{S}(\Omega)$ and $2 \leq q < \infty$,

$$\|[(X - \mathbb{E}[X])^\sharp]_2\|_{L^q(\Omega)} \lesssim q^{\frac{1}{2}} \|[D_0 X^\sharp]_2\|_{L^2(\mathbb{R}^d; L^q(\Omega))}. \quad \diamond$$

Proof. We start with the following observation: the definition (4.2) of the Malliavin derivative and the definition of stationary extensions lead to

$$T_x D_y X = D_{y+x} T_x X, \quad (4.8)$$

and thus, as T_x is an isometry on $L^q(\Omega)$,

$$\begin{aligned} \|D_0 X^\sharp\|_{L^p(\mathbb{R}^d; L^q(\Omega))}^p &= \int_{\mathbb{R}^d} \mathbb{E}[|D_0 T_x X|^q]^{\frac{p}{q}} dx = \int_{\mathbb{R}^d} \mathbb{E}[|T_x D_{-x} X|^q]^{\frac{p}{q}} dx \\ &= \int_{\mathbb{R}^d} \mathbb{E}[|D_{-x} X|^q]^{\frac{p}{q}} dx = \|DX\|_{L^p(\mathbb{R}^d; L^q(\Omega))}^p. \end{aligned} \quad (4.9)$$

We turn to the proof of (i). For $q = 2$, local quadratic averages can be removed by stationarity and we find

$$\|X\|_{\mathbb{M}_{p,2}^k(\Omega)} = \|X\|_{L^2(\Omega)} + \sum_{j=1}^k \|D_0^j X^\sharp\|_{L^p((\mathbb{R}^d)^j; L^2(\Omega))}.$$

Combined with (4.9), this becomes

$$\|X\|_{\mathbb{M}_{p,2}^k(\Omega)} = \|X\|_{L^2(\Omega)} + \sum_{j=1}^k \|D^j X\|_{L^p((\mathbb{R}^d)^j; L^2(\Omega))}.$$

In particular, for $p = 2$, we deduce that the norms of $\mathbb{M}_{2,2}^k(\Omega)$ and $\mathbb{D}^{k,2}(\Omega)$ are Lipschitz-equivalent, which proves (i).

Next, for $p \leq r$ and $q \geq s$, item (ii) follows Jensen's inequality $L^q(\Omega) \subset L^s(\Omega)$, together with the following discrete $\ell^p - \ell^r$ inequality, taking advantage of local averages,

$$\begin{aligned} \|[D_0^j X^\sharp]_2\|_{L^p((\mathbb{R}^d)^j; L^q(\Omega))} &\simeq \left(\sum_{z \in \frac{1}{C}\mathbb{Z}^d} \|[D_0^j X^\sharp]_2(z)\|_{L^q(\Omega)}^p \right)^{\frac{1}{p}} \\ &\lesssim \left(\sum_{z \in \frac{1}{C}\mathbb{Z}^d} \|[D_0^j X^\sharp]_2(z)\|_{L^q(\Omega)}^r \right)^{\frac{1}{r}} \simeq \|[D_0^j X^\sharp]_2\|_{L^r((\mathbb{R}^d)^j; L^q(\Omega))}. \end{aligned} \quad (4.10)$$

We turn to the proof of (iii). Starting point is the Helffer–Sjöstrand formula in Lemma 4.2. Using (4.8), recalling that \mathcal{L} commutes with stationarity in the sense of Lemma 4.2(i), and that T is a group of isometries, it becomes

$$\begin{aligned} \text{Cov}[Y; X] &= \mathbb{E} \left[\int_{\mathbb{R}^d} (D_x Y) (1 + \mathcal{L})^{-1} (D_x X) dx \right] \\ &= \mathbb{E} \left[\int_{\mathbb{R}^d} (T_{-x} D_x Y) (1 + \mathcal{L})^{-1} (T_{-x} D_x X) dx \right] \\ &= \mathbb{E} \left[\int_{\mathbb{R}^d} (D_0 Y^\sharp)(x) (1 + \mathcal{L})^{-1} (D_0 X^\sharp)(x) dx \right], \end{aligned}$$

and thus, by Hölder's inequality,

$$\text{Cov}[Y; X] \leq \|[D_0 Y^\sharp]_2\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} \|[(1 + \mathcal{L})^{-1} D_0 X^\sharp]_2\|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

Appealing to Lemma 4.2(ii) then yields the conclusion (iii).

We turn to the proof of (iv). Note that some care is needed to deal with local quadratic averages. Let $X \in \mathcal{S}(\Omega)$ with $\mathbb{E}[X] = 0$. Starting point is the decomposition

$$\|[X^\sharp]_2\|_{L^q(\Omega)}^q = \mathbb{E}[(X^\sharp]_2)^{q-1}] \mathbb{E}[(X^\sharp]_2) + \text{Cov}[(X^\sharp]_2)^{q-1}; (X^\sharp]_2), \quad (4.11)$$

and we separately analyze the two right-hand side terms. On the one hand, using Jensen's inequality and Poincaré's inequality of Lemma 4.2(iii), we find

$$\begin{aligned} \mathbb{E}[(X^\sharp]_2)^{q-1}] \mathbb{E}[(X^\sharp]_2) &\leq \|[X^\sharp]_2\|_{L^q(\Omega)}^{q-1} \|X\|_{L^2(\Omega)} \\ &\leq \|[X^\sharp]_2\|_{L^q(\Omega)}^{q-1} \|DX\|_{L^2(\Omega; L^2(\mathbb{R}^d))}, \end{aligned}$$

and thus, as in the proof of (i),

$$\mathbb{E}[(X^\sharp]_2)^{q-1}] \mathbb{E}[(X^\sharp]_2) \leq \|[X^\sharp]_2\|_{L^q(\Omega)}^{q-1} \|[D_0 X^\sharp]_2\|_{L^2(\mathbb{R}^d; L^2(\Omega))}. \quad (4.12)$$

On the other hand, the Helffer–Sjöstrand identity of Lemma 4.2(iii) together with identity (4.8) yields

$$\begin{aligned} \text{Cov} \left[([X^\sharp]_2^b)^{q-1}; [X^\sharp]_2^b \right] &= \mathbb{E} \left[\int_{\mathbb{R}^d} D_y \left(([X^\sharp]_2^b)^{q-1} \right) (1 + \mathcal{L})^{-1} D_y \left([X^\sharp]_2^b \right) dy \right] \\ &= \mathbb{E} \left[\int_{\mathbb{R}^d} D_0 \left(([X^\sharp]_2)^{q-1} \right) (1 + \mathcal{L})^{-1} D_0 \left([X^\sharp]_2 \right) \right], \end{aligned}$$

and thus, computing $D_0 \left(([X^\sharp]_2)^{q-1} \right) = (q-1) ([X^\sharp]_2)^{q-2} D_0 \left([X^\sharp]_2 \right)$, using Hölder's inequality, appealing to Lemma 4.2(ii), and noting that $|D_0 \left([X^\sharp]_2 \right)| \leq [D_0 X^\sharp]_2$, we find

$$\text{Cov} \left[([X^\sharp]_2^b)^{q-1}; [X^\sharp]_2^b \right] \leq (q-1) \| [X^\sharp]_2^b \|_{L^q(\Omega)}^{q-2} \| [D_0 X^\sharp]_2 \|_{L^2(\mathbb{R}^d; L^q(\Omega))}^2.$$

Inserting this into (4.11) together with (4.12), the claim (iv) follows. \square

Next, as a consequence of Theorem 3.1 and Corollary 4.1, we establish a corresponding regularity estimate for the stationary Helmholtz projection \mathcal{H}^{st} in Malliavin–Sobolev spaces; recall the definition (4.1) and $(\mathcal{H}^{\text{st}} X)^\sharp = \mathcal{H} X^\sharp$.

Corollary 4.4 (Stationary $\mathbb{M}_{p,q}^1$ regularity). *For all $X \in \mathcal{S}(\Omega)$, $1 < p, q < \infty$, and $\delta > 0$,*

$$\| \mathcal{H}^{\text{st}} X \|_{\mathbb{M}_{p,q}^1(\Omega)} \lesssim_{p,q,\delta} \| X \|_{\mathbb{M}_{p,q+\delta}^1(\Omega)}. \quad \diamond$$

Proof. Let $X \in \mathcal{S}(\Omega)$, $1 < p, q < \infty$, and $\delta > 0$ be fixed. Appealing to the stationary L^q regularity estimate of Corollary 4.1, we find

$$\| [(\mathcal{H}^{\text{st}} X)^\sharp]_2 \|_{L^q(\Omega)} \lesssim_{q,\delta} \| [X^\sharp]_2 \|_{L^{q+\delta}(\Omega)}, \quad (4.13)$$

and it remains to show

$$\| [D_0(\mathcal{H}^{\text{st}} X)^\sharp]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{q,\delta} \| X \|_{\mathbb{M}_{p,q+\delta}^1(\Omega)}. \quad (4.14)$$

By definition, $(\mathcal{H}^{\text{st}} X)^\sharp = \nabla w_X$ is the unique stationary gradient solution in $L_{\text{loc}}^2(\mathbb{R}^d; L^2(\Omega))$ of

$$-\nabla \cdot \mathbf{a} \nabla w_X = \nabla \cdot X^\sharp, \quad \text{in } \mathbb{R}^d,$$

with $\mathbb{E} [\nabla w_X] = 0$, and we deduce the following relation by taking the Malliavin derivative in this equation,

$$-\nabla \cdot \mathbf{a} \nabla D_0 w_X = \nabla \cdot D_0 X^\sharp + \nabla \cdot (D_0 \mathbf{a}) \nabla w_X.$$

The annealed L^p regularity estimate of Theorem 3.1 then gives

$$\begin{aligned} \| [D_0(\mathcal{H}^{\text{st}} X)^\sharp]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} &= \| [\nabla D_0 w_X]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \\ &\lesssim_{p,q,\delta} \| [D_0 X^\sharp]_2 \|_{L^p(\mathbb{R}^d; L^{q+\delta}(\Omega))} + \| [(D_0 \mathbf{a}) \nabla w_X]_2 \|_{L^p(\mathbb{R}^d; L^{q+\delta}(\Omega))}. \end{aligned}$$

In view of (2.1)–(2.2), we find $|D_0 \mathbf{a}(x)| \lesssim |\mathcal{C}_0(x)|$, and the integrability condition (2.3) then allows to bound the above by

$$\| [D_0(\mathcal{H}^{\text{st}} X)^\sharp]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{p,q,\delta} \| [D_0 X^\sharp]_2 \|_{L^p(\mathbb{R}^d; L^{q+\delta}(\Omega))} + \| [(\mathcal{H}^{\text{st}} X)^\sharp]_2 \|_{L^{q+\delta}(\Omega)}. \quad (4.15)$$

Combining this with (4.13), and recalling the definition (4.7) of the norm of $\mathbb{M}_{p,q}^1(\Omega)$, the claim (4.14) follows. \square

Remark 4.5. We comment on our definition of refined Malliavin–Sobolev spaces and we compare it with the slightly different choice in [2]. Removing local quadratic averages in the definition (4.7), and using identity (4.9), the norm of $\mathbb{M}_{p,q}^k(\Omega)$ would be reduced to the following,

$$\|X\|_{L^q(\Omega)} + \sum_{j=1}^k \|D^j X\|_{L^p((\mathbb{R}^d)^j; L^q(\Omega))},$$

which would then coincide with the choice in [2] provided that $L^p((\mathbb{R}^d)^j; L^q(\Omega))$ is further replaced by $L^q(\Omega; L^p((\mathbb{R}^d)^j))$. This makes two differences with [2]: we add local quadratic averages and we reverse spatial and probabilistic norms. This choice is designed to ensure the validity of the above stationary regularity result for \mathcal{H}^{st} , cf. Corollary 4.4. \diamond

4.4. Schwartz distributions on the probability space. We denote by $\mathcal{S}'(\Omega)$ the space of continuous linear functionals on $\mathcal{S}(\Omega)$, which is viewed as a space of Schwartz-like distributions on the probability space.³ It is easily checked that $\mathcal{S}'(\Omega)$, endowed with the weak-* topology, is separable and that $\mathcal{S}(\Omega)$ embeds as a dense linear subspace. For $X \in \mathcal{S}'(\Omega)$, the stationary extension $X^\sharp \in C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$ and the stationary gradient $\nabla^{\text{st}} X \in \mathcal{S}'(\Omega)$ are defined by duality via the following relations for all $Y \in \mathcal{S}(\Omega)$,

$$\begin{aligned} \langle X^\sharp(x), Y \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)} &:= \langle X, Y^\sharp(-x) \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)}, \\ \langle \nabla^{\text{st}} X, Y \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)} &:= -\langle X, \nabla^{\text{st}} Y \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)}. \end{aligned}$$

Conversely, an element $\psi \in C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$ is said to be stationary if it coincides with the stationary extension of an element $\psi^b \in \mathcal{S}'(\Omega)$. For $X \in \mathcal{S}'(\Omega)$ and $Y \in \mathcal{S}(\Omega)$, we use for simplicity the abusive notation $\mathbb{E}[XY] = \langle X, Y \rangle_{\mathcal{S}'(\Omega), \mathcal{S}(\Omega)}$ for the duality product.

As for usual Schwartz distributions, we may refine the space $\mathcal{S}'(\Omega)$ by considering dual Malliavin–Sobolev spaces: we denote by $(\mathbb{M}_{p,q}^k)'(\Omega) \subset \mathcal{S}'(\Omega)$ the dual space of $\mathbb{M}_{p,q}^k(\Omega)$, that is, the space of continuous linear functionals on $\mathbb{M}_{p,q}^k(\Omega)$. For all $1 \leq p, q < \infty$ and integers $k \geq 0$, the space $(\mathbb{M}_{p,q}^k)'(\Omega)$ is weakly-* separable and $\mathcal{S}(\Omega)$ embeds as a dense linear subspace. Such dual spaces were already used in [2], and might also be compared to some extent to Kondratiev and Hida spaces, see e.g. [21].

5. WEAK CORRECTOR THEORY

This section is devoted to the proof of Theorem 3: while only the first $\ell = \lceil \frac{d}{2} \rceil$ correctors can be constructed as stationary fields with bounded moments, see Theorem 3.2, we show that twice as many stationary correctors can be constructed in a distributional sense, which is expected to be optimal and is the cornerstone of our approach to the Bourgain–Spencer conjecture. More precisely, we prove the following result in dual Malliavin–Sobolev spaces.

Theorem 5.1 (Weak correctors). *Let $d > 1$. Higher-order weak correctors $\{\varphi^n\}_{0 \leq n \leq d}$, flux correctors $\{\sigma^n\}_{0 \leq n \leq d}$, fluxes $\{q^n\}_{1 \leq n \leq d}$, and effective tensors $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq d}$ are uniquely well-defined iteratively as follows:*

³The topology on $\mathcal{S}(\Omega)$ is chosen as follows: a sequence $(X_\varepsilon)_\varepsilon \subset \mathcal{S}(\Omega)$ is said to converge to X in $\mathcal{S}(\Omega)$ as $\varepsilon \downarrow 0$ if there exists $n \geq 1$ such that for all ε small enough the element X_ε can be represented as $X_\varepsilon = g_\varepsilon(\xi(h_\varepsilon^1), \dots, \xi(h_\varepsilon^n))$ with $h_\varepsilon^j \rightarrow h^j$ in $C_c^\infty(\mathbb{R}^d)^\kappa$, $g_\varepsilon \rightarrow g$ in $C_b^\infty(\mathbb{R}^n)$, and $X = g(\xi(h^1), \dots, \xi(h^n))$.

- $\varphi^0 := 1$ and for all $1 \leq n \leq d$ we define $\varphi^n := (\varphi_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\varphi_{i_1 \dots i_n}^n$ is the unique distributional solution in $W_{\text{loc}}^{1, \infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$ for some $1 < p, q < \infty$ of

$$-\nabla \cdot \mathbf{a} \nabla \varphi_{i_1 \dots i_n}^n = \nabla \cdot ((\mathbf{a} \varphi_{i_1 \dots i_{n-1}}^{n-1} - \sigma_{i_1 \dots i_{n-1}}^{n-1}) e_{i_n}), \quad \text{in } \mathbb{R}^d, \quad (5.1)$$

such that φ^n is stationary with $\mathbb{E}[\varphi^n] = 0$ for all $n < d$, and such that $\nabla \varphi^d$ is stationary with $\mathbb{E}[\nabla \varphi^d] = 0$ and with anchoring $\int_B \varphi^d = 0$.

- $\sigma^0 := 0$ and for all $1 \leq n \leq d$ we define $\sigma^n := (\sigma_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\sigma_{i_1 \dots i_n}^n$ is the unique distributional solution in $W_{\text{loc}}^{1, \infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega)^{d \times d})$ for some $1 < p, q < \infty$, with skew-symmetric matrix values, of

$$-\Delta \sigma_{i_1 \dots i_n}^n = \nabla \times q_{i_1 \dots i_n}^n, \quad \nabla \cdot \sigma_{i_1 \dots i_n}^n = q_{i_1 \dots i_n}^n, \quad \text{in } \mathbb{R}^d, \quad (5.2)$$

such that σ^n is stationary with $\mathbb{E}[\sigma^n] = 0$ for all $n < d$, and such that $\nabla \sigma^d$ is stationary with $\mathbb{E}[\nabla \sigma^d] = 0$ and with anchoring $\int_B \sigma^d = 0$.

- For all $1 \leq n \leq d$ we define $q^n := (q_{i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ with $q_{i_1 \dots i_n}^n$ given by

$$q_{i_1 \dots i_n}^n := \mathbf{a} \nabla \varphi_{i_1 \dots i_n}^n + (\mathbf{a} \varphi_{i_1 \dots i_{n-1}}^{n-1} - \sigma_{i_1 \dots i_{n-1}}^{n-1}) e_{i_n} - \bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n e_{i_n}. \quad (5.3)$$

- For all $1 \leq n \leq d$ we define $\bar{\mathbf{a}}^n := (\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n)_{1 \leq i_1, \dots, i_{n-1} \leq d}$ with $\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n$ the matrix given by

$$\bar{\mathbf{a}}_{i_1 \dots i_{n-1}}^n e_{i_n} := \mathbb{E}[\mathbf{a}(\nabla \varphi_{i_1 \dots i_n}^n + \varphi_{i_1 \dots i_{n-1}}^{n-1} e_{i_n})]. \quad (5.4)$$

For $n < \ell = \lceil \frac{d}{2} \rceil$, these weak correctors φ^n, σ^n coincide with the strong correctors in Theorem 3.2. In addition, the following estimates hold:

- (i) Weak corrector estimates: for all $1 \leq n < d$, $X \in \mathcal{S}(\Omega)$, $r < \frac{d}{n}$, and $\delta > 0$,

$$|\mathbb{E}[X(\varphi^n, \sigma^n)(x)]| + |\mathbb{E}[X \nabla(\varphi^{n+1}, \sigma^{n+1})(x)]| \lesssim_{r, \delta} \| [D_0 X^\sharp]_2 \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))},$$

and at critical order $n = d$, for all $X \in \mathcal{S}(\Omega)$, $r < \frac{d}{d-\eta}$, and $\eta, \delta > 0$,

$$|\mathbb{E}[X(\varphi^d, \sigma^d)(x)]| \lesssim_{r, \eta, \delta} \langle x \rangle^\eta \| [D_0 X^\sharp]_2 \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))}.$$

- (ii) Weak fluctuation scaling: for all $1 \leq n \leq d$, $X \in \mathcal{S}(\Omega)$, $g \in C_c^\infty(\mathbb{R}^d)$, p, r with $\frac{1}{p} + \frac{1}{r} > 1 + \frac{n-1}{d}$, and $\delta > 0$,

$$\left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g(\nabla \varphi^n, \nabla \sigma^n) \right] \right| \lesssim_{p, r, \delta} \|g\|_{L^p(\mathbb{R}^d)} \| [D_0 X^\sharp]_2 \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))}.$$

For $n \leq \ell = \lceil \frac{d}{2} \rceil$, the space $L^{2+\delta}(\Omega)$ can be replaced by $L^{1+\delta}(\Omega)$ in these estimates. \diamond

Remark 5.2 (Explicit 1D case). In dimension $d = 1$, the first corrector is explicit, see Remark 2.1(b). It is easily checked to be uniquely defined as the stationary solution φ^1 of (5.1) in $W_{\text{loc}}^{1, \infty}(\mathbb{R}; \mathcal{S}'(\Omega))$ with $\mathbb{E}[\varphi^1] = 0$, and its explicit formula entails for all $X \in \mathcal{S}(\Omega)$, $g \in C_c^\infty(\mathbb{R}^d)$, and $\frac{1}{p} + \frac{1}{r} \geq 1$,

$$\begin{aligned} |\mathbb{E}[X \varphi^1(x)]| &\lesssim \|DX\|_{L^1(\mathbb{R}^d; L^1(\Omega))}, \\ \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \nabla \varphi^1 \right] \right| &\lesssim \|g\|_{L^p(\mathbb{R}^d)} \|DX\|_{L^r(\mathbb{R}^d; L^1(\Omega))}. \end{aligned} \quad \diamond$$

5.1. Periodic approximation. In order to prove Theorem 5.1, we proceed by periodic approximation. For $L \geq 3$, we replace the covariance function \mathcal{C} by its L -periodization,

$$\mathcal{C}_L(x) := \sum_{z \in LZ^d} \mathcal{C}(x+z),$$

we consider the \mathbb{R}^k -valued centered stationary Gaussian random field G_L on \mathbb{R}^d with covariance function \mathcal{C}_L , and we define the associated coefficient field

$$\mathbf{a}_L(x) := a_0(G_L(x)). \quad (5.5)$$

As the covariance function \mathcal{C}_L is L -periodic, we note that G_L and \mathbf{a}_L are L -periodic almost surely. In this periodic setting, using Poincaré's inequality on the periodic cell $Q_L = [-\frac{L}{2}, \frac{L}{2}]^d$, the associated periodic correctors are obviously well-defined to all orders.

Lemma 5.3 (Periodized correctors). *Let $d \geq 1$. Higher-order periodized correctors $\{\varphi_L^n\}_{n \geq 0}$, flux correctors $\{\sigma_L^n\}_{n \geq 0}$, fluxes $\{q_L^n\}_{n \geq 1}$, and effective tensors $\{\bar{\mathbf{a}}_L^n\}_{n \geq 1}$ are uniquely well-defined iteratively as follows:*

- $\varphi_L^0 := 1$ and for all $n \geq 1$ we define $\varphi_L^n := (\varphi_{L;i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\varphi_{L;i_1 \dots i_n}^n$ is the unique weak solution in $H_{\text{per}}^1(Q_L; L^2(\Omega))$, with $\int_{\mathbb{T}_L^d} \varphi_{L;i_1 \dots i_n}^n = 0$, of

$$-\nabla \cdot \mathbf{a}_L \nabla \varphi_{L;i_1 \dots i_n}^n = \nabla \cdot ((\mathbf{a}_L \varphi_{L;i_1 \dots i_{n-1}}^{n-1} - \sigma_{L;i_1 \dots i_{n-1}}^{n-1}) e_{i_n}), \quad \text{in } Q_L. \quad (5.6)$$

- $\sigma_L^0 := 0$ and for all $n \geq 1$ we define $\sigma_L^n := (\sigma_{L;i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ where $\sigma_{L;i_1 \dots i_n}^n$ is the unique weak solution in $H_{\text{per}}^1(Q_L; L^2(\Omega)^{d \times d})$, with $\int_{\mathbb{T}_L^d} \sigma_{L;i_1 \dots i_n}^n = 0$ and with skew-symmetric matrix values, of

$$-\Delta \sigma_{L;i_1 \dots i_n}^n = \nabla \times q_{L;i_1 \dots i_n}^n, \quad \nabla \cdot \sigma_{L;i_1 \dots i_n}^n = q_{L;i_1 \dots i_n}^n, \quad \text{in } Q_L. \quad (5.7)$$

- For all $n \geq 1$ we define $q_L^n := (q_{L;i_1 \dots i_n}^n)_{1 \leq i_1, \dots, i_n \leq d}$ with $q_{L;i_1 \dots i_n}^n$ given by

$$q_{L;i_1 \dots i_n}^n := \mathbf{a}_L \nabla \varphi_{L;i_1 \dots i_n}^n + (\mathbf{a}_L \varphi_{L;i_1 \dots i_{n-1}}^{n-1} - \sigma_{L;i_1 \dots i_{n-1}}^{n-1}) e_{i_n} - \bar{\mathbf{a}}_{L;i_1 \dots i_{n-1}}^n e_{i_n}. \quad (5.8)$$

- For all $n \geq 1$ we define $\bar{\mathbf{a}}_L^n := (\bar{\mathbf{a}}_{L;i_1 \dots i_{n-1}}^n)_{1 \leq i_1, \dots, i_{n-1} \leq d}$ with $\bar{\mathbf{a}}_{L;i_1 \dots i_{n-1}}^n$ the matrix given by

$$\bar{\mathbf{a}}_{L;i_1 \dots i_{n-1}}^n e_{i_n} := \int_{Q_L} \mathbf{a}_L (\nabla \varphi_{L;i_1 \dots i_n}^n + \varphi_{L;i_1 \dots i_{n-1}}^{n-1} e_{i_n}). \quad (5.9) \quad \diamond$$

Proof. Let indices i_1, \dots, i_n be fixed and omitted in the notation. If $\varphi_L^{n-1}, \sigma_L^{n-1}$ are defined in $H_{\text{per}}^1(Q_L; L^2(\Omega))$, then the Lax–Milgram theorem together with Poincaré's inequality ensures that the corrector equation (5.6) admits a unique solution φ_L^n in $H_{\text{per}}^1(Q_L; L^2(\Omega))$ with $\int_{Q_L} \varphi_L^n = 0$. Next, the same argument ensures that the flux corrector equation $-\Delta \sigma_L^n = \nabla \times q_L^n$ admits a unique solution σ_L^n in $H_{\text{per}}^1(Q_L; L^2(\Omega))$ with $\int_{Q_L} \sigma_L^n = 0$. Taking the divergence in this flux corrector equation and noting that the corrector equation (5.6) entails $\nabla \cdot q_L^n = 0$, we find $-\Delta(\nabla \cdot \sigma_L^n) = -\Delta q_L^n$, which proves the relation $\nabla \cdot \sigma_L^n = q_L^n$ since both quantities have vanishing average. Energy estimates for φ_L^n, σ_L^n and Poincaré's inequality give

$$L^{-1} \|(\varphi_L^n, \sigma_L^n)\|_{L^2(Q_L)} \lesssim \|\nabla(\varphi_L^n, \sigma_L^n)\|_{L^2(Q_L)} \lesssim \|(\varphi_L^{n-1}, \sigma_L^{n-1})\|_{L^2(Q_L)}.$$

The conclusion follows by iteration. \square

We may naturally couple the periodized field G_L with G via the underlying white noise ξ in the representation (2.2). Indeed, denoting by $\mathcal{C}_{0,L}$ the L -periodization of \mathcal{C}_0 ,

$$\mathcal{C}_{0,L}(x) := \sum_{z \in L\mathbb{Z}^d} \mathcal{C}_0(x+z),$$

and noting that $\mathcal{C}_L(x) = \int_{Q_L} \mathcal{C}_{0,L}(x-y) \mathcal{C}_{0,L}(y) dy$, the following representation formula holds,

$$G_L(x) = \int_{Q_L} \mathcal{C}_{0,L}(x-\cdot) \xi. \quad (5.10)$$

In particular, the periodized field G_L is now constructed on the same probability space (Ω, \mathbb{P}) as G , and is $\sigma(\xi|_{Q_L})$ -measurable. We denote by $L^q(\Omega_L)$ the subspace of $\sigma(\xi|_{Q_L})$ -measurable elements of $L^q(\Omega)$, and we define the dense subspace $\mathcal{S}(\Omega_L)$ as in (2.4) with test functions h_j supported in Q_L . Periodized correctors φ_L^n, σ_L^n are elements of $H_{\text{per}}^1(Q_L; L^2(\Omega_L))$.

The periodized Gaussian field G_L is stationary in the sense that its law is invariant under spatial translations, but its stationarity is clearly no longer described via the same action T as in Section 4.1. Denote by $\xi_L := \sum_{z \in L\mathbb{Z}^d} (\xi \mathbf{1}_{Q_L})(\cdot+z)$ the periodization of the restriction $\xi|_{Q_L}$. Since $\xi|_{Q_L} = \xi_L|_{Q_L}$, the translation $x \mapsto \xi_L(\cdot+x)|_{Q_L}$ induces an action $T_L = \{T_{L;x}\}_{x \in \mathbb{R}^d}$ of the additive group $(\mathbb{R}^d, +)$ on the space of $\sigma(\xi|_{Q_L})$ -measurable random variables. As the law of $\xi_L(\cdot+x)|_{Q_L}$ does not depend on the shift $x \in \mathbb{R}^d$, the map $T_{L;x}$ is an isometry on $L^q(\Omega_L)$ for all $1 \leq q \leq \infty$. Also note that, for any $\sigma(\xi|_{Q_L})$ -measurable random variable X_L , the map $(x, \omega) \mapsto (T_{L;x} X_L)(\omega)$ is L -periodic in x , and is stochastically continuous and jointly measurable on $\mathbb{R}^d \times \Omega$.

In this periodized setting, stationarity is defined as follows: an L -periodic random field $\psi_L \in L_{\text{per}}^1(Q_L; L^1(\Omega_L))$ is stationary if there exists a random variable $X_L \in L^1(\Omega_L)$ such that $\psi_L(x, \omega) = (T_{L;x} X_L)(\omega)$. As before, for all q , this provides a canonical isomorphism between random variables in $L^q(\Omega_L)$ and L -periodic stationary random fields in $L_{\text{per}}^q(Q_L; L^q(\Omega_L))$: for $X_L \in L^q(\Omega_L)$ we define its L -periodic stationary extension $X_L^\sharp(x, \omega) = (T_{L;x} X_L)(\omega)$, and for an L -periodic stationary field $\psi_L \in L_{\text{per}}^q(Q_L; L^q(\Omega_L))$ we define $\psi_L^\flat \in L^q(\Omega_L)$ such that $\psi_L(x, \omega) = (T_{L;x} \psi_L^\flat)(\omega)$.

5.2. Weak bounds on periodized correctors. The following result states that the weak corrector estimates of Theorem 5.1 hold for periodized correctors uniformly with respect to the period. The argument is as follows: to estimate weak expressions like $\mathbb{E}[X_L \varphi_L^n]$, since the corrector φ_L^n roughly takes the form $[(-\nabla \cdot \mathbf{a}_L \nabla)^{-1} \nabla \cdot]^n \mathbf{a}_L$, we may migrate half of the iterated heterogeneous Riesz transforms $(-\nabla \cdot \mathbf{a}_L \nabla)^{-1} \nabla \cdot$ to the test function X_L , so that in the end we only need strong L^2 estimates on φ_L^m for $m \leq \frac{n}{2}$, which then allows to define twice as many weak correctors. We already mentioned this useful symmetrization trick in a related work with Otto [14, Remark 2.5]; see also [1, Theorem 3.5].

Proposition 5.4 (Weak bounds on periodized correctors).

(i) Weak corrector estimates: for all $1 \leq n < d$, $X_L \in \mathcal{S}(\Omega_L)$, $r < \frac{d}{n}$, and $\delta > 0$,

$$\begin{aligned} |\mathbb{E}[X_L(\varphi_L^n, \sigma_L^n)(x)]| + |\mathbb{E}[X_L \nabla(\varphi_L^{n+1}, \sigma_L^{n+1})(x)]| \\ \leq_{r,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}, \end{aligned} \quad (5.11)$$

and at critical order $n = d$, for all $X_L \in \mathcal{S}(\Omega_L)$, $r < \frac{d}{d-\eta}$, and $\eta, \delta > 0$,

$$\left| \mathbb{E} \left[X_L \left((\varphi_L^d, \sigma_L^d)(x) - \int_B (\varphi_L^d, \sigma_L^d) \right) \right] \right| \lesssim_{r, \eta, \delta} \langle x \rangle^\eta \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}. \quad (5.12)$$

(ii) Weak fluctuation scaling: for all $1 \leq n \leq d$, $X_L \in \mathcal{S}(\Omega_L)$, $g_L \in C_{\text{per}}^\infty(Q_L)$, p, r with $\frac{1}{p} + \frac{1}{r} > 1 + \frac{n-1}{d}$, and $\delta > 0$,

$$\left| \mathbb{E} \left[X_L \int_{Q_L} g_L (\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| \lesssim_{p, \delta} \| g_L \|_{L_{\text{per}}^p(Q_L)} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}. \quad (5.13)$$

For $n \leq \ell = \lceil \frac{d}{2} \rceil$, the space $L^{2+\delta}(\Omega_L)$ can be replaced by $L^{1+\delta}(\Omega_L)$ in these estimates. \diamond

Proof. We split the proof into five steps.

Step 1. Symmetrization: for all $n \geq \ell$, $X_L \in \mathcal{S}(\Omega_L)$, $g_L \in C_{\text{per}}^\infty(Q_L)$, and $1 \leq p, q \leq \infty$,

$$\begin{aligned} \left| \mathbb{E} \left[X_L \int_{\mathbb{T}_L^d} g_L (\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| &\lesssim \| [D_0 (\mathcal{R}_L^* \mathcal{K}_L^*)^{n-\ell} (X_L^\sharp *_L g_L)]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \\ &\quad \times \| [D_0 (\mathcal{K}_L \mathcal{R}_L)^{\ell-1} \mathcal{K}_L 1]_2 \|_{L^{p'}(Q_L; L^{q'}(\Omega_L))}, \end{aligned} \quad (5.14)$$

where $*_L$ stands for the convolution of L -periodic functions, and where we have defined the linear operator

$$\mathcal{K}_L := (\mathcal{H}_L^\circ, \mathcal{H}_L, \mathcal{H}^\circ \mathbf{a}_L, \mathcal{H}_L \mathbf{a}_L, \mathcal{H}_L^\circ \mathbf{a}_L \mathcal{H}_L, \mathcal{H}_L^\circ \mathbf{a}_L \mathcal{H}_L \mathbf{a}_L), \quad (5.15)$$

in terms of the following Helmholtz projections and Riesz transforms, acting on L -periodic functions,

$$\mathcal{H}_L^\circ := \nabla(-\Delta)^{-1} \nabla \cdot, \quad \mathcal{H}_L := \nabla(-\nabla \cdot \mathbf{a}_L \nabla)^{-1} \nabla \cdot, \quad \mathcal{R}_L := \Delta^{-1} \nabla \cdot,$$

and where we omit matrix contractions in the notation. Note that in (5.14) we do not share evenly the number of Riesz transforms on both sides, as we wish to have as little as possible such operators on the test function $X_L^\sharp *_L g_L$ in order to obtain the best norm.

We start by using stationarity in form of

$$\begin{aligned} \mathbb{E} \left[X_L \int_{Q_L} g_L (\nabla \varphi_L^n, \nabla \sigma_L^n) \right] &= \mathbb{E} \left[\left(\int_{Q_L} X_L^\sharp(-x) g_L(x) dx \right) (\nabla \varphi_L^n, \nabla \sigma_L^n)^b \right] \\ &= \mathbb{E} \left[(X_L^\sharp *_L g_L)^b (\nabla \varphi_L^n, \nabla \sigma_L^n)^b \right]. \end{aligned} \quad (5.16)$$

In terms of the operators $\mathcal{H}_L^\circ, \mathcal{H}_L, \mathcal{R}_L$, the equations for higher-order periodized correctors, flux correctors, and fluxes in Definition 5.3 take on the following form,

$$\begin{aligned} \nabla \varphi_L^n &= \mathcal{H}_L \mathbf{a}_L \mathcal{R}_L \nabla \varphi_L^{n-1} - \mathcal{H}_L \mathcal{R}_L \nabla \sigma_L^{n-1}, \\ \nabla \sigma_L^n &= \mathcal{H}_L^\circ \mathbf{a}_L \nabla \phi_L^n + \mathcal{H}_L^\circ \mathbf{a}_L \mathcal{R}_L \nabla \phi_L^{n-1} - \mathcal{H}_L^\circ \mathcal{R}_L \nabla \sigma_L^{n-1}, \end{aligned}$$

and thus by iteration,

$$(\nabla \varphi_L^n, \nabla \sigma_L^n) = (\mathcal{K}_L \mathcal{R}_L)^{n-1} \mathcal{K}_L 1,$$

where we omit matrix contractions in the notation. Inserting this into (5.16), and denoting by $\mathcal{K}_L^{\text{st}}, \mathcal{R}_L^{\text{st}}$ the corresponding lifted operators on $L^2(\Omega_L)$, we find

$$\left| \mathbb{E} \left[X_L \int_{Q_L} g_L (\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| \lesssim \left| \mathbb{E} \left[(X_L^\sharp *_L g_L)^b (\mathcal{K}_L^{\text{st}} \mathcal{R}_L^{\text{st}})^{n-1} \mathcal{K}_L^{\text{st}} 1 \right] \right|,$$

or equivalently, using adjoints,

$$\left| \mathbb{E} \left[X_L \int_{Q_L} g_L (\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| \lesssim |\mathbb{E} [((\mathcal{R}_L^{\text{st},*} \mathcal{K}_L^{\text{st},*})^{n-\ell} (X_L^\sharp *_L g_L)^b) ((\mathcal{K}_L^{\text{st}} \mathcal{R}_L^{\text{st}})^{\ell-1} \mathcal{K}_L^{\text{st}} 1)]|.$$

Now appealing to Lemma 4.3(iii) (restricted to $\mathcal{S}(\Omega_L)$), the claim (5.14) follows.

Step 2. Annealed estimates: for all $X_L \in \mathcal{S}(\Omega_L)$, $1 < p, q < \infty$, and $\delta > 0$,

$$\| [D_0 \mathcal{K}_L X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_{p,q,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^{p \wedge 2}(Q_L; L^{(q \vee 2) + \delta}(\Omega_L))}, \quad (5.17)$$

and for all $\frac{d}{d-1} < p < \infty$ and $1 \leq q \leq \infty$,

$$\| [D_0 \mathcal{R}_L X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_p \| [D_0 X_L^\sharp]_2 \|_{L^{\frac{dp}{d+p}}(Q_L; L^q(\Omega_L))}. \quad (5.18)$$

By the definition (5.15) of \mathcal{K}_L , the result (5.17) follows from the following three estimates, for all $1 < p, q < \infty$ and $\delta > 0$,

$$\| [D_0 \mathcal{H}_L^\circ X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_{p,q} \| [D_0 X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))}, \quad (5.19)$$

$$\| [D_0(\mathbf{a}_L X_L^\sharp)]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_{p,q} \| [D_0 X_L^\sharp]_2 \|_{L^{p \wedge 2}(Q_L; L^{q \vee 2}(\Omega_L))} + |\mathbb{E}[X_L]|, \quad (5.20)$$

$$\| [D_0 \mathcal{H}_L X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_{p,q,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^{p \wedge 2}(Q_L; L^{(q+\delta) \vee 2}(\Omega_L))}. \quad (5.21)$$

We start with the proof of (5.18). As the convolution kernel for the Riesz transform \mathcal{R}_L is bounded by $x \mapsto C|(x)_L|^{1-d}$, where we have set $(x)_L := x \bmod Q_L$, this estimate follows from the Hardy–Littlewood–Sobolev inequality in the form

$$\begin{aligned} \| [D_0 \mathcal{R}_L X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} &= \| [\mathcal{R}_L D_0 X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \\ &\lesssim \left(\int_{Q_L} \left(\int_{Q_L} |(x-y)_L|^{1-d} \| [D_0 X_L^\sharp]_2(y) \|_{L^q(\Omega_L)} dy \right)^p dx \right)^{\frac{1}{p}} \\ &\lesssim_p \| [D_0 X_L^\sharp]_2 \|_{L^{\frac{dp}{d+p}}(Q_L; L^q(\Omega_L))}. \end{aligned}$$

We turn to the proof of (5.19). In view of Banach-valued Fourier multiplier theorems, e.g. in form of the extrapolation result in [25, Theorem 3.15], this follows from the maximal L^p regularity for the Helmholtz projection \mathcal{H}_L° with values in $L^q(\Omega_L)$.

We turn to the proof of (5.20). In view of (5.5) and (5.10), we find $|D_0 \mathbf{a}_L(x)| \lesssim |\mathcal{C}_{0,L}(x)|$, and the integrability condition (2.3) yields $\int_{Q_L} |\mathcal{C}_{0,L}|_\infty \leq \int_{\mathbb{R}^d} |\mathcal{C}_0|_\infty \leq 1$. We deduce

$$\begin{aligned} &\| [D_0(\mathbf{a}_L X_L^\sharp)]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \\ &\lesssim \| [D_0 X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} + \| [(D_0 \mathbf{a}_L) X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \\ &\lesssim \| [D_0 X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} + \| [X_L^\sharp]_2^b \|_{L^q(\Omega_L)}. \end{aligned}$$

Decomposing $\| [X_L^\sharp]_2^b \|_{L^q(\Omega_L)} \leq \| [(X_L - \mathbb{E}[X_L])^\sharp]_2^b \|_{L^q(\Omega_L)} + |\mathbb{E}[X_L]|$, appealing to Poincaré's inequality in form of Lemma 4.3(iv) (restricted to $\mathcal{S}(\Omega_L)$), and using the discrete $\ell^p - \ell^r$ inequality as in (4.10), the claim (5.20) follows.

It remains to establish (5.21). We start from (4.15) in the proof of Corollary 4.4 (restricted to $\mathcal{S}(\Omega_L)$), in form of

$$\| [D_0 \mathcal{H}_L X_L^\sharp]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \lesssim_{p,q,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^p(Q_L; L^{q+\delta}(\Omega_L))} + \| [\mathcal{H}_L X_L^\sharp]_2^b \|_{L^{q+\delta}(\Omega_L)}.$$

Noting that $\mathcal{H}_L X_L^\sharp = \mathcal{H}_L(X_L - \mathbb{E}[X_L])^\sharp$, appealing to the stationary L^q regularity estimate of Corollary 4.1, and using Poincaré’s inequality in form of Lemma 4.3(iv), the last right-hand side term is bounded by

$$\begin{aligned} \|[\mathcal{H}_L X_L^\sharp]_2^\flat\|_{L^{q+\delta}(\Omega_L)} &\lesssim_{q,\delta} \|[(X_L - \mathbb{E}[X_L])^\sharp]_2^\flat\|_{L^{q+2\delta}(\Omega_L)} \\ &\lesssim_{q,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^2(Q_L; L^{(q+2\delta)\vee 2}(\Omega_L))}, \end{aligned}$$

and the claim (5.21) follows.

Step 3. Weak fluctuation scaling (5.13).

For $\ell < n \leq d$, combining (5.14) with (5.17)–(5.18), we find for all $p < \frac{d}{n-1}$ and $\delta > 0$,

$$\left| \mathbb{E} \left[X_L \int_{Q_L} g_L(\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| \lesssim_{p,\delta} \| [D_0(X_L^\sharp *_L g_L)]_2 \|_{L^p(Q_L; L^{2+\delta}(\Omega_L))}. \quad (5.22)$$

Writing $D_0(X_L^\sharp *_L g_L) = g_L *_L (D_0 X_L^\sharp)$ and appealing to Young’s convolution inequality, the conclusion (5.13) follows.

The case $n \leq \ell$ is simpler and is essentially already contained in [14, Lemma 7.1]. More precisely, it suffices to avoid the symmetrization trick of Step 1: we replace (5.14) by

$$\begin{aligned} \left| \mathbb{E} \left[X_L \int_{Q_L} g_L(\nabla \varphi_L^n, \nabla \sigma_L^n) \right] \right| &\lesssim \| [D_0(X_L^\sharp *_L g_L)]_2 \|_{L^p(Q_L; L^q(\Omega_L))} \\ &\quad \times \| [D_0(\mathcal{K}_L \mathcal{R}_L)^{n-1} \mathcal{K}_L 1]_2 \|_{L^{p'}(Q_L; L^{q'}(\Omega_L))}, \end{aligned}$$

which can now be estimated by using (5.17)–(5.18) as above.

Step 4. Weak corrector estimates (5.11).

Let $1 \leq n < d$. The bound on $\nabla(\varphi_L^{n+1}, \sigma_L^{n+1})$ already follows from (5.22) after replacing g_L by a Dirac measure, and it remains to estimate $(\varphi_L^n, \sigma_L^n)$ itself. In view of the anchoring $\int_{Q_L} (\varphi_L^n, \sigma_L^n) = 0$, we may decompose

$$\begin{aligned} (\varphi_L^n, \sigma_L^n)(x) &= (\varphi_L^n, \sigma_L^n)(x) - \int_{Q_L} (\varphi_L^n, \sigma_L^n) \\ &= \int_0^L \left(\int_{Q_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^n, \nabla \sigma_L^n)(y) dy \right) dt, \end{aligned} \quad (5.23)$$

and we separately analyze the contribution of the integral for $r \in [0, 1]$ and for $r \in [1, L]$. On the one hand, appealing to (5.13) with $p = 1$, we find for all $r < \frac{d}{n-1}$ and $\delta > 0$,

$$\begin{aligned} &\mathbb{E} \left[X_L \int_0^1 \left(\int_{Q_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^n, \nabla \sigma_L^n)(y) dy \right) dt \right] \\ &\lesssim_{r,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))} \int_0^1 \int_{Q_t(x)} \left| \frac{x-y}{t} \right| dy dt \\ &\leq \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}. \end{aligned} \quad (5.24)$$

On the other hand, appealing again to (5.13), we obtain for all p, r with $\frac{1}{p} + \frac{1}{r} > 1 + \frac{n-1}{d}$, and $\delta > 0$,

$$\begin{aligned} \mathbb{E} \left[X_L \int_1^L \left(\int_{Q_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^n, \nabla \sigma_L^n)(y) dy \right) dt \right] \\ \lesssim_{p,r,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))} \int_1^L t^{-d(1-\frac{1}{p})} dt, \end{aligned}$$

and thus, for all $r < \frac{d}{n}$ and $\delta > 0$, choosing $p > \frac{d}{d-1}$ such that $\frac{1}{p} + \frac{1}{r} > 1 + \frac{n-1}{d}$,

$$\mathbb{E} \left[X_L \int_1^L \left(\int_{B_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^n, \nabla \sigma_L^n)(y) dy \right) dt \right] \lesssim_{r,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}.$$

Combining this with (5.23) and (5.24), the conclusion (5.11) follows. In the case $n \leq \ell$, as in (5.13), the space $L^{2+\delta}(\Omega_L)$ can be replaced by $L^{1+\delta}(\Omega_L)$.

Step 5. Weak sublinearity estimate (5.12) for φ_L^d .

We decompose

$$\begin{aligned} (\varphi_L^d, \sigma_L^d)(x) - \int_B (\varphi_L^d, \sigma_L^d) \\ = \left((\varphi_L^d, \sigma_L^d)(x) - \int_{B(x)} (\varphi_L^d, \sigma_L^d) \right) + \left(\int_{B(x)} (\varphi_L^d, \sigma_L^d) - \int_B (\varphi_L^d, \sigma_L^d) \right) \\ = \int_0^1 \left(\int_{B_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^d, \nabla \sigma_L^d)(y) dy \right) dt + \left(\int_{B(x)} (\varphi_L^d, \sigma_L^d) - \int_B (\varphi_L^d, \sigma_L^d) \right), \quad (5.25) \end{aligned}$$

and we analyze the two right-hand side terms separately. The first one is estimated as in (5.24), for all $r < \frac{d}{d-1}$ and $\delta > 0$,

$$\mathbb{E} \left[X_L \int_0^1 \left(\int_{B_t(x)} \frac{x-y}{t} \cdot (\nabla \varphi_L^d, \nabla \sigma_L^d)(y) dy \right) dt \right] \lesssim_{r,\delta} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}. \quad (5.26)$$

We turn to the second right-hand side term in (5.25). Denoting by $\chi_L \in H_{\text{per}}^1(Q_L)$ the unique solution of

$$-\Delta \chi_L = \frac{\mathbf{1}_B}{|B|} - L^{-d}, \quad \text{in } Q_L,$$

and setting $\nabla \chi_{x,L} := \nabla \chi_L(\cdot - x) - \nabla \chi_L$, we can write

$$\mathbb{E} \left[X_L \left(\int_{B(x)} (\varphi_L^d, \sigma_L^d) - \int_B (\varphi_L^d, \sigma_L^d) \right) \right] = \mathbb{E} \left[X_L \int_{Q_L} \nabla \chi_{x,L} \cdot (\nabla \varphi_L^d, \nabla \sigma_L^d) \right].$$

Appealing to (5.13), we deduce for all p, r with $\frac{1}{p} + \frac{1}{r} > 1 + \frac{d-1}{d}$, and $\delta > 0$,

$$\begin{aligned} \mathbb{E} \left[X_L \left(\int_{B(x)} (\varphi_L^d, \sigma_L^d) - \int_B (\varphi_L^d, \sigma_L^d) \right) \right] \\ \lesssim_{p,r,\delta} \| \nabla \chi_{x,L} \|_{L^p(Q_L)} \| [D_0 X_L^\sharp]_2 \|_{L^r(Q_L; L^{2+\delta}(\Omega_L))}. \quad (5.27) \end{aligned}$$

Noting that a direct computation yields

$$\|\nabla\chi_{x,L}\|_{L^p(\mathbb{R}^d)} \lesssim \begin{cases} 1 & : p > \frac{d}{d-1}, \\ \log(2+|x|)^{1-\frac{1}{d}} & : p = \frac{d}{d-1}, \\ \langle x \rangle^{d(\frac{1}{p}+\frac{1}{d}-1)} & : p < \frac{d}{d-1}, \end{cases}$$

and combining (5.25), (5.26), and (5.27), the conclusion (5.12) easily follows. In the case $n \leq \ell$, as in (5.13), the space $L^{2+\delta}(\Omega_L)$ can be replaced by $L^{1+\delta}(\Omega_L)$. \square

5.3. Proof of Theorem 5.1. It remains to pass to the infinite-period limit in the periodized corrector estimates of Proposition 5.4, and to show that the limiting correctors are indeed uniquely defined in a weak sense. We start with the following uniqueness statement in dual Malliavin–Sobolev spaces.

Lemma 5.5 (Uniqueness). *If ψ belongs to $W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$ for some $1 < p, q < \infty$, if $\nabla\psi$ is stationary with $\mathbb{E}[\nabla\psi] = 0$, and if the equation $-\nabla \cdot \mathbf{a}\nabla\psi = 0$ is satisfied in \mathbb{R}^d in the distributional sense, then there holds $\nabla\psi = 0$.* \diamond

Proof. Let $1 < p, q < \infty$ be fixed. We split the proof into three steps.

Step 1. Special case $\mathbf{a} = \text{Id}$: If ψ belongs to $W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$, if $\nabla\psi$ is stationary with $\mathbb{E}[\nabla\psi] = 0$, and if the equation $-\Delta\psi = 0$ is satisfied in \mathbb{R}^d in the distributional sense, then there holds $\nabla\psi = 0$.

Given $X \in \mathcal{S}(\Omega)$, we consider the function $u_X = \mathbb{E}[X\psi] \in C^\infty(\mathbb{R}^d)$. As $\nabla\psi$ is stationary, we can write

$$\nabla u_X(x) = \mathbb{E}[X\nabla\psi(x)] = \mathbb{E}[(T_{-x}X)(\nabla\psi)^{\flat}]. \quad (5.28)$$

For $(\nabla\psi)^{\flat} \in (\mathbb{M}_{p,q}^1)'(\Omega)$, as the norm of $T_{-x}X$ in $\mathbb{M}_{p,q}^1(\Omega)$ is bounded independently of x , we deduce that ∇u_X is uniformly bounded. As the relation $-\Delta\psi = 0$ implies that u_X is harmonic, we deduce that ∇u_X is constant. In particular, identity (5.28) then yields for all $R > 0$,

$$\nabla u_X(x) = f_{B_R} \nabla u_X = \mathbb{E}[(f_{B_R} X^{\sharp})(\nabla\psi)^{\flat}].$$

Noting that $f_{B_R} X^{\sharp}$ converges to $\mathbb{E}[X]$ in $\mathbb{M}_{p,q}^1(\Omega)$ as $R \uparrow \infty$, and recalling $\mathbb{E}[\nabla\psi] = 0$, we deduce

$$\nabla u_X(x) = \mathbb{E}[X] \mathbb{E}[(\nabla\psi)^{\flat}] = 0.$$

This means $\mathbb{E}[X\nabla\psi] = \nabla u_X = 0$, and the arbitrariness of X yields the claim $\nabla\psi = 0$.

Step 2. Density result: If ψ belongs to $W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$ with $\nabla\psi$ stationary and $\mathbb{E}[\nabla\psi] = 0$, then there exists a sequence $(X_n)_n \subset \mathcal{S}(\Omega)$ such that $\nabla^{\text{st}} X_n$ converges to $(\nabla\psi)^{\flat}$ weakly-* in $(\mathbb{M}_{p,q}^1)'(\Omega)$.

Let $\psi \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$ with $\nabla\psi$ stationary and $\mathbb{E}[\nabla\psi] = 0$. As $(\nabla\psi)^{\flat} \in (\mathbb{M}_{p,q}^1)'(\Omega)^d$, by density, there is a sequence $(Y_m)_m \subset \mathcal{S}(\Omega)^d$ that converges to $(\nabla\psi)^{\flat}$ weakly-* in $(\mathbb{M}_{p,q}^1)'(\Omega)$. Next, for all $m, r \geq 1$, we define $X_{m,r} \in H^1(\Omega)$ as the unique solution of

$$\left(\frac{1}{r} - \Delta^{\text{st}}\right)X_{m,r} = -\nabla^{\text{st}} \cdot Y_m. \quad (5.29)$$

The energy estimate for this equation takes the form

$$r^{-\frac{1}{2}}\|X_{m,r}\|_{L^2(\Omega)} + \|\nabla^{\text{st}} X_{m,r}\|_{L^2(\Omega)} \lesssim \|Y_m\|_{L^2(\Omega)},$$

and the proof of Corollary 4.4 (in the simpler case $\mathbf{a} = \text{Id}$) yields, up to a duality argument,

$$\|\nabla^{\text{st}} X_{m,r}\|_{(\mathbb{M}_{p,q}^1)'(\Omega)} \lesssim_{p,q} \|Y_m\|_{(\mathbb{M}_{p,q}^1)'(\Omega)}.$$

A weak compactness argument then allows to find a diagonal subsequence $(\nabla^{\text{st}} X_n)_n \subset (\nabla^{\text{st}} X_{m,r})_{m,r}$ that converges weakly- $*$ in $(\mathbb{M}_{p,q}^1)'(\Omega)$ to some limit L with $\mathbb{E}[L] = 0$, in such a way that we may pass to the limit in (5.29) and obtain

$$-\nabla^{\text{st}} \cdot L = -\nabla^{\text{st}} \cdot (\nabla\psi)^{\flat}. \quad (5.30)$$

In addition, the gradient structure of $(\nabla^{\text{st}} X_n)_n$ allows us to write the limit as $L = (\nabla\phi)^{\flat}$ for some $\phi \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$. The relation (5.30) then becomes

$$-\nabla^{\text{st}} \cdot (\nabla\phi)^{\flat} = -\nabla^{\text{st}} \cdot (\nabla\psi)^{\flat}.$$

In view of Step 1, we can conclude $L^{\sharp} = \nabla\phi = \nabla\psi$. Further approximating the X_n 's by elements in $\mathcal{S}(\Omega)$, the claimed density result follows.

Step 3. Conclusion.

Let ψ belong to $W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; (\mathbb{M}_{p,q}^1)'(\Omega))$ with $\nabla\psi$ stationary and $\mathbb{E}[\nabla\psi] = 0$, and assume that $-\nabla \cdot \mathbf{a}\nabla\psi = 0$ is satisfied in \mathbb{R}^d in the distributional sense. In particular, $(\nabla\psi)^{\flat}$ belongs to $(\mathbb{M}_{p,q}^1)'(\Omega)$ and satisfies for all $X \in \mathcal{S}(\Omega)$,

$$\mathbb{E}[\nabla^{\text{st}} X \cdot \mathbf{a}^{\flat}(\nabla\psi)^{\flat}] = 0. \quad (5.31)$$

Given $Y \in \mathcal{S}(\Omega)^d$, consider the unique solution $w_Y \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; L^2(\Omega))$ of

$$-\nabla \cdot \mathbf{a}^* \nabla w_Y = \nabla \cdot Y^{\sharp},$$

such that ∇w_Y is stationary with $\mathbb{E}[\nabla w_Y] = 0$ and with anchoring $\int_B w_Y = 0$. For all $X \in \mathcal{S}(\Omega)$, this equation yields

$$\mathbb{E}[(\nabla w_Y)^{\flat} \cdot \mathbf{a}^{\flat} \nabla^{\text{st}} X] = -\mathbb{E}[Y \cdot \nabla^{\text{st}} X], \quad (5.32)$$

and we note that Corollary 4.4 with $Y \in \mathcal{S}(\Omega)^d$ ensures that $w_Y \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^d; \mathbb{M}_{p,q}^1(\Omega))$ for all $1 < p, q < \infty$. Appealing to the density result of Step 2, we deduce from (5.31)–(5.32),

$$0 = \mathbb{E}[(\nabla w_Y)^{\flat} \cdot \mathbf{a}^{\flat}(\nabla\psi)^{\flat}] = -\mathbb{E}[Y \cdot (\nabla\psi)^{\flat}].$$

and the arbitrariness of Y yields the conclusion $\nabla\psi = 0$. \square

With this uniqueness result at hand, we may now pass to the infinite-period limit in the estimates of Proposition 5.4 and conclude with the proof of Theorem 5.1.

Proof of Theorem 5.1. For all $X \in \mathcal{S}(\Omega)$ and $g \in C_c^\infty(\mathbb{R}^d)$, there is some large enough $L_0 > 0$ such that $X \in \mathcal{S}(\Omega_L)$ and $g \in C_{\text{per}}^\infty(Q_L)$ for all $L \geq L_0$. Therefore, the bounds of Proposition 5.4 ensure that for all $n \leq d$ the sequence $(\varphi_L^n, \sigma_L^n)_{L \geq 1}$ is uniformly bounded in $C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$. By a weak- $*$ compactness argument, we deduce that up to extraction of a subsequence $(\varphi_L^n, \sigma_L^n)$ converges weakly- $*$ to some (φ^n, σ^n) in $C^\infty(\mathbb{R}^d; \mathcal{S}'(\Omega))$, such that (φ^n, σ^n) is stationary with $\mathbb{E}[(\varphi^n, \sigma^n)] = 0$ for all $n < d$, and such that $(\nabla\varphi^d, \nabla\sigma^d)$ is stationary with $\mathbb{E}[(\nabla\varphi^d, \nabla\sigma^d)] = 0$ and anchoring $\int_B (\varphi^d, \sigma^d) = 0$. Passing to the limit in the periodized corrector equations (5.6)–(5.9) and in the estimates (5.11)–(5.13), we easily deduce that the limiting collection $\{\varphi^n, \sigma^n\}_{1 \leq n \leq d}$ satisfies equations (5.1)–(5.4) in the distributional sense as well as the corresponding limiting estimates. Finally, Lemma 5.5 ensures that those limiting objects are uniquely defined. \square

6. EFFECTIVE DESCRIPTION OF ENSEMBLE AVERAGES

This section is devoted to the proof of Theorem 2, that is, the first half of the Bourgain–Spencer conjecture. More precisely, we establish the following error estimates.

Theorem 6.1 (Effective description of ensemble averages). *Let $d > 1$, let the higher-order effective tensors $\{\bar{\mathbf{a}}^n\}_{1 \leq n \leq d}$ be defined in Theorem 5.1, and let the associated higher-order homogenized solution operators be given in Definition 3.3. Then, for all $f \in C_c^\infty(\mathbb{R}^d)^d$, we have for all $n < d$, $\frac{d}{d-n} \vee 2 < p < \infty$, and $\eta > 0$,*

$$\begin{aligned} & \left\| \left[\mathbb{E}[\nabla u_{\varepsilon, f}] - \nabla \bar{\mathcal{U}}_\varepsilon^n[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} + \left\| \left[\mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon, f}] - \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{\mathcal{U}}_\varepsilon^n[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} \\ & \lesssim_{p, \eta} \varepsilon^n \|\langle \cdot \rangle^\eta \langle \nabla \rangle^{2n-1} f\|_{L^p(\mathbb{R}^d)}, \end{aligned} \quad (6.1)$$

while at critical order $n = d$ we have for all $\frac{d}{\eta} < p < \infty$ and $0 < \eta \ll 1$,

$$\begin{aligned} & \left\| \left[\mathbb{E}[\nabla u_{\varepsilon, f}] - \nabla \bar{\mathcal{U}}_\varepsilon^d[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} + \left\| \left[\mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon, f}] - \bar{\mathcal{A}}_\varepsilon^d \nabla \bar{\mathcal{U}}_\varepsilon^d[f] \right]_{2; \varepsilon} \right\|_{L^p(\mathbb{R}^d)} \\ & \lesssim_{p, \eta} \varepsilon^{d-\eta} \|\langle \cdot \rangle^\eta \langle \nabla \rangle^{2d-1} f\|_{L^p(\mathbb{R}^d)}. \quad \diamond \end{aligned} \quad (6.2)$$

Proof. We consider the ensemble-averaged field and the ensemble-averaged flux separately, and we split the proof into four steps.

Step 1. Preliminary: Given a test function $g \in C_c^\infty(\mathbb{R}^d)^d$, we consider the unique almost sure gradient solution $\nabla v_{\varepsilon, g}$ in $L^2(\mathbb{R}^d)^d$ of the auxiliary equation

$$-\nabla \cdot \mathbf{a}^*(\frac{\cdot}{\varepsilon}) \nabla v_{\varepsilon, g} = \nabla \cdot g, \quad \text{in } \mathbb{R}^d, \quad (6.3)$$

and we show that for all $h \in C_c^\infty(\mathbb{R}^d)$, $X \in \mathcal{S}(\Omega)$, $1 < r < \frac{d}{d-\eta}$, and $0 < \eta, \delta \ll 1$,

$$\begin{aligned} A_\varepsilon(X, h, g) & := \mathbb{E} \left[X \int_{\mathbb{R}^d} \varepsilon^d (\varphi^d, \sigma^d) \left(\frac{\cdot}{\varepsilon} \right) h \nabla v_{\varepsilon, g} \right] \\ & \lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \left(\|X\|_{L^\infty(\Omega)} + \|[D_0 X^\sharp]_\infty\|_{L^r(\mathbb{R}^d; L^\infty(\Omega))} \right) \|\langle \cdot \rangle^\eta h\|_{L^{r'}(\mathbb{R}^d)} \|g\|_{2+\delta; \varepsilon} \|g\|_{L^r(\mathbb{R}^d)}. \end{aligned} \quad (6.4)$$

Starting point is the weak corrector estimate of Theorem 5.1(i), which yields for all $r < \frac{d}{d-\eta}$ and $\eta, \delta > 0$,

$$\begin{aligned} A_\varepsilon(X, h, g) & \leq \varepsilon^d \int_{\mathbb{R}^d} |h(x)| \left| \mathbb{E} \left[X \nabla v_{\varepsilon, g}(x) (\varphi^d, \sigma^d) \left(\frac{x}{\varepsilon} \right) \right] \right| dx \\ & \lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \int_{\mathbb{R}^d} \langle x \rangle^\eta |h(x)| \\ & \quad \times \left(\int_{\mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0(X^\sharp(y')) T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dy \right)^{\frac{1}{r}} dx. \end{aligned}$$

Using Leibniz' rule for D_0 , we split the right-hand side into two terms: $A_\varepsilon^1(X, h, g)$ and $A_\varepsilon^2(X, h, g)$, where the first one corresponds to $(D_0 X^\sharp(y')) T_{y'} \nabla v_{\varepsilon, g}(x)$, while the second

one corresponds to $X^\sharp(y')(D_0 T_{y'} \nabla v_{\varepsilon, g}(x))$. Straightforward estimates lead to

$$\begin{aligned} A_\varepsilon^1(X, h, g) &\lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \| [D_0 X^\sharp]_\infty \|_{L^r(\mathbb{R}^d; L^\infty(\Omega))} \int_{\mathbb{R}^d} \langle x \rangle^\eta |h(x)| \| \nabla v_{\varepsilon, g}(x) \|_{L^{2+\delta}(\Omega)} dx, \\ A_\varepsilon^2(X, h, g) &\lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \| X \|_{L^\infty(\Omega)} \\ &\quad \times \int_{\mathbb{R}^d} \langle x \rangle^\eta |h(x)| \left(\int_{\mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dy \right)^{\frac{1}{r}} dx, \end{aligned}$$

and thus, by Hölder's inequality,

$$A_\varepsilon^1(X, h, g) \lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \| [D_0 X^\sharp]_\infty \|_{L^r(\mathbb{R}^d; L^\infty(\Omega))} \| \langle \cdot \rangle^\eta h \|_{L^{r'}(\mathbb{R}^d)} \| \nabla v_{\varepsilon, g} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))}, \quad (6.5)$$

$$\begin{aligned} A_\varepsilon^2(X, h, g) &\lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \| X \|_{L^\infty(\Omega)} \| \langle \cdot \rangle^\eta h \|_{L^{r'}(\mathbb{R}^d)} \\ &\quad \times \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}}. \quad (6.6) \end{aligned}$$

We now estimate the two terms separately, and we start with A_ε^1 . In order to apply annealed maximal regularity theory, we smuggle in local averages at the scale ε , for $r \leq 2$,

$$\| \nabla v_{\varepsilon, g} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))} \leq \| [\nabla v_{\varepsilon, g}]_{2+\delta; \varepsilon} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))},$$

and we need to replace $[\cdot]_{2+\delta; \varepsilon}$ by local quadratic averages $[\cdot]_{2; \varepsilon}$. For that purpose, since $\nabla v_{\varepsilon, g}$ satisfies the elliptic equation (6.3), we appeal to Meyers' perturbative argument in the following form, for $0 < \delta \ll 1$,

$$\left(\int_{B_\varepsilon(x)} |\nabla v_{\varepsilon, g}|^{2+\delta} \right)^{\frac{1}{2+\delta}} \lesssim \left(\int_{B_{2\varepsilon}(x)} |\nabla v_{\varepsilon, g}|^2 \right)^{\frac{1}{2}} + \left(\int_{B_{2\varepsilon}(x)} |g|^{2+\delta} \right)^{\frac{1}{2+\delta}},$$

so that the above becomes

$$\| \nabla v_{\varepsilon, g} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))} \lesssim \| [\nabla v_{\varepsilon, g}]_{2; \varepsilon} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))} + \| [g]_{2+\delta; \varepsilon} \|_{L^r(\mathbb{R}^d)}.$$

We are now in position to apply the (ε -rescaled) annealed maximal regularity estimate of Theorem 3.1, which yields for all $1 < r \leq 2$,

$$\| \nabla v_{\varepsilon, g} \|_{L^r(\mathbb{R}^d; L^{2+\delta}(\Omega))} \lesssim_r \| [g]_{2+\delta; \varepsilon} \|_{L^r(\mathbb{R}^d)}. \quad (6.7)$$

Inserting this into (6.5), we deduce for all $1 < r < \frac{d}{d-\eta}$ and $0 < \eta, \delta \ll 1$,

$$A_\varepsilon^1(X, h, g) \lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \| [D_0 X^\sharp]_\infty \|_{L^r(\mathbb{R}^d; L^\infty(\Omega))} \| \langle \cdot \rangle^\eta h \|_{L^{r'}(\mathbb{R}^d)} \| [g]_{2+\delta; \varepsilon} \|_{L^r(\mathbb{R}^d)}. \quad (6.8)$$

We turn to the estimate on A_ε^2 . Acting with T and taking the Malliavin derivative in the equation (6.3) for $\nabla v_{\varepsilon, g}$, we find

$$-\nabla \cdot (\mathbf{a}^*(\frac{\cdot}{\varepsilon} + y)(D_0 T_y \nabla v_{\varepsilon, g})) = \nabla \cdot ((D_0 \mathbf{a}^*(\frac{\cdot}{\varepsilon} + y)) T_y \nabla v_{\varepsilon, g}). \quad (6.9)$$

Smuggling in local averages at the scale ε , using Meyers' argument and the annealed maximal regularity as above for equation (6.9), we find for all $1 < r \leq 2$,

$$\begin{aligned} & \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}} \\ & \leq \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B_\varepsilon(x) \times B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x')|^{2+\delta} dx' dy' \right)^{\frac{1}{2+\delta}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}} \\ & \lesssim_{r, \delta} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B_\varepsilon(x) \times B(y)} |(D_0 \mathbf{a}^*(\frac{x'}{\varepsilon} + y)) T_{y'} \nabla v_{\varepsilon, g}(x')|^{2+\delta} dx' dy' \right)^{\frac{1}{2+\delta}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}}. \end{aligned}$$

Recalling that $|D_0 \mathbf{a}^*(x)| \lesssim |\mathcal{C}_0(x)|$ and using the integrability condition (2.3), straightforward computations lead to

$$\begin{aligned} & \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}} \\ & \lesssim_{r, \delta} \|\nabla v_{\varepsilon, g}\|_{2+2\delta; \varepsilon} \|L^r(\mathbb{R}^d; L^{2+2\delta}(\Omega))\|. \end{aligned}$$

Using again Meyers' argument and the annealed maximal regularity similarly as above, we conclude for all $1 < r \leq 2$,

$$\left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left\| \left(\int_{B(y)} |D_0 T_{y'} \nabla v_{\varepsilon, g}(x)|^2 dy' \right)^{\frac{1}{2}} \right\|_{L^{2+\delta}(\Omega)}^r dx dy \right)^{\frac{1}{r}} \lesssim_{r, \delta} \|g\|_{2+2\delta; \varepsilon} \|L^r(\mathbb{R}^d)\|.$$

Inserting this into (6.6), we deduce for all $1 < r < \frac{d}{d-\eta}$ and $0 < \eta, \delta \ll 1$,

$$A_\varepsilon^2(X, h, g) \lesssim_{r, \eta, \delta} \varepsilon^{d-\eta} \|X\|_{L^\infty(\Omega)} \|\langle \cdot \rangle^\eta h\|_{L^{r'}(\mathbb{R}^d)} \|g\|_{2+2\delta; \varepsilon} \|L^r(\mathbb{R}^d)\|.$$

Combined with (6.8), this yields the claim (6.4).

Step 2. Approximation of ensemble-averaged field: for all $n < d$, $\frac{d}{d-n} \vee 2 < p < \infty$, and $\delta > 0$,

$$\|[\mathbb{E}[\nabla u_{\varepsilon, f}] - \nabla \bar{U}_\varepsilon^n[f]]\|_{2-\delta; \varepsilon} \|L^p(\mathbb{R}^d)\| \lesssim_{p, \delta} \varepsilon^n \|\langle \nabla \rangle^{2n-1} f\|_{L^p(\mathbb{R}^d)}, \quad (6.10)$$

while at critical order $n = d$ we have for all $\frac{d}{n} < p < \infty$ and $0 < \eta, \delta \ll 1$,

$$\|[\mathbb{E}[\nabla u_{\varepsilon, f}] - \nabla \bar{U}_\varepsilon^d[f]]\|_{2-\delta; \varepsilon} \|L^p(\mathbb{R}^d)\| \lesssim_{p, \eta, \delta} \varepsilon^{d-\eta} \|\langle \cdot \rangle^\eta \langle \nabla \rangle^{2d-1} f\|_{L^p(\mathbb{R}^d)}. \quad (6.11)$$

Starting point is Proposition 3.4, where identity (3.10) can be continued to higher orders in terms of the weak correctors defined in Theorem 5.1: the following equation holds in the distributional sense on $\mathbb{R}^d \times \Omega$ for all $1 \leq n \leq d$,

$$\begin{aligned} -\nabla \cdot \mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla (u_{\varepsilon, f} - \mathcal{F}_\varepsilon^n[\bar{U}_\varepsilon^n[f]]) &= \nabla \cdot \left(\sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \right) \\ &+ \nabla \cdot (\varepsilon^n (\mathbf{a} \varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n) (\frac{\cdot}{\varepsilon}) \nabla \nabla_{i_1 \dots i_n}^n \bar{U}_\varepsilon^n[f]). \quad (6.12) \end{aligned}$$

Given a test function $g \in C_c^\infty(\mathbb{R}^d)^d$, consider the solution $\nabla v_{\varepsilon, g}$ of the auxiliary problem (6.3). Testing the latter with $u_{\varepsilon, f} - \mathcal{F}_\varepsilon^n[\bar{U}_\varepsilon^n[f]]$, we find in $\mathcal{S}'(\Omega)$,

$$\int_{\mathbb{R}^d} g \cdot \nabla (u_{\varepsilon, f} - \mathcal{F}_\varepsilon^n[\bar{U}_\varepsilon^n[f]]) = - \int_{\mathbb{R}^d} \nabla v_{\varepsilon, g} \cdot \mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla (u_{\varepsilon, f} - \mathcal{F}_\varepsilon^n[\bar{U}_\varepsilon^n[f]]),$$

where both sides are indeed well-defined in $\mathcal{S}'(\Omega)$ in view of the result (6.4) of Step 1 (together with corresponding estimates for correctors $\varphi^n, \nabla\varphi^{n+1}$ with $n < d$). Next, testing equation (6.12) with $v_{\varepsilon,g}$, we deduce in $\mathcal{S}'(\Omega)$,

$$\begin{aligned} \int_{\mathbb{R}^d} g \cdot \nabla(u_{\varepsilon,f} - \mathcal{F}_{\varepsilon}^n[\bar{\mathcal{U}}_{\varepsilon}^n[f]]) &= \int_{\mathbb{R}^d} \nabla v_{\varepsilon,g} \cdot \sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \\ &\quad + \varepsilon^n \int_{\mathbb{R}^d} \nabla v_{\varepsilon,g} \cdot (\mathbf{a}\varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n)(\frac{\cdot}{\varepsilon}) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_{\varepsilon}^n[f]. \end{aligned}$$

Taking the expectation and noting that the definition of the higher-order two-scale expansion (3.7) yields

$$\mathbb{E}[\nabla \mathcal{F}_{\varepsilon}^n[\bar{\mathcal{U}}_{\varepsilon}^n[f]]] = \nabla \bar{\mathcal{U}}_{\varepsilon}^n[f],$$

we find

$$\begin{aligned} \int_{\mathbb{R}^d} g \cdot (\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_{\varepsilon}^n[f]) &= \mathbb{E} \left[\int_{\mathbb{R}^d} \nabla v_{\varepsilon,g} \cdot \sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \right] \\ &\quad + \mathbb{E} \left[\varepsilon^n \int_{\mathbb{R}^d} \nabla v_{\varepsilon,g} \cdot (\mathbf{a}\varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n)(\frac{\cdot}{\varepsilon}) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_{\varepsilon}^n[f] \right]. \end{aligned}$$

We focus on the proof of (6.11) for $n = d$, while the proof of (6.10) is similar. We analyze the two right-hand side terms in the above identity separately. For the first term, we use Hölder's inequality and we appeal to the annealed maximal regularity theory of Theorem 3.1 in form of $\|[\nabla v_{\varepsilon,g}]_{2;\varepsilon}\|_{L^r(\mathbb{R}^d; L^2(\Omega))} \lesssim \|g\|_{2;\varepsilon}\|_{L^r(\mathbb{R}^d)}$. For the second term, we apply the result (6.4) of Step 1. For all $1 < r < \frac{d}{d-\eta}$ and $0 < \eta, \delta \ll 1$, we deduce

$$\begin{aligned} \left| \int_{\mathbb{R}^d} g \cdot (\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_{\varepsilon}^d[f]) \right| &\lesssim_{r,\eta,\delta} \sum_{k=2}^d \sum_{l=d+2-k}^d \varepsilon^{k+l-2} \|g\|_{2;\varepsilon}\|_{L^r(\mathbb{R}^d)} \|\nabla^k \tilde{u}_f^l\|_{L^{r'}(\mathbb{R}^d)} \\ &\quad + \varepsilon^{d-\eta} \|g\|_{2+\delta;\varepsilon}\|_{L^r(\mathbb{R}^d)} \| \langle \cdot \rangle^{\eta} \nabla^{d+1} \bar{\mathcal{U}}_{\varepsilon}^d[f] \|_{L^{r'}(\mathbb{R}^d)}. \end{aligned}$$

Further using standard weighted L^p regularity for the higher-order homogenized equations in Definition 3.3, we are led to

$$\left| \int_{\mathbb{R}^d} g \cdot (\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_{\varepsilon}^d[f]) \right| \lesssim_{r,\eta,\delta} \varepsilon^{d-\eta} \|g\|_{2+\delta;\varepsilon}\|_{L^r(\mathbb{R}^d)} \| \langle \cdot \rangle^{\eta} \langle \nabla \rangle^{2d-1} f \|_{L^{r'}(\mathbb{R}^d)}.$$

Taking the supremum over g , this yields the claim (6.11) for the ensemble-averaged field.

Step 3. Approximation of ensemble-averaged flux: for all $n < d$, $\frac{d}{d-n} \vee 2 < p < \infty$, and $\delta > 0$,

$$\| [\mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f}] - \bar{\mathcal{A}}_{\varepsilon}^n \nabla \bar{\mathcal{U}}_{\varepsilon}^n[f]]_{2-\delta;\varepsilon} \|_{L^p(\mathbb{R}^d)} \lesssim_{p,\delta} \varepsilon^n \| \langle \nabla \rangle^{2n-1} f \|_{L^p(\mathbb{R}^d)}, \quad (6.13)$$

while at critical order $n = d$ we have for all $\frac{d}{\eta} < p < \infty$ and $0 < \eta, \delta \ll 1$,

$$\| [\mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f}] - \bar{\mathcal{A}}_{\varepsilon}^d \nabla \bar{\mathcal{U}}_{\varepsilon}^d[f]]_{2-\delta;\varepsilon} \|_{L^p(\mathbb{R}^d)} \lesssim_{p,\eta,\delta} \varepsilon^{d-\eta} \| \langle \cdot \rangle^{\eta} \langle \nabla \rangle^{2d-1} f \|_{L^p(\mathbb{R}^d)}. \quad (6.14)$$

Starting point is the same identity (6.12) as in Step 1, which we now test with the unique almost sure gradient solution $\nabla w_{\varepsilon,g}$ in $L^2(\mathbb{R}^d)^d$ of the auxiliary problem

$$-\nabla \cdot \mathbf{a}^*(\frac{\cdot}{\varepsilon}) \nabla w_{\varepsilon,g} = \nabla \cdot \mathbf{a}^*(\frac{\cdot}{\varepsilon}) g, \quad \text{in } \mathbb{R}^d.$$

This yields the following identity in $\mathcal{S}'(\Omega)$,

$$\begin{aligned} \int_{\mathbb{R}^d} g \cdot \mathbf{a}(\cdot/\varepsilon) (\nabla u_{\varepsilon,f} - \nabla \mathcal{F}_\varepsilon^n[\bar{\mathcal{U}}_\varepsilon^n[f]]) &= \int_{\mathbb{R}^d} \nabla w_{\varepsilon,g} \cdot \sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \\ &+ \varepsilon^n \int_{\mathbb{R}^d} \nabla w_{\varepsilon,g} \cdot (\mathbf{a}\varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n)(\cdot/\varepsilon) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_\varepsilon^n[f]. \end{aligned}$$

Taking the expectation and noting that the definitions of the higher-order two-scale expansion (3.7) and of higher-order effective tensors (5.4) yield

$$\mathbb{E} [\mathbf{a}(\cdot/\varepsilon) \nabla \mathcal{F}_\varepsilon^n[\bar{\mathcal{U}}_\varepsilon^n[f]]] = \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{\mathcal{U}}_\varepsilon^n[f] + \mathbb{E} [\varepsilon^n (\mathbf{a}\varphi_{i_1 \dots i_n}^n)(\cdot/\varepsilon) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_\varepsilon^n[f]],$$

the above identity becomes

$$\begin{aligned} \int_{\mathbb{R}^d} g \cdot (\mathbb{E}[\mathbf{a}(\cdot/\varepsilon) \nabla u_{\varepsilon,f}] - \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{\mathcal{U}}_\varepsilon^n[f]) \\ = \mathbb{E} \left[\int_{\mathbb{R}^d} \nabla w_{\varepsilon,g} \cdot \sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^k \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{u}_f^l \right] \\ + \mathbb{E} \left[\varepsilon^n \int_{\mathbb{R}^d} \nabla w_{\varepsilon,g} \cdot (\mathbf{a}\varphi_{i_1 \dots i_n}^n - \sigma_{i_1 \dots i_n}^n)(\cdot/\varepsilon) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_\varepsilon^n[f] \right] \\ + \mathbb{E} \left[\varepsilon^n \int_{\mathbb{R}^d} g \cdot (\mathbf{a}\varphi_{i_1 \dots i_n}^n)(\cdot/\varepsilon) \nabla \nabla_{i_1 \dots i_n}^n \bar{\mathcal{U}}_\varepsilon^n[f] \right]. \end{aligned}$$

Noting that the result (6.4) of Step 1 also holds for $\nabla v_{\varepsilon,g}$ replaced by $\nabla w_{\varepsilon,g}$, and arguing as in Step 2, the claim (6.13)–(6.14) easily follows.

Step 4. Conclusion.

It remains to replace sub-quadratic averages $[\cdot]_{2-\delta}$ in (6.10)–(6.11) and in (6.13)–(6.14) by quadratic averages. We focus on the proof for the ensemble-averaged field, while the proof is identical for the flux. For that purpose, we use the following interpolation inequality: for all $h \in C_c^\infty(\mathbb{R}^d)$ and $0 < \delta \leq 1$, setting $\kappa_\delta := (1 - \frac{1}{2}\delta^2)(1 + \delta)^{-1}$,

$$[h]_2 \leq [h]_{2-\delta^2}^{\kappa_\delta} [h]_{2+\delta}^{1-\kappa_\delta},$$

which yields by Hölder's inequality, for all p ,

$$\|[h]_2\|_{L^p(\mathbb{R}^d)} \leq \|[h]_{2-\delta^2}\|_{L^p(\mathbb{R}^d)}^{\kappa_\delta} \|[h]_{2+\delta}\|_{L^p(\mathbb{R}^d)}^{1-\kappa_\delta}.$$

Applying this inequality to $h = \mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_\varepsilon^n[f]$, appealing to (6.10)–(6.11) with δ replaced by δ^2 , and using (6.7) for $u_{\varepsilon,f}$ together with the standard L^p regularity for the higher-order homogenized equations in form of

$$\begin{aligned} \|[\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_\varepsilon^n[f]]_{2+\delta;\varepsilon}\|_{L^p(\mathbb{R}^d)} &\leq \|[\nabla u_{\varepsilon,f}]_{2+\delta;\varepsilon}\|_{L^p(\mathbb{R}^d; L^2(\Omega))} + \|[\nabla \bar{\mathcal{U}}_\varepsilon^n[f]]_{2+\delta;\varepsilon}\|_{L^p(\mathbb{R}^d)} \\ &\lesssim_p \| \langle \nabla \rangle^{n-1} f \|_{2+\delta;\varepsilon} \|_{L^p(\mathbb{R}^d)}, \end{aligned}$$

we deduce for all $n < d$, $\frac{d}{d-n} \vee 2 < p < \infty$, and $\delta > 0$,

$$\|[\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_\varepsilon^n[f]]_{2;\varepsilon}\|_{L^p(\mathbb{R}^d)} \lesssim_{p,\delta} \varepsilon^{\kappa_\delta n} \| \langle \nabla \rangle^{2n-1} f \|_{L^p(\mathbb{R}^d)}, \quad (6.15)$$

and similarly, at critical order $n = d$, for all $\frac{d}{\eta} < p < \infty$ and $0 < \eta, \delta \ll 1$,

$$\|[\mathbb{E}[\nabla u_{\varepsilon,f}] - \nabla \bar{\mathcal{U}}_\varepsilon^d[f]]_{2;\varepsilon}\|_{L^p(\mathbb{R}^d)} \lesssim_{p,\eta,\delta} \varepsilon^{\kappa_\delta(d-\eta)} \| \langle \cdot \rangle^\eta \langle \nabla \rangle^{2d-1} f \|_{L^p(\mathbb{R}^d)}. \quad (6.16)$$

Noting that $\kappa_\delta \geq 1 - 2\delta$, the conclusion (6.2) for $n = d$ follows from (6.16) up to renaming η, δ . Regarding the conclusion (6.1), it is first deduced from (6.15) with suboptimal rate $\varepsilon^{(1-2\delta)n}$, while the optimal rate ε^n is easily recovered by interpolation. \square

APPENDIX A. INTRINSIC DESCRIPTION OF WEAK FLUCTUATIONS

Aside from the Bourgain–Spencer conjecture on ensemble averages, another natural question concerns the intrinsic description of fluctuations $\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]$ in the weak sense of $\mathcal{S}'(\Omega)$. The proof of Theorem 2 ensures that such “weak” fluctuations are (almost) of order $O(\varepsilon^d)$: for all $f \in C_c^\infty(\mathbb{R}^d)^d$, $X \in \mathcal{R}(\Omega)$, and $\eta > 0$,

$$\left\| \left[\mathbb{E}[X(\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}])] \right]_{2,\varepsilon} \right\|_{L^\infty(\mathbb{R}^d)} \lesssim_{X,f,\eta} \varepsilon^{d-\eta},$$

This contrasts with the usual CLT scaling result, e.g. [14], which states that fluctuations are of order $O(\varepsilon^{d/2})$ in the usual strong sense: for all $f, g \in C_c^\infty(\mathbb{R}^d)^d$ and $q < \infty$,

$$\mathbb{E} \left[\left| \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]) \right|^q \right]^{\frac{1}{q}} \lesssim_{f,g,q} \varepsilon^{\frac{d}{2}}.$$

Taking inspiration from our recent work with Gloria and Otto [13, 14] on “strong” fluctuations, we show that “weak” fluctuations can similarly be described intrinsically (almost) to relative order $O(\varepsilon^{d/2})$. This result only involves the standard strong correctors $\{\varphi^n\}_{1 \leq n \leq \ell}$, and we do not know whether this could be improved.

Theorem A.1 (Two-scale expansion for “weak” fluctuations). *For all $f \in C_c^\infty(\mathbb{R}^d)^d$, there exists an ℓ th-order differential operator $\Xi_\varepsilon^{\circ,\ell}[\nabla \cdot]$ with ε -rescaled stationary random coefficients expressed in terms of correctors $\{\varphi^n\}_{1 \leq n \leq \ell}$, see Definition A.3 below, such that there holds for all $X \in \mathcal{R}(\Omega)$, $g \in C_c^\infty(\mathbb{R}^d)^d$, and $\eta > 0$,*

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} g \cdot \varepsilon^{-d} \mathbb{E}[X(\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}])] - \int_{\mathbb{R}^d} (\nabla \bar{U}_\varepsilon^\ell)^*[g] \cdot \varepsilon^{-d} \mathbb{E}[X \Xi_\varepsilon^{\circ,\ell}[\nabla \bar{U}_\varepsilon^\ell f]] \right| \\ & + \left| \int_{\mathbb{R}^d} g \cdot \varepsilon^{-d} \mathbb{E}[X(\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f} - \mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f}])] \right. \\ & \quad \left. - \int_{\mathbb{R}^d} (\text{Id} + \bar{\mathcal{A}}_\varepsilon^\ell \nabla \bar{U}_\varepsilon^\ell)^*[g] \cdot \varepsilon^{-d} \mathbb{E}[X \Xi_\varepsilon^{\circ,\ell}[\nabla \bar{U}_\varepsilon^\ell f]] \right| \\ & \lesssim_\eta \varepsilon^{\frac{d}{2}-\eta} \|[DX]_2\|_{L^1(\mathbb{R}^d; L^2(\Omega))} \|\mu_d \langle \nabla \rangle^{3\ell-2} g\|_{(L^4 \cap L^\infty)(\mathbb{R}^d)} \|\mu_d \langle \nabla \rangle^{3\ell-2} f\|_{(L^4 \cap L^\infty)(\mathbb{R}^d)}, \end{aligned}$$

where the weight μ_d is given by

$$\mu_d(x) := \begin{cases} \langle x \rangle^{\frac{1}{2}} & : d \text{ odd}, \\ \log(2 + |x|)^{\frac{1}{2}} & : d \text{ even}. \end{cases} \quad \diamond$$

A.1. Intrinsic description of strong fluctuations. As first noticed by Gu and Mourrat [20], random fluctuations of $\nabla u_{\varepsilon,f}$ are not described by the usual two-scale expansion: the Gaussian limit of the centered rescaled observable $\varepsilon^{-d/2} \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}])$ differs from the limit of $\varepsilon^{-d/2} \int_{\mathbb{R}^d} g \cdot \nabla \varphi_i^1(\frac{\cdot}{\varepsilon}) \nabla_i \bar{u}_f^1$. In [13], with Gloria and Otto, we establish the first *pathwise* theory of fluctuations, providing an intrinsic description of fluctuations based on a suitably modified notion of two-scale expansion (see also the related heuristics

in [20]). More precisely, fluctuations of $\nabla u_{\varepsilon,f}$ coincide to first order with fluctuations of some deterministic Helmholtz projection of the corresponding *homogenization commutator*

$$\Xi_{\varepsilon}^1[\nabla u_{\varepsilon,f}] := (\mathbf{a}(\frac{\cdot}{\varepsilon}) - \bar{\mathbf{a}}^1) \nabla u_{\varepsilon,f},$$

for which the usual two-scale expansion is shown to be accurate: leading-order fluctuations are then governed by the so-called *standard* homogenization commutator,

$$\Xi_{\varepsilon}^{\circ,1}[\nabla \bar{u}_f^1] := (\mathbf{a}(\frac{\cdot}{\varepsilon}) - \bar{\mathbf{a}}^1) (\nabla \varphi_i^1(\frac{\cdot}{\varepsilon}) + e_i) \nabla_i \bar{u}_f^1.$$

We emphasize that the two-scale expansion of $\nabla u_{\varepsilon,f}$ via its commutator differs from the naive two-scale expansion (1.5).

As we showed in [14], this theory is naturally extended to higher orders. For $0 \leq n \leq \ell$ we define the following n th-order homogenization commutator, as inspired by the higher-order constitutive law (3.8),

$$\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}] := (\mathbf{a}(\frac{\cdot}{\varepsilon}) - \bar{\mathcal{A}}_{\varepsilon}^n) \nabla u_{\varepsilon,f}. \quad (\text{A.1})$$

We first recall that higher-order fluctuations of $\nabla u_{\varepsilon,f}$ coincide with fluctuations of some deterministic Helmholtz projection of this commutator; see [14, proof of Proposition 3.2].

Lemma A.2 (Reduction to commutators; [14]). *For all $1 \leq n \leq \ell$ and $f, g \in C_c^{\infty}(\mathbb{R}^d)^d$, we have*

$$\begin{aligned} & \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]) - \int_{\mathbb{R}^d} (\nabla \bar{U}_{\varepsilon}^n)^*[g] \cdot (\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}]]) \\ &= \int_{\mathbb{R}^d} \left(\sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^{*,k} \nabla \nabla_{i_1 \dots i_{k-1}}^{k-1} \tilde{v}_g^l \right) \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]), \quad (\text{A.2}) \end{aligned}$$

which entails for all $q < \infty$,

$$\begin{aligned} & \mathbb{E} \left[\left| \varepsilon^{-\frac{d}{2}} \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon} - \mathbb{E}[\nabla u_{\varepsilon}]) \right. \right. \\ & \quad \left. \left. - \varepsilon^{-\frac{d}{2}} \int_{\mathbb{R}^d} (\nabla \bar{U}_{\varepsilon}^n)^*[g] \cdot (\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}]] \right|^q \right]^{\frac{1}{q}} \lesssim_{f,g,p} \varepsilon^n. \quad \diamond \end{aligned}$$

Next, we recall the definition of the n th-order standard commutator as the suitable two-scale expansion of the commutator $\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}]$. It is obtained by inserting the n th-order two-scale expansion $\mathcal{F}_{\varepsilon}^n[\cdot]$ into $\Xi_{\varepsilon}^n[\nabla \cdot]$ and by truncating the obtained differential operator to order n ; see [14, Section 3].

Definition A.3 (Standard higher-order homogenization commutators). *For $0 \leq n \leq \ell$, the n th-order standard commutator $\Xi_{\varepsilon}^{\circ,n}[\nabla \cdot]$ is an n th-order differential operator with ε -rescaled stationary random coefficients in $H_{\text{loc}}^{1-n}(\mathbb{R}^d; L^2(\Omega))$, expressed in terms of correctors $\{\varphi^k\}_{1 \leq k \leq n}$. Applied to $\bar{w} \in C_c^{\infty}(\mathbb{R}^d)$, it is defined by*

$$\Xi_{\varepsilon}^{\circ,n}[\nabla \bar{w}](x) := \Xi_{\varepsilon}^n[\nabla \mathcal{F}_{\varepsilon}^n[T_x^n \bar{w}]](x),$$

where $T_x^n \bar{w}$ is the n th-order Taylor polynomial of \bar{w} at x . Note that $\mathbb{E}[\Xi_{\varepsilon}^{\circ,n}[\nabla \cdot]] = 0$. \diamond

Our main result in [14, Corollary 1] then shows that the two-scale expansion for commutators $\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_{\varepsilon}^n[\nabla u_{\varepsilon,f}]] \approx \Xi_{\varepsilon}^{\circ,n}[\nabla \bar{U}_{\varepsilon}^n[f]]$ is accurate to order $O(\varepsilon^n)$ in the fluctuation scaling. Due to the maximal number $\ell = \lceil \frac{d}{2} \rceil$ of stationary “strong” correctors,

this saturates at order $O(\varepsilon^{d/2})$, which is expected to be optimal as fluctuations become non-Gaussian at that order [14].

Proposition A.4 (Two-scale expansion of commutators; [14]). *For all $f, g \in C_c^\infty(\mathbb{R}^d)$ and $q < \infty$, we have*

$$\begin{aligned} \mathbb{E} \left[\left| \varepsilon^{-\frac{d}{2}} \int_{\mathbb{R}^d} g \cdot (\Xi_\varepsilon^\ell[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^\ell[\nabla u_{\varepsilon,f}]]) - \varepsilon^{-\frac{d}{2}} \int_{\mathbb{R}^d} g \cdot \Xi_\varepsilon^{\circ,\ell}[\nabla \bar{U}_\varepsilon^\ell[f]] \right|^q \right]^{\frac{1}{q}} \\ \lesssim_{f,g,q} \varepsilon^{\frac{d}{2}} \times \begin{cases} 1 & : d \text{ odd}, \\ |\log \varepsilon|^{\frac{1}{2}} & : d \text{ even}. \end{cases} \quad \diamond \end{aligned}$$

A.2. Proof of Theorem A.1. We split the proof into two steps.

Step 1. Reduction to commutators: for all $1 \leq n \leq \ell$, $p > 1$, and $\delta > 0$,

$$\begin{aligned} \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]) \right] - \mathbb{E} \left[X \int_{\mathbb{R}^d} (\nabla \bar{U}_\varepsilon^n)^*[g] \cdot (\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}]]) \right] \right| \\ \lesssim_{p,\delta} \varepsilon^{n+\frac{d}{p}} \|DX\|_{L^p(\mathbb{R}^d; L^{1+\delta}(\Omega))} \|\langle \nabla \rangle^{2n-1} g\|_{L^{2p'}(\mathbb{R}^d)} \|f\|_{L^{2p'}(\mathbb{R}^d)}, \quad (\text{A.3}) \end{aligned}$$

and similarly, for fluctuations of the flux,

$$\begin{aligned} \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \cdot (\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f} - \mathbb{E}[\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f}]) \right] \right. \\ \left. - \mathbb{E} \left[X \int_{\mathbb{R}^d} (\text{Id} + \bar{\mathcal{A}}_\varepsilon^n \nabla \bar{U}_\varepsilon^n)^*[g] \cdot (\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}]]) \right] \right| \\ \lesssim_{p,q} \varepsilon^{n+\frac{d}{p}} \|DX\|_{L^p(\mathbb{R}^d; L^{1+\delta}(\Omega))} \|\langle \nabla \rangle^{3n-2} g\|_{L^{2p'}(\mathbb{R}^d)} \|f\|_{L^{2p'}(\mathbb{R}^d)}. \quad (\text{A.4}) \end{aligned}$$

Noting that $\mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla u_{\varepsilon,f} = \Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] + \bar{\mathcal{A}}_\varepsilon^n \nabla u_{\varepsilon,f}$, the claim (A.4) follows from (A.3), and it remains to prove the latter. Given a test function $X \in \mathcal{S}(\Omega)$, starting from identity (A.2) and appealing to the Helffer–Sjöstrand identity, see Lemma 4.2, together with Hölder’s inequality, we find for all $1 \leq p, q \leq \infty$,

$$\begin{aligned} \mathbb{E} \left[X \int_{\mathbb{R}^d} g \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]) \right] - \mathbb{E} \left[X \int_{\mathbb{R}^d} (\nabla \bar{U}_\varepsilon^n)^*[g] \cdot (\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}]]) \right] \\ = \mathbb{E} \left[X \int_{\mathbb{R}^d} H_\varepsilon^n \cdot (\nabla u_{\varepsilon,f} - \mathbb{E}[\nabla u_{\varepsilon,f}]) \right] \\ \leq \|DX\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \left\| \int_{\mathbb{R}^d} H_\varepsilon^n \cdot D \nabla u_{\varepsilon,f} \right\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))}, \quad (\text{A.5}) \end{aligned}$$

where we have set for abbreviation

$$H_\varepsilon^n := \sum_{k=2}^n \sum_{l=n+2-k}^n \varepsilon^{k+l-2} \bar{\mathbf{a}}_{i_1 \dots i_{k-1}}^{*,k} \nabla \nabla^{k-1} \bar{v}_{i_1 \dots i_{k-1}}^l g.$$

Consider the unique almost sure gradient solution ∇h_ε^n in $L^2(\mathbb{R}^d)^d$ of the auxiliary equation

$$-\nabla \cdot \mathbf{a}^*(\frac{\cdot}{\varepsilon}) \nabla h_\varepsilon^n = \nabla \cdot H_\varepsilon^n, \quad \text{in } \mathbb{R}^d,$$

and note that taking the Malliavin derivative in the equation for $\nabla u_{\varepsilon,f}$ yields

$$-\nabla \cdot \mathbf{a}(\frac{\cdot}{\varepsilon}) \nabla D_z u_{\varepsilon,f} = \nabla \cdot (D_z \mathbf{a}(\frac{\cdot}{\varepsilon})) \nabla u_{\varepsilon,f}.$$

Then computing

$$\int_{\mathbb{R}^d} H_\varepsilon^n \cdot D_z \nabla u_{\varepsilon,f} = - \int_{\mathbb{R}^d} \nabla h_\varepsilon^n \cdot \mathbf{a}(\frac{\cdot}{\varepsilon}) D_z \nabla u_{\varepsilon,f} = \int_{\mathbb{R}^d} \nabla h_\varepsilon^n \cdot (D_z \mathbf{a}(\frac{\cdot}{\varepsilon})) \nabla u_{\varepsilon,f},$$

and recalling that $|D_z \mathbf{a}(\frac{x}{\varepsilon})| \lesssim |\mathcal{C}_0(z - \frac{x}{\varepsilon})|$, we obtain from Hölder's inequality and from the integrability condition (2.3),

$$\begin{aligned} & \left\| \int_{\mathbb{R}^d} H_\varepsilon^n \cdot D \nabla u_{\varepsilon,f} \right\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} \\ & \lesssim \left(\int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} [\mathcal{C}_0]_\infty(\cdot - \frac{x}{\varepsilon}) \|\nabla h_\varepsilon^n \nabla u_{\varepsilon,f}\|_{1;\varepsilon}(x) \, dx \right|^{p'} \right)^{\frac{1}{p'}} \\ & \lesssim \varepsilon^{\frac{d}{p}} \|\nabla h_\varepsilon^n \nabla u_{\varepsilon,f}\|_{1;\varepsilon} \| \cdot \|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))}. \end{aligned}$$

Using again Hölder's inequality, together with the annealed L^p regularity of Theorem 3.1, this becomes for $p, q > 1$,

$$\begin{aligned} \left\| \int_{\mathbb{R}^d} H_\varepsilon^n \cdot D \nabla u_{\varepsilon,f} \right\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} & \lesssim \varepsilon^{\frac{d}{p}} \|\nabla h_\varepsilon^n\|_{2;\varepsilon} \| \cdot \|_{L^{2p'}(\mathbb{R}^d; L^{2q'}(\Omega))} \|\nabla u_{\varepsilon,f}\|_{2;\varepsilon} \| \cdot \|_{L^{2p'}(\mathbb{R}^d; L^{2q'}(\Omega))} \\ & \lesssim_{p,q} \varepsilon^{\frac{d}{p}} \|H_\varepsilon^n\|_{L^{2p'}(\mathbb{R}^d)} \|f\|_{L^{2p'}(\mathbb{R}^d)}. \end{aligned}$$

Inserting this into (A.5), and applying the standard constant-coefficient maximal L^p regularity for higher-order homogenized equations, the claim (A.3) follows.

Step 2. Two-scale expansion of commutators: for all $1 \leq n \leq \ell$, $1 < p < \frac{2d}{2d-1}$, and $\delta > 0$,

$$\begin{aligned} & \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \cdot \left((\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}]] - \Xi_\varepsilon^{\circ,n}[\nabla \bar{U}_\varepsilon^n[f]]) \right) \right] \right| \\ & \lesssim_{p,\delta} \varepsilon^{\frac{d}{p}+n} \mu_{d,n}(\frac{1}{\varepsilon}) \|DX\|_{L^p(\mathbb{R}^d; L^{1+\delta}(\Omega))} \\ & \quad \times \|\mu_{d,n} \langle \nabla \rangle^n g\|_{(L^4 \cap L^{2p'}) (\mathbb{R}^d)} \|\mu_{d,n} \langle \nabla \rangle^{3n-2} f\|_{(L^4 \cap L^{2p'}) (\mathbb{R}^d)}, \end{aligned} \quad (\text{A.6})$$

where the weight $\mu_{d,n}$ is given by $\mu_{d,n} = 1$ if $n < \ell$, and by $\mu_{d,n} = \mu_d$ if $n = \ell$. Combined with the result (A.3)–(A.4) of Step 1, and with the standard constant-coefficient maximal L^p regularity for homogenized equations, this yields the conclusion.

Recalling that $\mathbb{E}[\Xi_\varepsilon^{\circ,n}[\nabla \bar{U}_\varepsilon^n[f]]] = 0$, appealing to the Helffer–Sjöstrand identity, see Lemma 4.2(iii), together with Hölder's inequality, we find for all $1 \leq p, q \leq \infty$,

$$\begin{aligned} & \left| \mathbb{E} \left[X \int_{\mathbb{R}^d} g \cdot \left((\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \mathbb{E}[\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}]] - \Xi_\varepsilon^{\circ,n}[\nabla \bar{U}_\varepsilon^n[f]]) \right) \right] \right| \\ & \leq \|DX\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \left\| \int_{\mathbb{R}^d} g \cdot D(\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \Xi_\varepsilon^{\circ,n}[\nabla \bar{U}_\varepsilon^n[f]]) \right\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))}. \end{aligned}$$

Repeating the analysis in [14, Theorem 1(ii)] yields the following estimate for the last right-hand side term: for all $1 \leq n \leq \ell$, $1 < p < \frac{d}{n-1}$, and $q > 1$,

$$\begin{aligned} & \left\| \int_{\mathbb{R}^d} g \cdot D(\Xi_\varepsilon^n[\nabla u_{\varepsilon,f}] - \Xi_\varepsilon^{\circ,n}[\nabla \bar{U}_\varepsilon^n f]) \right\|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} \\ & \lesssim_{p,q} \varepsilon^{\frac{d}{p}+n} \mu_{d,n}(\tfrac{1}{\varepsilon}) \|\mu_{d,n}(|\cdot|) \langle \nabla \rangle^n g\|_{(L^{\frac{2dp'}{d+(n-1)p'} \cap L^{2p'}})(\mathbb{R}^d)} \\ & \quad \times \|\mu_{d,n}(|\cdot|) \langle \nabla \rangle^{2n-1}(f, \nabla \bar{U}_\varepsilon^n f)\|_{(L^{\frac{2dp'}{d+(n-1)p'} \cap L^{2p'}})(\mathbb{R}^d)}. \end{aligned}$$

Appealing to the standard (weighted) constant-coefficient maximal L^p regularity for higher-order homogenized equations, the claim (A.6) follows. \square

ACKNOWLEDGEMENTS

The author thanks Antoine Gloria, Marius Lemm, and Felix Otto for many motivating and inspiring discussions around the Bourgain–Spencer conjecture, and acknowledges financial support from the CNRS-Momentum program.

REFERENCES

- [1] A. Abdulle and T. Pouchon. Effective models and numerical homogenization for wave propagation in heterogeneous media on arbitrary timescales. *Found. Comput. Math.*, 20:1505–1547, 2020.
- [2] A. Andersson, R. Kruse, and S. Larsson. Duality in refined Sobolev-Malliavin spaces and weak approximation of SPDE. *Stoch. Partial Differ. Equ. Anal. Comput.*, 4(1):113–149, 2016.
- [3] S. Armstrong, T. Kuusi, and J.-C. Mourrat. *Quantitative stochastic homogenization and large-scale regularity*, volume 352 of *Grundlehren der Mathematischen Wissenschaften*. Springer, 2019.
- [4] S. N. Armstrong, T. Kuusi, and J.-C. Mourrat. The additive structure of elliptic homogenization. *Invent. Math.*, 208:999–1154, 2017.
- [5] S. N. Armstrong and J.-C. Mourrat. Lipschitz regularity for elliptic equations with random coefficients. *Arch. Ration. Mech. Anal.*, 219(1):255–348, 2016.
- [6] S. N. Armstrong and C. K. Smart. Quantitative stochastic homogenization of convex integral functionals. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(2):423–481, 2016.
- [7] P. Bella, B. Fehrman, J. Fischer, and F. Otto. Stochastic homogenization of linear elliptic equations: Higher-order error estimates in weak norms via second-order correctors. *SIAM J. Math. Anal.*, 49(6):4658–4703, 2017.
- [8] J. Bourgain. On a homogenization problem. *J. Stat. Phys.*, 172(2):314–320, 2018.
- [9] M. Duerinckx. On the size of chaos via Glauber calculus in the classical mean-field dynamics. *Commun. Math. Phys.*, 2021. In press.
- [10] M. Duerinckx, J. Fischer, and A. Gloria. Scaling limit of the homogenization commutator for Gaussian coefficient fields. Preprint, arXiv:1910.04088.
- [11] M. Duerinckx, A. Gloria, and M. Lemm. A remark on a surprising result by Bourgain in homogenization. *Comm. Part. Diff. Eq.*, 44(2):1345–1357, 2019.
- [12] M. Duerinckx, A. Gloria, and F. Otto. Robustness of the pathwise structure of fluctuations in stochastic homogenization. *Probab. Theory Related Fields*, 178:531–566, 2020.
- [13] M. Duerinckx, A. Gloria, and F. Otto. The structure of fluctuations in stochastic homogenization. *Comm. Math. Phys.*, 377:259–306, 2020.
- [14] M. Duerinckx and F. Otto. Higher-order pathwise theory of fluctuations in stochastic homogenization. *Stoch. Partial Differ. Equ. Anal. Comput.*, 8:625–692, 2020.
- [15] M. Duerinckx and C. Shirley. A new spectral analysis of stationary random Schrödinger operators. Preprint, arXiv:2004.12025.
- [16] A. Gloria, S. Neukamm, and F. Otto. Quantitative estimates in stochastic homogenization for correlated coefficient fields. Preprint, arXiv:1910.05530.

- [17] A. Gloria, S. Neukamm, and F. Otto. An optimal quantitative two-scale expansion in stochastic homogenization of discrete elliptic equations. *M2AN Math. Model. Numer. Anal.*, 48(2):325–346, 2014.
- [18] A. Gloria, S. Neukamm, and F. Otto. A regularity theory for random elliptic operators. *Milan J. Math.*, 88(1):99–170, 2020.
- [19] Y. Gu. High order correctors and two-scale expansions in stochastic homogenization. *Probab. Theory Related Fields*, 169(3-4):1221–1259, 2017.
- [20] Y. Gu and J.-C. Mourrat. Scaling limit of fluctuations in stochastic homogenization. *Multiscale Model. Simul.*, 14(1):452–481, 2016.
- [21] H. Holden, B. Øksendal, J. Ubøe, and T. Zhang. *Stochastic partial differential equations. A modeling, white noise functional approach*. Probability and its Applications. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [22] V. V. Jikov, S. M. Kozlov, and O. A. Oleĭnik. *Homogenization of differential operators and integral functionals*. Springer-Verlag, Berlin, 1994.
- [23] J. Kim and M. Lemm. On the Averaged Green’s Function of an Elliptic Equation with Random Coefficients. *Arch. Ration. Mech. Anal.*, 234:1121–1166, 2019.
- [24] S. M. Kozlov. The averaging of random operators. *Mat. Sb. (N.S.)*, 109(151)(2):188–202, 327, 1979.
- [25] P. C. Kunstmann and L. Weis. Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^∞ -functional calculus. In *Functional analytic methods for evolution equations*, volume 1855 of *Lecture Notes in Math.*, pages 65–311. Springer, Berlin, 2004.
- [26] J.-C. Mourrat and F. Otto. Correlation structure of the corrector in stochastic homogenization. *Ann. Probab.*, 44(5):3207–3233, 2016.
- [27] I. Nourdin and G. Peccati. *Normal approximations with Malliavin calculus. From Stein’s method to universality*, volume 192 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012.
- [28] D. Nualart. *The Malliavin calculus and related topics*. Springer-Verlag, Berlin, second edition, 2006.
- [29] G. C. Papanicolaou and S. R. S. Varadhan. Boundary value problems with rapidly oscillating random coefficients. In *Random fields, Vol. I, II (Esztergom, 1979)*, volume 27 of *Colloq. Math. Soc. János Bolyai*, pages 835–873. North-Holland, Amsterdam, 1981.
- [30] G. Peccati and M. Reitzner, editors. *Stochastic analysis for Poisson point processes*, volume 7 of *Bocconi & Springer Series*. Bocconi University Press, Springer, 2016.
- [31] I. M. Sigal. Homogenization problem. Unpublished preprint.

(Mitia Duerinckx) UNIVERSITÉ PARIS-SACLAY, CNRS, LABORATOIRE DE MATHÉMATIQUES D’ORSAY, 91400 ORSAY, FRANCE & UNIVERSITY OF CALIFORNIA, LOS ANGELES, DEPARTMENT OF MATHEMATICS, CA 90095, USA & UNIVERSITÉ LIBRE DE BRUXELLES, DÉPARTEMENT DE MATHÉMATIQUE, 1050 BRUSSELS, BELGIUM

Email address: mitia.duerinckx@u-psud.fr