CHERENKOV RADIATION WITH MASSIVE BOSONS AND QUANTUM FRICTION

MITIA DUERINCKX AND CHRISTOPHER SHIRLEY

ABSTRACT. This work is devoted to several translation-invariant models in non-relativistic quantum field theory (QFT), describing a non-relativistic quantum particle interacting with a quantized relativistic field of bosons. In this setting, we aim at the rigorous study of Cherenkov radiation or friction effects at small disorder, which amounts to the metastability of the embedded mass shell of the bare non-relativistic particle when the coupling to the quantized field is turned on. Although this problem is naturally approached by means of Mourre's celebrated commutator method, important regularity issues are known to be inherent to QFT models and restrict the application of the method. In this perspective, we introduce a novel non-standard procedure to construct Mourre conjugate operators, which differs from second quantization and allows to circumvent many regularity issues. To show its versatility, we apply this construction to the Nelson model with massive bosons, to Fröhlich's polaron model, and to a quantum friction model with massless bosons introduced by Bruneau and De Bièvre: for each of those examples, we improve on previous results.

MSC-class: 81T10; 81V73; 47B25; 81Q10; 47A55; 81Q15; 47A40; 81U24.

CONTENTS

1. Introduction and main results	2
1.1. General overview	2
1.2. Translation-invariant massive Nelson model	4
1.3. Quantum friction model	10
1.4. Link to random Schrödinger operators	14
2. Quantum friction model	15
2.1. Construction of conjugate operator	15
2.2. Mourre estimate	21
2.3. Regularity of the interaction	22
2.4. Consequences of Mourre estimate	24
3. Translation-invariant massive Nelson model	25
3.1. Spectrum of uncoupled Hamiltonians	25
3.2. A first construction of conjugate operator	28
3.3. Mourre estimate	30
3.4. Modification procedure and improved regularity	35
3.5. Consequences of Mourre estimate	40
Appendix A. Mourre's commutator method	41
Acknowledgements	46
References	46

1. INTRODUCTION AND MAIN RESULTS

1.1. General overview. This work is devoted to several models in non-relativistic quantum field theory (QFT), describing a non-relativistic quantum particle interacting with a quantized relativistic field of bosons, and we focus on translation-invariant models where the total momentum is conserved. In this context, we aim at the rigorous study of Cherenkov radiation or friction effects: if the initial momentum |P| of the non-relativistic particle exceeds some threshold $|P_{\star}|$ (more precisely, if its initial energy exceeds the minimal energy for one-boson states), then the particle is expected to dissipate energy and to slow down by emitting so-called Cherenkov radiation. In terms of spectral theory, this dissipative phenomenon is translated into the continuity of the energy-momentum spectrum and the absence of embedded mass shell in some region with $|P| > |P_{\star}|$.

We focus on the perturbative regime corresponding to a weak particle-field coupling. In that setting, the starting point is that, for large momentum $|P| > |P_{\star}|$, the mass shell $E = \frac{1}{2}P^2$ of the bare non-relativistic particle is embedded in the continuous spectrum of the uncoupled model. In link with Fermi's golden rule, this mass shell is then indeed expected to be metastable and to disappear as the coupling to the quantized field is turned on. Such spectral results are naturally complemented with scattering resonance descriptions. As interaction Hamiltonians in QFT models are typically not relatively compact, even the perturbative analysis at weak coupling is a nontrivial problem for which still only partial results are available [3, 12, 37, 18, 11]. Note that a different line of research has aimed to rather describe the reduced dynamics of the non-relativistic particle in the kinetic limit: in [49, 19, 13], it is shown to take form of a Boltzmann equation describing the slowdown of the particle. Although supporting the same thesis, such results are limited to diagonal time regimes: they provide a kinetic description only on some suitable timescale, typically for times of order $t = O(g^{-2})$ in terms of the coupling constant $g \ll 1$, and they do not imply any detailed spectral information.

In recent decades, there has been much interest in spectral and scattering theory for QFT models, aiming to adapt the various techniques originally developed for the study of N-particle Schrödinger operators [31]. In particular, Mourre's commutator method [39, 2, 31] has emerged as a fundamental tool to explore the nature of the essential spectrum of such Hamiltonians. This method is more general than related dilation-analyticity techniques and further provides direct insight into time-dependent scattering theory [41, 48, 8]. The starting point is the validity of a so-called Mourre estimate: given a self-adjoint operator H and a spectral interval $J \subset \sigma(H)$, we aim to construct a so-called 'conjugate' operator A such that

$$\mathbb{1}_{J}(H)[H, iA]\mathbb{1}_{J}(H) \ge c_{0}\mathbb{1}_{J}(H) + K,$$

for some constant $c_0 > 0$ and some compact operator K. Conjugate operators can be viewed as quantum analogues of escape functions for Hamiltonian dynamics. Under suitable regularity assumptions (such as H-boundedness of [H, iA]), the Mourre estimate can be used to infer detailed spectral information on H. We also recall the following perturbative version: if a Mourre estimate holds for H and if an H-bounded perturbation V satisfies a suitable regularity condition (such as H-boundedness of [V, iA]), then the perturbed operator H + gV also satisfies a Mourre estimate on any spectral subinterval provided that g is small enough. Perturbative Mourre theory then allows to study the spectrum of perturbed Hamiltonians at weak coupling: it can be used to infer the metastability of embedded bound states under Fermi's golden rule, and time-dependent scattering results are further known to hold under additional regularity assumptions, see [8]. In a nutshell, these techniques allow to reduce the problem to constructing a conjugate operator for the uncoupled Hamiltonian such that the required regularity conditions are satisfied. We refer to Appendix A for a brief overview of Mourre's theory with precise definitions.

Yet, important difficulties arise when applying this method to QFT models: due to the infinite-dimensionality of the underlying Fock space, commutators of the Hamiltonian with natural choices of conjugate operators are typically not relatively bounded. This lack of regularity has led in particular to the development of so-called "singular" Mourre theory in some settings, see [47, 38, 27, 28, 20], but the latter gives no access to time-dependent scattering theory. In the present contribution, we develop a new strategy to construct conjugate operators for QFT models at weak coupling, which solves many regularity issues and thus allows to take advantage of the full power of regular Mourre theory in several new cases. The construction is based on the following two steps:

Step 1: We start with a generic construction for a tentative conjugate operator for the uncoupled model of interest: given a Hamiltonian with a convex symbol h(k), we consider the generator of dilations around the energy minimizer $h(k_0) = \min_k h(k)$, that is, the operator

$$A_{k_0} := \frac{i}{2} \big((k - k_0) \cdot \nabla_k + \nabla_k \cdot (k - k_0) \big).$$
(1.1)

By convexity, this obviously yields a Mourre inequality in form of

$$[h(k), iA_{k_0}] = (k - k_0) \cdot \nabla h(k) \ge 0.$$
(1.2)

In case of QFT models, however, this construction leads to regularity issues: commutators either with the uncoupled Hamiltonian or with the perturbation are generally uncontrolled. A typical issue is that the constructed conjugate (1.1) may depend on the number of bosons via the energy minimizer k_0 and therefore not lead to a second-quantization operator.

Step 2: Although conjugates that are not second-quantization operators lead to regularity issues in general, second-quantization operators are not the only type of admissible operators. As inspired by our previous work [17], we devise a new construction to replace second quantization, which may a priori seem quite iconoclastic: instead of summing operators over different boson variables, we consider some 'signed maximum' of operators, see Section 2.1.2. Using this to modify the tentative construction in Step 1, we manage to solve many regularity issues.

To show the wide applicability of this procedure to construct conjugate operators, we illustrate it on two paradigmatic models.

— Translation-invariant Nelson model with massive bosons: Previous results on this model were restricted to the energy-momentum spectrum below the two-boson threshold and were further limited by the lack of regularity [3, 35, 30, 37, 18]. Our new construction (1.1) amounts to expressing relative boson momenta in the frame that minimizes the kinetic energy, which leads us to new conjugates that allow to study the spectrum for the first time beyond the two-boson threshold. In addition, we manage to avoid regularity issues and to exploit the full power of Mourre's theory: it leads us to a better understanding of Cherenkov radiation, in particular by deriving scattering resonance descriptions. Our construction is quite general and may be of independent interest for other massive QFT models: the same analysis can be repeated, for instance, for Fröhlich's polaron model [36].

— Quantum friction model with massless bosons introduced in [6, 11]: Previous results on this model were limited by the lack of regularity [11]. By our new construction of conjugate operators, we manage to cure all regularity issues and to exploit the full power of Mourre's theory. This considerably improves on previous results on the topic, by establishing in particular the first scattering resonance description.

We expect these results to provide valuable inspiration to further understand Cherenkov radiation in the *massless* Nelson model, for which the only known result at the moment is a weak form of instability for the embedded mass shell [12]. Another perspective concerns the large-coupling regime, for which very little is known even for the massive Nelson model beyond the two-boson threshold, cf. [37, 18], and our new constructions can also be expected to inspire further progress in that case.

In the next two subsections, we introduce in full detail the two models that we focus on in this work, and we formulate our main results, the proofs of which are postponed to Sections 2 and 3.

1.2. Translation-invariant massive Nelson model. The Nelson model was introduced in [40] as a toy model in QFT for a free non-relativistic quantum particle interacting linearly with a quantized radiation field of relativistic scalar bosons; see e.g. [36] and references therein. While very complete results are available in the confined setting [15, 4, 29, 16, 28, both for massive and massless bosons, the understanding remains quite limited in the translation-invariant setting that we consider here. For massive bosons, a detailed description of the bottom of the energy-momentum spectrum was obtained in [50, 35, 36], but the structure of the essential spectrum is only understood below the two-boson threshold, both at weak [33, 3] and large coupling [37, 18]. For massless bosons, the bottom of the spectrum is studied in [24, 25, 42, 26, 43] and the upper spectrum in [9]in the case $|P| < |P_{\star}|$, but no spectral result seems available for $|P| > |P_{\star}|$: the only known result related to Cherenkov radiation is a weak form of instability for the embedded mass shell [12]. In the sequel, we shall focus on the case of massive bosons in the weakcoupling regime and use Mourre's theory to investigate the essential spectrum around the embedded mass shell beyond the two-boson threshold, aiming at a detailed understanding of Cherenkov radiation in that case.

1.2.1. *Description of the model.* The state space for the Nelson model is given by the product Hilbert space

$$\mathcal{H} := \mathcal{H}^{\mathrm{p}} \otimes \mathcal{H}^{\mathrm{f}}, \tag{1.3}$$

where:

- $\mathcal{H}^{p} := L^{2}(\mathbb{R}^{d})$ is the state space for the non-relativistic quantum particle, and we denote respectively by x and $p = \frac{1}{i} \nabla_{x}$ the particle position and momentum coordinates;
- \mathcal{H}^{f} is the state space for the quantized radiation field and takes form of the bosonic Fock space

$$\mathcal{H}^{\mathrm{f}} := \Gamma_s(\mathfrak{h}) := \bigoplus_{n=0}^{\infty} \Gamma_s^{(n)}(\mathfrak{h}),$$

constructed on the single-boson space $\mathfrak{h} := L^2(\mathbb{R}^d)$. In other words, we set $\Gamma_s^{(0)}(\mathfrak{h}) := \mathbb{C}\Omega$ with Ω the vacuum state, and for $n \geq 1$ the *n*-boson state space is the *n*-fold symmetric tensor product

$$\Gamma_s^{(n)}(\mathfrak{h}) := \mathfrak{h}^{\otimes_s n}.$$

We work in the momentum representation, with $k \in \mathbb{R}^d$ standing for the momentum coordinate of the field bosons.

On this bosonic Fock space \mathcal{H}^{f} , we use standard notation for creation and annihilation operators $\{a^*(k)\}_{k\in\mathbb{R}^d}$ and $\{a(k)\}_{k\in\mathbb{R}^d}$, which obey the canonical commutation relations

$$[a^*(k), a^*(k')] = [a(k), a(k')] = 0, \qquad [a(k), a^*(k')] = \delta(k - k'), \qquad a(k)\Omega = 0.$$

We also write $d\Gamma(A)$ for the second quantization of an operator A on \mathfrak{h} , and in particular $N := d\Gamma(1)$ is the number operator on \mathcal{H}^{f} . In this setting, we consider the following translation-invariant Hamiltonian,

$$H_g := H^{\mathbf{p}} \otimes \mathbb{1}_{\mathcal{H}^{\mathbf{f}}} + \mathbb{1}_{\mathcal{H}^{\mathbf{p}}} \otimes H^{\mathbf{f}} + g\Phi(\rho_x) \quad \text{on } \mathcal{H},$$
(1.4)

where:

 the Hamiltonian of the free quantum particle is given by the standard non-relativistic dispersion relation

$$H^{\mathbf{p}} := \frac{1}{2}p^2 \qquad \text{on } \mathcal{H}^{\mathbf{p}};$$

— the free field Hamiltonian is given by second quantization,

$$H^{\mathrm{f}} := \mathrm{d}\Gamma(\omega) = \int_{\mathbb{R}^d} \omega(k) \, a^*(k) a(k) \, \mathrm{d}k \qquad \mathrm{on} \ \mathcal{H}^{\mathrm{f}},$$

where for bosons of mass $m \ge 0$ the single-boson dispersion relation reads

$$\omega(k) := \sqrt{m^2 + |k|^2}; \tag{1.5}$$

— the real number g is the coupling constant for the particle with the bosonic field;

— the interaction Hamiltonian is given by a translation-invariant field operator

$$\Phi(\rho_x) := \int_{\mathbb{R}^d} \rho(k) \left(a^*(k) e^{-ik \cdot x} + a(k) e^{ik \cdot x} \right) \mathrm{d}k, \tag{1.6}$$

for some real-valued interaction kernel $\rho \in L^2(\mathbb{R}^d)$ with $\rho \neq 0$.

Our results on this model will be restricted to the case of massive bosons m > 0 in the weak-coupling regime $|g| \ll 1$. We could also treat the case of a single-boson dispersion relation of the form $\omega(k) = m + |k|$ with m > 0. Regarding the interaction kernel ρ , we shall need to assume strong enough regularity both in infrared and ultraviolet domains, typically requiring ρ to have both some H^s regularity and some polynomial decay.

Before studying this model, we recall its well-posedness properties. For that purpose, we first define the vector subspace

$$\mathcal{C}^{\mathrm{f}} := \mathrm{d}\Gamma_{\mathrm{fin}}(C^{\infty}_{c}(\mathbb{R}^{d})) \subset \mathcal{H}^{\mathrm{f}},$$

where for a vector subspace $\mathfrak{g} \subset \mathfrak{h}$ we denote by $\Gamma_{\text{fin}}(\mathfrak{g})$ the algebraic direct sum of the algebraic tensor products $\mathfrak{g}^{\otimes_s n}$. In these terms, the uncoupled Nelson Hamiltonian

$$H_0 = \mathcal{H}^{\mathrm{p}} \otimes \mathbb{1}_{H^{\mathrm{f}}} + \mathbb{1}_{H^{\mathrm{p}}} \otimes \mathcal{H}^{\mathrm{t}}$$

is clearly essentially self-adjoint on $C_c^{\infty}(\mathbb{R}^d) \otimes \mathcal{C}^{\mathrm{f}}$ (henceforth, tensor products between spaces that are not complete are implicitly understood in the algebraic sense). Besides, as $\rho \in \mathrm{L}^2(\mathbb{R}^d)$, standard estimates ensure that the field operator $\Phi(\rho_x)$ is $(\mathbb{1}_{\mathcal{H}^{\mathrm{p}}} \otimes N^{1/2})$ bounded. In case of massive bosons m > 0, as $\mathrm{d}\Gamma(\omega) \ge mN$, this entails that $\Phi(\rho_x)$ is an infinitesimal perturbation of H_0 . The Kato–Rellich theorem then ensures that for all g the coupled Nelson Hamiltonian H_g is self-adjoint on the same domain $\mathcal{D}(H_0)$ and essentially self-adjoint on the same core $C_c^{\infty}(\mathbb{R}^d) \otimes \mathcal{C}^{\mathrm{f}}$. 1.2.2. Translation invariance. By definition, cf. (1.4), the Hamiltonian H_g is translationinvariant in the sense that it commutes with the total momentum operator

$$P_{ ext{tot}} := p \otimes \mathbb{1}_{\mathcal{H}^{ ext{f}}} + \mathbb{1}_{\mathcal{H}^{ ext{p}}} \otimes \mathrm{d}\Gamma(k) \qquad ext{on } \mathcal{H}.$$

This allows to decompose H_g as a direct integral with respect to the spectrum of the latter. More precisely, in terms of the following unitary transformation, which goes back to Lee, Low, and Pines [32],

$$U: \mathcal{H} \to \int_{\mathbb{R}^d}^{\oplus} \mathcal{H}^{\mathrm{f}} \, \mathrm{d}P, \qquad U := (F \otimes \mathrm{Id}_{\mathcal{H}^{\mathrm{f}}}) \circ \Gamma(e^{ik \cdot x}).$$

where F stands for the Fourier transform on \mathcal{H}^{p} and where Γ is the second quantization functor, we obtain the decomposition

$$UH_g U^* = \int_{\mathbb{R}^d}^{\oplus} H_g(P) \, \mathrm{d}P \qquad \text{on } \int_{\mathbb{R}^d}^{\oplus} \mathcal{H}^{\mathrm{f}} \, \mathrm{d}P, \tag{1.7}$$

where for all $P \in \mathbb{R}^d$ the fiber Hamiltonian $H_q(P)$ takes the form

$$H_g(P) := \frac{1}{2}(P - \mathrm{d}\Gamma(k))^2 + H^{\mathrm{f}} + g\Phi(\rho) \qquad \text{on } \mathcal{H}^{\mathrm{f}}, \tag{1.8}$$

in terms of the fiber interaction Hamiltonian

$$\Phi(\rho) := \int_{\mathbb{R}^d} \rho(k) \left(a^*(k) + a(k) \right) \, \mathrm{d}k.$$
(1.9)

We recall well-posedness properties of these fiber Hamiltonians. First, for P = 0, the uncoupled fiber Hamiltonian $H_0(0) = \frac{1}{2} d\Gamma(k)^2 + H^f$ is essentially self-adjoint on C^f . Next, for any $P \in \mathbb{R}^d$, noting that $\frac{1}{2}(P - d\Gamma(k))^2 - \frac{1}{2}d\Gamma(k)^2 = \frac{1}{2}|P|^2 - P \cdot d\Gamma(k)$ is an infinitesimal perturbation of $H_0(0)$, the Kato–Rellich theorem ensures that the uncoupled fiber Hamiltonian

$$H_0(P) = \frac{1}{2}(P - d\Gamma(k))^2 + H^{f}$$

is also essentially self-adjoint on \mathcal{C}^{f} and that its domain is independent of P,

$$\mathcal{D} := \mathcal{D}(H_0(P)) = \mathcal{D}(H_0(0)) = \mathcal{D}(\mathrm{d}\Gamma(k)^2) \cap \mathcal{D}(\mathrm{d}\Gamma(\omega)).$$
(1.10)

Besides, as $\rho \in L^2(\mathbb{R}^d)$, standard estimates ensure that the field operator $\Phi(\rho)$ is $N^{1/2}$ bounded. In case of massive bosons m > 0, as $d\Gamma(\omega) \ge mN$, this entails that $\Phi(\rho)$ is an infinitesimal perturbation of $H_0(P)$. The Kato–Rellich theorem then ensures that for all gthe coupled fiber Hamiltonian $H_g(P)$ is self-adjoint on the same domain \mathcal{D} and essentially self-adjoint on the same core \mathcal{C}^{f} .

1.2.3. Energy-momentum spectrum. In this translation-invariant setting, the natural object of study is the energy-momentum spectrum $\{(P, E) : E \in \sigma(H_g(P))\}$, where $\sigma(H_g(P))$ is the spectrum of the fiber Hamiltonian $H_g(P)$ at fixed total momentum P. We start by recalling the explicit structure of this spectrum for the uncoupled Hamiltonian.

Lemma 1.1 (Spectrum of uncoupled Nelson model). Consider the translation-invariant Nelson model with massive bosons m > 0, cf. (1.3)–(1.9). Given a total momentum $P \in \mathbb{R}^d$, the spectrum of the uncoupled fiber Hamiltonian $H_0(P)$ is given by

$$\sigma_{\rm pp}(H_0(P)) = \{\frac{1}{2}P^2\}, \quad \sigma_{\rm ac}(H_0(P)) = [E_0(P), \infty), \quad \sigma_{\rm sc}(H_0(P)) = \emptyset, \tag{1.11}$$

where the eigenvalue $\frac{1}{2}P^2$ is simple and is associated with the vacuum state Ω , and where the bottom of the absolutely continuous spectrum is given by

$$E_0(P) := \frac{1}{2}c(P)^2 + \sqrt{m^2 + (|P| - c(P))^2}, \qquad (1.12)$$

in terms of the unique solution $c(P) \in [0,1)$ of the implicit equation

$$c(P) = \frac{|P| - c(P)}{\sqrt{m^2 + (|P| - c(P))^2}}$$

Moreover, there is a unique critical value $|P_{\star}| > 1$ such that

$$E_0(P_\star) = \frac{1}{2} P_\star^2, \tag{1.13}$$

and the following alternative then holds:

- for $|P| < |P_{\star}|$, the fiber Hamiltonian $H_0(P)$ has an isolated ground state at $\frac{1}{2}P^2$; - for $|P| > |P_{\star}|$, the fiber Hamiltonian $H_0(P)$ has no ground state and its eigenvalue is
- for $|P| > |P_{\star}|$, the fiber Hamiltonian $H_0(P)$ has no ground state and its eigenvalue is embedded in the absolutely continuous spectrum. \diamond

Before turning to our main results on coupled Hamiltonians, we further elaborate on this statement and emphasize the layered structure of the spectrum. By definition (1.8), the uncoupled fiber Hamiltonian commutes with the number operator N and thus splits as a direct sum on many-boson state spaces,

$$H_0(P) = \bigoplus_{n=0}^{\infty} H_0^{(n)}(P) \quad \text{on} \quad \mathcal{H}^{\mathrm{f}} = \bigoplus_{n=0}^{\infty} \Gamma_s^{(n)}(\mathfrak{h}), \quad (1.14)$$

in terms of the restrictions

$$H_0^{(n)}(P) := H_0(P)|_{\Gamma_s^{(n)}(\mathfrak{h})}$$

While the eigenvalue $\frac{1}{2}P^2$ is associated with the vacuum state Ω and corresponds to a free non-relativistic particle with momentum P, the absolutely continuous spectrum corresponds to states supporting at least one boson,

$$\sigma_{\rm ac}(H_0(P)) = \operatorname{adh} \bigcup_{n=1}^{\infty} \sigma_{\rm ac}(H_0^{(n)}(P)), \qquad (1.15)$$

where adh stands for the closure. For all $n \ge 1$, the restriction $H_0^{(n)}(P)$ is a multiplication operator in momentum coordinates, with symbol

$$H_0^{(n)}(P;k_1,\ldots,k_n) := \frac{1}{2} \left(P - \sum_{j=1}^n k_j \right)^2 + \sum_{j=1}^n \omega(k_j).$$
(1.16)

Its spectrum is absolutely continuous and coincides with the essential image of the symbol,

$$\sigma_{\rm ac}\big(H_0^{(n)}(P)\big) = \big[E_0^{(n)}(P),\infty\big), \qquad \sigma_{\rm pp}\big(H_0^{(n)}(P)\big) = \sigma_{\rm sc}\big(H_0^{(n)}(P)\big) = \varnothing,$$

in terms of the so-called n-boson energy threshold

$$E_0^{(n)}(P) := \min_{k_1,\dots,k_n \in \mathbb{R}^d} H_0^{(n)}(P;k_1,\dots,k_n).$$
(1.17)

In the case of massive bosons m > 0, it is easily checked that

$$E_0^{(n)}(P) < E_0^{(n+1)}(P)$$
 for all n , and $E_0^{(n)}(P) \uparrow \infty$ as $n \uparrow \infty$,

cf. Lemma 3.2 below. In view of (1.15), this ensures in particular that the bottom of the absolutely continuous spectrum is

$$E_0(P) := E_0^{(1)}(P) = \min_{n \ge 1} E_0^{(n)}(P),$$

and we then recover the expression (1.12) by computing the minimum of (1.16) for n = 1. Due to the layered structure of the spectrum, cf. (1.15), our results in the sequel are naturally restricted away from energy thresholds.

1.2.4. Main results. We may now formulate our main results on the massive Nelson model. Our starting point is the following perturbative Mourre commutator result: for $|P| > |P_{\star}|$, as the eigenvalue $\frac{1}{2}P^2$ is embedded in the essential spectrum, a Mourre estimate is proven around and above the eigenvalue away from energy thresholds, as well as below the twoboson threshold, and this is complemented with a regularity statement for the interaction Hamiltonian. We refer to Appendix A for standard definitions and notation related to Mourre's theory.

Theorem 1.2 (Mourre estimate for Nelson model). Consider the translation-invariant Nelson model with massive bosons m > 0, cf. (1.3)–(1.9). Given a total momentum $|P| > |P_{\star}|$, define $n_P \geq 1$ such that

$$\frac{1}{2}|P|^2 \in \left[E_0^{(n_P)}(P), E_0^{(n_P+1)}(P)\right),\tag{1.18}$$

where we recall definitions (1.13) and (1.17). Then, for n = 1 as well as for any $n \ge n_P$, we can construct an operator $A_{P,n}$ on \mathcal{H}^{f} , essentially self-adjoint on \mathcal{C}^{f} , with the following properties.

- (i) The uncoupled fiber Hamiltonian $H_0(P)$ is of class $C^{\infty}(A_{P,n})$. Moreover, the unitary group generated by $A_{P,n}$ leaves the domain of $H_0(P)$ invariant, and the iterated commutators $\operatorname{ad}_{iA_{P,n}}^s(H_0(P))$ extend as $H_0(P)$ -bounded operators for all $s \geq 0$.
- (ii) For all $\varepsilon > 0$ and all energy intervals $I \subset \left[E_0^{(n)}(P) + \varepsilon, E_0^{(n+1)}(P)\right)$, the following Mourre estimate holds with respect to $A_{P,n}$ on I,

$$\mathbb{1}_{I}(H_{0}(P)) \left[H_{0}(P), iA_{P,n} \right] \mathbb{1}_{I}(H_{0}(P)) \geq \varepsilon \bar{\Pi}_{\Omega} \, \mathbb{1}_{I}(H_{0}(P)) \, \bar{\Pi}_{\Omega},$$

where $\overline{\Pi}_{\Omega}$ is the orthogonal projection on $\mathbb{C}\Omega^{\perp}$. In particular, the Mourre estimate is strict if the interval I does not contain the eigenvalue $\frac{1}{2}P^2$.

(iii) The fiber interaction Hamiltonian $\Phi(\rho)$ satisfies the following regularity condition: if the interaction kernel ρ belongs to $H^{\nu}(\mathbb{R}^d)$ with $\langle k \rangle^{\nu} \nabla^{\nu} \rho \in L^2(\mathbb{R}^d)$ for some $\nu \geq 1$, then the iterated commutators $\operatorname{ad}_{iA_{P,n}}^s(\Phi(\rho))$ extend as $H_0(P)^{1/2}$ -bounded operators for all $1 \leq s \leq \nu$.

Compared to previous work on the topic [3, 37, 18], this provides the first Mourre estimate above the two-boson threshold. In addition, the C^{∞} -regularity stated in item (i) allows us to exploit for the first time the full power of Mourre's theory, cf. Appendix A, while previous constructions were restricted to C^2 -regularity. As a corollary, we deduce the following description of the essential spectrum of fiber Hamiltonians at weak coupling, which proves in particular the instability of the mass shell $E = \frac{1}{2}P^2$ of the free non-relativistic particle when coupled to the bosonic field. This is further complemented with a dynamical resonance description, which exploits the C^{∞} -regularity and is thus new even below the two-boson threshold. It constitutes a precise formulation of Cherenkov radiation for the massive Nelson model.

Corollary 1.3 (Cherenkov radiation for Nelson model). Consider the translation-invariant Nelson model with massive bosons m > 0, cf. (1.3)–(1.9), and assume that the interaction kernel ρ belongs to $H^2(\mathbb{R}^d)$ with $\langle k \rangle^2 \nabla^2 \rho \in L^2(\mathbb{R}^d)$. Given a total momentum $|P| > |P_{\star}|$, define $n_P \geq 1$ as in (1.18), and assume that Fermi's condition holds,

$$\gamma_P := \frac{1}{2} (2\pi)^{1-d} \int_{\{k: \frac{1}{2}(P-k)^2 + \omega(k) = \frac{1}{2}P^2\}} \frac{|\rho(k)|^2}{|k-P+\nabla\omega(k)|} \, \mathrm{d}\mathcal{H}_{d-1}(k) > 0, \qquad (1.19)$$

where \mathcal{H}_{d-1} stands for the (d-1)th-dimensional Hausdorff measure. (This condition holds in particular if ρ does not vanish.) Then, the following properties hold.

(i) Absence of embedded mass shell:

For all $g \neq 0$, the essential spectrum of the fiber Hamiltonian $H_g(P)$ is purely absolutely continuous below $E_0^{(2)}(P)$ and above $E_0^{(n_P)}(P)$ away from thresholds: more precisely, there is a sequence $(C_{P,n})_n$ such that $H_g(P)$ has purely absolutely continuous spectrum in

$$I_{g}(P) := \left(E_{0}^{(1)}(P) + \sqrt{g}C_{P,1}, E_{0}^{(2)}(P) - gC_{P,1}\right)$$
$$\bigcup_{n \ge n_{P}} \left(E_{0}^{(n)}(P) + \sqrt{g}C_{P,n}, E_{0}^{(n+1)}(P) - gC_{P,n}\right).$$

(*ii*) Quasi-exponential decay law:

Further assume that for some $\nu \geq 0$ the interaction kernel ρ belongs to $H^{5+\nu}(\mathbb{R}^d)$ with $\langle k \rangle^{5+\nu} \nabla^{5+\nu} \rho \in L^2(\mathbb{R}^d)$. Then, there is $g_0 > 0$ such that, for all smooth cut-off functions h supported in $I_{g_0}(P)$ and equal to 1 in a neighborhood of the uncoupled eigenvalue $\frac{1}{2}P^2$, there holds for all $t \geq 0$ and $|g| \leq g_0$,

$$\left| \left\langle \Omega, e^{-itH_g(P)} h(H_g(P)) \Omega \right\rangle - e^{-itz_g(P)} \right| \lesssim_{g_0,\rho,h} \begin{cases} g^2 |\log g| \langle t \rangle^{-\nu}, & \text{if } \nu \ge 0; \\ g^2 \langle t \rangle^{-(\nu-1)}, & \text{if } \nu \ge 1; \end{cases}$$

where the dynamical resonance $z_g(P)$ is given by Fermi's golden rule,

$$z_g(P) = \frac{1}{2}P^2 - g^2(\theta_P + i\gamma_P),$$

where $\gamma_P > 0$ is defined in (1.19) and where the real part $\theta_P \in \mathbb{R}$ takes the form

$$\theta_P := (2\pi)^{-d} \text{ p.v.} \int_{\mathbb{R}} (t - \frac{1}{2}P^2)^{-1} \left(\int_{\{k : \frac{1}{2}(P-k)^2 + \omega(k) = t\}} \frac{|\rho(k)|^2}{|k - P + \nabla \omega(k)|} \, \mathrm{d}\mathcal{H}_{d-1}(k) \right) \, \mathrm{d}t.$$

(Henceforth, we use the notation $\leq_{g_0,\rho,h}$ for $\leq C \times$ up to some constant C > 0 that depends on g_0 and on controlled norms of ρ, h .)

In particular, for all $u_{\circ} \in L^{2}(\mathbb{R}^{d})$ with Fourier transform compactly supported in the set $\{P : |P| > |P_{\star}|\}$, provided that ρ does not vanish and satisfies the requirements of (ii) for some $\nu \geq 0$, there holds uniformly for all $t \geq 0$,

$$\langle (\delta_x \otimes \Omega), e^{-itH_g}(u_\circ \otimes \Omega) \rangle = \int_{\mathbb{R}^d} \hat{u}_\circ(P) e^{ix \cdot P - itz_g(P)} \, \mathrm{d}P + o_g(1),$$

where $o_g(1)$ tends to 0 in $L^{\infty}_x(\mathbb{R}^d)$ as $g \downarrow 0$ (depending on ρ, u_{\circ}).

We briefly comment on possible extensions and open problems. First note that the above result gives curiously no access to the spectrum in the energy interval $(E_0^{(2)}(P), E_0^{(n_P)}(P))$ between the two-boson threshold and the last threshold below the embedded eigenvalue:

 \Diamond

although we also expect the same behavior in this interval (away from thresholds), a different type of construction seems to be needed for conjugate operators. Another question concerns the use of the above Mourre estimate to further investigate asymptotic completeness: the extension of [30, 18] beyond the two-boson threshold in the weak-coupling regime is postponed to a future work. Finally, a last question concerns the validity of the Mourre estimate at large coupling: this was solved in [37] below the two-boson threshold and we may expect our present contribution to give valuable inspiration to get beyond that.

The Nelson model belongs to the class of so-called translation-invariant Pauli–Fierz models. These also include Fröhlich's polaron model in solid-state physics [23, 21], as well as non-relativistic QFT models with vector bosons. Our findings are easily extended to those settings:

— The polaron model introduced by Fröhlich [23, 21] describes an electron interacting with lattice vibrations of a polar crystal. These are naturally represented in terms of a Bose field over a crystalline lattice and we consider the continuum limit of the latter, thus treating the crystal as a polarizable continuum. The model then takes the same form as the Nelson model (1.3)–(1.4), where the single-boson dispersion relation is now taken to be constant, $\omega(k) = 1$, so that the free field Hamiltonian is the number operator $H^{f} = N$. We refer to [36] and references therein for a discussion of this model. Our analysis of the massive Nelson model can be repeated mutatis mutandis in this setting and we note that several calculations actually reduce dramatically: in particular, the critical value of the total momentum and the energy thresholds are simply

$$|P_{\star}| = \sqrt{2}$$
 and $E_0^{(n)}(P) = n$.

Corollary 1.3 yields the first rigorous justification of Cherenkov radiation for the polaron model (see formal discussion in [22, p.227–230]).

- Consider the translation-invariant non-relativistic QFT model for a non-relativistic quantum particle minimally coupled to a quantized radiation field of relativistic vector bosons [51, 34]. For d = 3, the single-boson space is then $\mathfrak{h} := L^2(\mathbb{R}^3 \times \{+, -\})$, where +/- stands for boson polarization, and we consider the Hamiltonian

$$H_{\alpha} := \frac{1}{2} \left(p - \alpha^{\frac{1}{2}} A_x \right)^2 + \mathbb{1} \otimes H^{\mathrm{f}},$$

where $\alpha \geq 0$ is the coupling constant, where the vector potential A_x is linear in creation and annihilation operators, and where the free field Hamiltonian is as before $H^{\rm f} = d\Gamma(\omega)$ with single-boson dispersion relation $\omega(k) = \sqrt{m^2 + k^2}$. In case of massive vector bosons m > 0, our analysis of the massive Nelson model is easily adapted to this other model under suitable regularity assumptions on the interaction kernel defining the vector potential; we skip the detail.

While we focus here on massive bosons, the case of massless bosons is different and will be commented at the end of the next section.

1.3. Quantum friction model. We turn to the quantum version [6, 11] of a translationinvariant Hamiltonian model for friction introduced by Bruneau and De Bièvre [7]. It describes a non-relativistic quantum particle moving through a translation-invariant medium consisting of uncoupled quantized vibration fields at each point in space. This model happens to be substantially simpler to study than the Nelson model, precisely due to the fact that vibration fields are uncoupled in space, and we shall thus be able to further treat the case of massless bosons for this model. In the sequel, we aim at a detailed understanding of friction effects in the weak coupling regime, improving on previous results in [11].

1.3.1. *Description of the model.* The state space for the model is given by the product Hilbert space

$$\mathcal{H} := \mathcal{H}^{\mathrm{p}} \otimes \mathcal{H}^{\mathrm{f}}, \tag{1.20}$$

where:

- $\mathcal{H}^{p} := L^{2}(\mathbb{R}^{d})$ is the state space for the non-relativistic quantum particle, and we denote respectively by x and $p = \frac{1}{i} \nabla_{x}$ the particle position and momentum coordinates;
- \mathcal{H}^{f} is the state space for the quantized vibration fields and takes form of the bosonic Fock space $\mathcal{H}^{\mathrm{f}} := \Gamma_s(\mathfrak{h})$ constructed on the single-boson space $\mathfrak{h} := \mathrm{L}^2(\mathbb{R}^q \times \mathbb{R}^d)$. We work in momentum representation, with $k \in \mathbb{R}^q$ standing for the momentum coordinate for vibrational degrees of freedom, and with $\xi \in \mathbb{R}^d$ standing for the momentum coordinate dual to the particle position x.

On this bosonic Fock space \mathcal{H}^{f} , we use standard notation for creation and annihilation operators $\{a^*(k,\xi)\}_{(k,\xi)\in\mathbb{R}^q\times\mathbb{R}^d}$ and $\{a(k,\xi)\}_{(k,\xi)\in\mathbb{R}^q\times\mathbb{R}^d}$, we use the notation $\mathrm{d}\Gamma(A)$ for the second quantization of an operator A on \mathfrak{h} , and in particular $N := \mathrm{d}\Gamma(1)$ is the number operator. In this setting, we consider the following translation-invariant Hamiltonian,

$$H_g := H^{\mathbf{p}} \otimes \mathbb{1}_{\mathcal{H}^{\mathbf{f}}} + \mathbb{1}_{\mathcal{H}^{\mathbf{p}}} \otimes H^{\mathbf{f}} + g\Phi(\rho_x) \quad \text{on } \mathcal{H},$$
(1.21)

where:

 — the Hamiltonian of the free quantum particle is given by the standard non-relativistic dispersion relation

$$H^{\mathbf{p}} := \frac{1}{2}p^2 \qquad \text{on } \mathcal{H}^{\mathbf{p}}$$

— the free field Hamiltonian is given by second quantization

$$H^{\mathrm{f}} := \mathrm{d}\Gamma(\omega) = \iint_{\mathbb{R}^{q} \times \mathbb{R}^{d}} \omega(k,\xi) a^{*}(k,\xi) a(k,\xi) \,\mathrm{d}k \,\mathrm{d}\xi \qquad \text{on } \mathcal{H}^{\mathrm{f}}$$

where the single-boson dispersion relation reads

$$\omega(k,\xi) := |k|, \tag{1.22}$$

which corresponds to massless bosons and is naturally independent of ξ as vibration fields at different values of x are not coupled;

- the real number g is the coupling constant for the particle with the vibration fields;
- the interaction Hamiltonian is given by a translation-invariant field operator

$$\Phi(\rho_x) := \iint_{\mathbb{R}^q \times \mathbb{R}^d} \rho(k,\xi) \Big(a^*(k,\xi) e^{-i\xi \cdot x} + a(k,\xi) e^{i\xi \cdot x} \Big) \, \mathrm{d}k \, \mathrm{d}\xi, \tag{1.23}$$

for some real-valued interaction kernel $\rho \in L^2(\mathbb{R}^q \times \mathbb{R}^d)$ with $\rho \neq 0$.

Our results on this model are restricted to the weak-coupling regime $|g| \ll 1$. Regarding the interaction kernel ρ in (1.23), we shall need strong ultraviolet regularity, but a quite general infrared behavior is allowed. More precisely, we consider the following assumption, for some $\nu \geq 0$,

(Reg_{ν}) There holds $(1 + |k|^{-\frac{1}{2}})(k \cdot \nabla_k)^{\alpha}(\xi \cdot \nabla_{\xi})^{\beta} \nabla_{\xi}^{\gamma} \rho \in L^2(\mathbb{R}^q \times \mathbb{R}^d)$ for all $\alpha, \beta, \gamma \ge 0$ with $\alpha + \beta + \gamma \le \nu$.

 \Diamond

Note that this holds for all ν for instance if ρ takes the particular form $\rho(k,\xi) = |k|^{\mu}\sigma(k,\xi)$ for some $\sigma \in \mathcal{S}(\mathbb{R}^q \times \mathbb{R}^d)$ and $\mu > -\frac{1}{2}(q-1)$. This restriction on the infrared parameter μ is essentially optimal in the sense that it precisely ensures that the interaction Hamiltonian $\Phi(\rho_x)$ is relatively bounded with respect to H_0 .

1.3.2. Translation invariance. As in (1.7) (see also [11, Section 3.2]), we can decompose the above Hamiltonian H_g as a direct integral

$$H_g \cong \int_{\mathbb{R}^d}^{\oplus} H_g(P) \, \mathrm{d}P \qquad \text{on } \int_{\mathbb{R}^d}^{\oplus} \mathcal{H}^{\mathrm{f}} \, \mathrm{d}P,$$

where for all $P \in \mathbb{R}^d$ the fiber Hamiltonian $H_q(P)$ takes the form

$$H_g(P) := \frac{1}{2} (P - d\Gamma(\xi))^2 + H^{\rm f} + g \Phi(\rho).$$
(1.24)

We recall well-posedness properties of these fiber Hamiltonians. First note that the uncoupled fiber Hamiltonian $H_0(P)$ is essentially self-adjoint on

$$\mathcal{C}^{\mathrm{f}} := \mathrm{d}\Gamma_{\mathrm{fin}}(C^{\infty}_{c}(\mathbb{R}^{q} \times \mathbb{R}^{d})),$$

and that its domain is independent of P,

$$\mathcal{D} := \mathcal{D}(H_0(P)) = \mathcal{D}(H_0(0)) = \mathcal{D}(\mathrm{d}\Gamma(\xi)^2) \cap \mathcal{D}(\mathrm{d}\Gamma(|k|)).$$
(1.25)

Besides, standard estimates ensure that $\Phi(\rho)$ is $(H^{f})^{1/2}$ -bounded provided that the interaction kernel satisfies Assumption (Reg₀). The Kato–Rellich theorem then ensures that for all g the coupled fiber Hamiltonian $H_g(P)$ is self-adjoint on the same domain \mathcal{D} and essentially self-adjoint on the same core \mathcal{C}^{f} .

1.3.3. *Energy-momentum spectrum.* We start by recalling the structure of the energymomentum spectrum of the uncoupled Hamiltonian.

Lemma 1.4 (Spectrum of uncoupled quantum friction model). Consider the quantum friction model (1.20)–(1.24). For any total momentum $P \in \mathbb{R}^d$, the spectrum of the uncoupled fiber Hamiltonian $H_0(P)$ is given by

$$\sigma_{\rm pp}(H_0(P)) = \{\frac{1}{2}P^2\}, \quad \sigma_{\rm ac}(H_0(P)) = [0,\infty), \quad \sigma_{\rm sc}(H_0(P)) = \emptyset,$$
(1.26)

where the eigenvalue $\frac{1}{2}P^2$ is simple and is associated with the vacuum state Ω .

We emphasize the key difference with the Nelson model: as vibration fields at different positions in space are not coupled, the propagation speed of bosons in space vanishes and the spectrum of the uncoupled fiber Hamiltonian is therefore $[0, \infty)$ for any total momentum P. In particular, the eigenvalue $\frac{1}{2}P^2$ is strictly embedded in absolutely continuous spectrum whenever $P \neq 0$. Hence, Cherenkov radiation is expected to occur whenever the non-relativistic particle has a non-vanishing momentum, which then results in a friction effect that tends to stop the particle.

1.3.4. Main results. We may now formulate our main results on the quantum friction model. Our starting point is the following perturbative Mourre commutator result. Note that our construction only yields a Mourre estimate above $\frac{1}{18}P^2$, but this is enough for our purposes as it covers a neighborhood of the embedded eigenvalue $\frac{1}{2}P^2$.

Theorem 1.5 (Mourre estimate for quantum friction model). Consider the quantum friction model (1.20)–(1.24). Given a total momentum $P \neq 0$ and given $\delta > 0$, we can construct a self-adjoint operator $A_{P,\delta}$ in \mathcal{H}^{f} , essentially self-adjoint on \mathcal{C}^{f} , with the following properties.

- (i) The uncoupled fiber Hamiltonian $H_0(P)$ is of class $C^{\infty}(A_{P;\delta})$. Moreover, the unitary group generated by $A_{P;\delta}$ leaves the domain of $H_0(P)$ invariant, and the iterated commutators $\operatorname{ad}_{iA_{P;\delta}}^s(H_0(P))$ extend as $H_0(P)$ -bounded operators for all $s \geq 0$.
- (ii) For all $\varepsilon > 0$ and all energy intervals $I \subset \left[\frac{1}{18}P^2 + \delta + \varepsilon, \infty\right)$, the following Mourre estimate holds with respect to $A_{P;\delta}$ on I,

$$\mathbb{1}_{I}(H_{0}(P))[H_{0}(P), iA_{P;\delta}] \mathbb{1}_{I}(H_{0}(P)) \geq \varepsilon \mathbb{1}_{I}(H_{0}(P)) - P^{2}\Pi_{\Omega},$$

where Π_{Ω} is the orthogonal projection on $\mathbb{C}\Omega$. In particular, the Mourre estimate is strict if the interval I does not contain the eigenvalue $\frac{1}{2}P^2$.

(iii) The fiber interaction Hamiltonian $\Phi(\rho)$ satisfies the following regularity condition: if the interaction kernel ρ satisfies (Reg_{ν}) for some $\nu \geq 1$, then the iterated commutators $\operatorname{ad}_{iA_{P,\delta}}^s(\Phi(\rho))$ extend as $H_0(P)^{1/2}$ -bounded operators for all $1 \leq s \leq \nu$.

This provides the first Mourre estimate with C^{∞} -regularity for this model. In contrast, in [11], the authors used the generator of radial translations $R := \frac{i}{2} d\Gamma(\frac{k}{|k|} \cdot \nabla_k + \nabla_k \cdot \frac{k}{|k|})$ as a natural conjugate and were confronted both with the singularity of this operator at small k and with the dramatic lack of associated regularity: the commutator with $H_0(P)$ is formally

$$[H_0(P), iR] = N,$$

which is positive on $\mathbb{C}\Omega^{\perp}$ but is not controlled by $H_0(P)$ in the case of massless bosons. To accommodate this difficulty, the authors of [11] had to appeal to the "singular" Mourre theory developed in [47, 38, 27, 28, 20], which is precisely meant for this situation but allows for weaker consequences than the regular theory. In particular, they deduced in [11] the instability of the embedded mass shell $E = \frac{1}{2}P^2$ at weak coupling, but no scattering resonance description was obtained. In addition, their restriction on the infrared behavior of the interaction kernel ρ was much stronger than what we impose here in Assumption (Reg_{ν}). Rather combining Theorem 1.5 above with the full power of regular Mourre theory, we are led to the following improved description.

Corollary 1.6 (Quantum friction). Consider the quantum friction model (1.20)–(1.24), and assume that the interaction kernel ρ satisfies Assumption (Reg₂). Given a total momentum $P \neq 0$, assume that Fermi's condition holds,

$$\gamma_P := \frac{1}{2} (2\pi)^{1-d} \int_{|k| \le \frac{1}{2} P^2} \int_{\{\xi : \frac{1}{2} (P-\xi)^2 = \frac{1}{2} P^2 - |k|\}} \frac{|\rho(k,\xi)|^2}{\sqrt{(P-\xi)^2 + 1}} d\mathcal{H}_{d-1}(\xi) dk > 0, \quad (1.27)$$

where \mathcal{H}_{d-1} stands for the (d-1)th-dimensional Hausdorff measure. (This condition holds in particular if ρ does not vanish.) Then, the following properties hold.

(i) Absence of embedded mass shell:

For all $g \neq 0$, the coupled fiber Hamiltonian $H_g(P)$ has purely absolutely continuous spectrum in

$$I_g(P) := \left(\frac{1}{18}P^2 + \sqrt{g}C_P, \infty\right).$$

 \Diamond

(*ii*) Quasi-exponential decay law:

Further assume that for some $\nu \geq 0$ the interaction kernel ρ satisfies Assumption (Reg_{5+ ν}). Then, there is $g_0 > 0$ such that, for all smooth cut-off functions h supported in $I_{g_0}(P)$ and equal to 1 in a neighborhood of the uncoupled eigenvalue $\frac{1}{2}P^2$, there holds for all $t \geq 0$ and $|g| \leq g_0$,

$$\left| \left\langle \Omega, e^{-itH_g(P)} h(H_g(P)) \Omega \right\rangle - e^{-itz_g(P)} \right| \lesssim_{g_0,\rho,h} \begin{cases} g^2 |\log g| \langle t \rangle^{-\nu}, & \text{if } \nu \ge 0; \\ g^2 \langle t \rangle^{-(\nu-1)}, & \text{if } \nu \ge 1; \end{cases}$$

where the dynamical resonance $z_g(P)$ is given by Fermi's golden rule,

$$z_g(P) = \frac{1}{2}P^2 - g^2(\theta_P + i\gamma_P),$$

where $\gamma_P > 0$ is defined in (1.27) and where the real part $\theta_P \in \mathbb{R}$ takes the form

$$\theta_P := (2\pi)^{-d} \text{ p. v.} \int_{\mathbb{R}} (t - \frac{1}{2}P^2)^{-1} \\ \times \left(\int_{|k| \le t} \int_{\{\xi : \frac{1}{2}(P-\xi)^2 = t - |k|} \frac{|\rho(k,\xi)|^2}{\sqrt{(P-\xi)^2 + 1}} \mathrm{d}\mathcal{H}_{d-1}(\xi) \mathrm{d}k \right) \mathrm{d}t.$$

In particular, for all $u_{\circ} \in L^{2}(\mathbb{R}^{d})$ with compactly supported Fourier transform, provided that ρ does not vanish and satisfies the requirements of (ii) for some $\nu \geq 0$, there holds uniformly for all $t \geq 0$,

$$\langle (\delta_x \otimes \Omega), e^{-itH_g}(u_\circ \otimes \Omega) \rangle = \int_{\mathbb{R}^d} \hat{u}_\circ(P) e^{ix \cdot P - itz_g(P)} \, \mathrm{d}P + o_g(1),$$

where $o_g(1)$ tends to 0 in $L^{\infty}_x(\mathbb{R}^d)$ as $g \downarrow 0$ (depending on ρ, u_{\circ}).

As this result concerns massless bosons, we may wonder whether our constructions could be adapted to some extent to the massless Nelson model. At the moment, however, we leave this as an open question. Compared to the quantum friction model, the difficulty is that for the Nelson model (1.3)–(1.4) the momentum coordinates k and ξ coincide: in view of our construction (2.15) of the conjugate operator below, we are then essentially reduced to controlling the commutator $\left[\sum_{j=1}^{n} |\nabla_{z_j}|, \max_j z_j\right]$ uniformly with respect to n, which we do not know how to do. Controlling this commutator may actually require to further adapt our construction of the conjugate operator. For massive bosons, the problem simplifies drastically as a bound O(n) on this commutator is sufficient, cf. Lemma 3.8.

1.4. Link to random Schrödinger operators. As is well known, e.g. [13, Section 1.3], [5], or [17, Lemma 5.6], random Schrödinger operators can be viewed as particular instances of Pauli–Fierz models. This comparison was actually our original motivation for the present contribution, in link with our previous work [17]. More precisely, given a stochastically translation-invariant Gaussian field V on \mathbb{R}^d , constructed on a probability space Ω , a Gaussian chaos decomposition of $L^2(\Omega)$ ensures that the random Schrödinger operator $-\frac{1}{2}\Delta + gV$ in $L^2(\mathbb{R}^d \times \Omega)$ is unitarily equivalent to the translation-invariant Hamiltonian

$$H_g := \frac{1}{2}p^2 \otimes \mathbb{1}_{\mathcal{H}^{\mathrm{f}}} + g\Phi(\rho_x) \quad \text{on } \mathcal{H} = \mathcal{H}^{\mathrm{p}} \otimes \mathcal{H}^{\mathrm{f}},$$

where $\mathcal{H}^{p} := L^{2}(\mathbb{R}^{d})$ and $\mathcal{H}^{f} := \Gamma_{s}(\mathfrak{h})$ with $\mathfrak{h} := L^{2}(\mathbb{R}^{d})$, and where the interaction kernel ρ is such that $\rho * \check{\rho}$ is the covariance function of V. In other words, the action of the Gaussian field is viewed as the interaction with a bosonic field. Via this isomorphism, the

fiber decomposition (1.7) for H_g is precisely equivalent to the Floquet–Bloch type fibration that we introduced in [17] (see also [5]),

$$H_g \cong \int_{\mathbb{R}^d}^{\oplus} H_g(P) \, \mathrm{d}P, \quad \text{where} \quad H_g(P) := \frac{1}{2}(P-k)^2 + g\Phi(\rho) \quad \text{on } \mathcal{H}^{\mathrm{f}}.$$

Compared to QFT models studied in the present work, the key difficulty is that degrees of freedom associated with the random potential V do not evolve in time, hence the dispersion relation for the corresponding bosons is trivial, $\omega = 0$, and the free field Hamiltonian vanishes. This makes the study of random Schrödinger operators particularly intricate from this perspecive: the perturbation $\Phi(\rho)$ is not relatively bounded with respect to $H_0(P)$, and its commutators with any Mourre conjugate operator for $H_0(P)$ would not be relatively bounded either. In other words, $\Phi(\rho)$ is not a regular perturbation in the sense of Mourre's theory. In [17], we proceeded by truncating $\Phi(\rho)$ to state spaces with a bounded number of bosons, depending on the size of the coupling constant g, and this allowed us to deduce a scattering resonance description at least up to the kinetic timescale $t \leq g^{-2}$. As explained in [17], if one had ideal estimates on the number operator along the dynamics, this could be extended to $t \ll g^{-4}$. Such improvements would be of tremendous interest in link with the quantum diffusion conjecture.

2. QUANTUM FRICTION MODEL

This section is devoted to the proof of our main results on the quantum friction model at weak coupling, cf. (1.20)-(1.24). We start with the construction of a suitable conjugate operator and with the proof of the Mourre estimate, thus establishing Theorem 1.5, before turning to consequences on the metastability of the embedded mass shell.

2.1. Construction of conjugate operator. The uncoupled fiber Hamiltonian splits as a sum of two multiplication operators,

$$H_0(P) = \frac{1}{2}(P - \mathrm{d}\Gamma(\xi))^2 + \mathrm{d}\Gamma(|k|),$$

where $\frac{1}{2}(P - d\Gamma(\xi))^2$ only involves the momentum coordinate $\xi \in \mathbb{R}^d$ dual to the spatial position and where $d\Gamma(|k|)$ only involves the momentum coordinate $k \in \mathbb{R}^q$ for vibrational degrees of freedom. We shall similarly construct a conjugate operator as a sum

$$A_P = B_P + D_1, \tag{2.1}$$

where B_P acts only on the variable ξ , and D_1 on the variable k. The commutator then splits formally as

$$[H_0(P), iA_P] = \frac{1}{2}[(P - d\Gamma(\xi))^2, iB_P] + [d\Gamma(|k|), iD_1],$$

so we are reduced to proving a Mourre estimate for both contributions separately. For the second contribution, $d\Gamma(|k|)$, recalling (1.1), a natural choice for the conjugate D_1 is the second quantization of the generator of dilations in the k-direction,

$$D_1 := \mathrm{d}\Gamma(d_1), \qquad d_1 := \frac{i}{2} \left(k \cdot \nabla_k + \nabla_k \cdot k \right), \tag{2.2}$$

which satisfies the following commutator identity,

$$[\mathrm{d}\Gamma(|k|), iD_1] = \mathrm{d}\Gamma(|k|). \tag{2.3}$$

It remains to construct a suitable conjugate B_P for $(P - d\Gamma(\xi))^2$. We start by briefly underlining the difficulty and motivating our unusual construction.

2.1.1. Motivation for construction of B_P . Recalling (1.1), a natural choice is to consider the generator of dilations in the ξ -direction around the minimizer of the symbol of $(P - d\Gamma(\xi))^2$. On the *n*-boson state space, the symbol takes the form

$$(\xi_1,\ldots,\xi_n) \mapsto \left(P-\sum_{j=1}^n \xi_j\right)^2,$$

which attains a minimum at

$$\xi_1 = \ldots = \xi_n = \frac{1}{n}P,\tag{2.4}$$

and we are then led to defining the following tentative conjugate operator on the n-boson state space,

$$\sum_{j=1}^{n} \frac{i}{2} \left((\xi_j - \frac{1}{n}P) \cdot \nabla_{\xi_j} + \nabla_{\xi_j} \cdot (\xi_j - \frac{1}{n}P) \right) = \sum_{j=1}^{n} \frac{i}{2} \left(\xi_j \cdot \nabla_{\xi_j} + \nabla_{\xi_j} \cdot \xi_j \right) - \frac{1}{n} \sum_{j=1}^{n} iP \cdot \nabla_{\xi_j}.$$

On the Fock space, this means

$$B_P := D_2 - N^{-\frac{1}{2}} \mathrm{d}\Gamma(iP \cdot \nabla_{\xi}) N^{-\frac{1}{2}}, \qquad (2.5)$$

in terms of the standard generator of dilations in the ξ -direction,

$$D_2 := \mathrm{d}\Gamma(d_2), \qquad d_2 := \frac{i}{2}(\xi \cdot \nabla_{\xi} + \nabla_{\xi} \cdot \xi), \qquad (2.6)$$

where we have implicitly defined the pseudo-inverse $N^{-\frac{1}{2}} := \bar{\Pi}_{\Omega} N^{-\frac{1}{2}} \bar{\Pi}_{\Omega}$ with some abuse of notation, recalling that $\bar{\Pi}_{\Omega} = 1 - \Pi_{\Omega}$ is the orthogonal projection on $\mathbb{C}\Omega^{\perp}$. For this choice, as expected from (1.2), we obtain

$$\frac{1}{2}[(P - d\Gamma(\xi))^2, iB_P] = \bar{\Pi}_{\Omega}(P - d\Gamma(\xi))^2 \bar{\Pi}_{\Omega} = (P - d\Gamma(\xi))^2 - P^2 \Pi_{\Omega}, \qquad (2.7)$$

which has exactly the desired behavior: indeed, combined with (2.3), it yields a Mourre estimate for $H_0(P)$ above the bottom of the spectrum.

Although this might look like the end of the story, this choice B_P does actually not suit our purposes as it behaves badly with respect to the fiber interaction Hamiltonian $\Phi(\rho)$: a direct computation shows that the commutator $[\Phi(\rho), iB_P]$ is not even relatively bounded when restricted to any fixed n-boson state space. The core of the problem is that B_P is not a second-quantization operator due to its dependence on the number operator, cf. (2.5), which is a direct consequence of the fact that the minimizer of the energy symbol depends the number n of bosons, cf. (2.4). While this issue seems unavoidable, we note that B_P is actually not the only possible choice. On the *n*-boson state space, the operator $N^{-1/2} d\Gamma(iP \cdot \nabla_{\xi}) N^{-1/2}$ in the definition (2.5) of B_P amounts to the arithmetic average $\frac{1}{n}\sum_{j=1}^{n}iP\cdot\nabla_{\xi_j}$ of coordinates $\{iP\cdot\nabla_{\xi_j}\}_{1\leq j\leq n}$, which we shall replace by the signed maximum of coordinates. This highly non-standard choice is directly inspired by our previous work [17] and we show that it does essentially not change the commutator relation (2.7), while behaving much better with the field operator $\Phi(\rho)$. The reason for this is intuitively clear: as expressed in Lemma 2.2 below, the maximum of a set coordinates satisfies better 'locality' properties with respect to creation and annihilation of coordinates than their arithmetic average.

2.1.2. Signed maximum and regularization. We turn to the construction of the suitable notion of signed maximum of coordinates $\{iP \cdot \nabla_{\xi_j}\}_{1 \leq j \leq n}$ on the *n*-boson state space, which will be used to replace the arithmetic mean $\frac{1}{n}\sum_{j=1}^{n}iP \cdot \nabla_{\xi_j}$ in (2.5). First, instead of momentum representation on \mathfrak{h} , we use position representation: we denote by $y := i\nabla_{\xi}$ the position coordinate dual to ξ , and we set $z := P \cdot y$ for the coordinate in the direction of the total momentum P. For all $n \geq 1$, we define the function $m_n : \mathbb{R}^n \to \mathbb{R}$ as the signed maximum of coordinates: for all $z_1, \ldots, z_n \in \mathbb{R}$, we set

$$m_n(z_1,\ldots,z_n) := z_{j_0}$$

where the index j_0 is chosen such that $|z_{j_0}| = \max_j |z_j|$. This is obviously well-defined on \mathbb{R}^n up to a null set. Equivalently, we can write

$$m_n(z_1, \dots, z_n) := (\max_j |z_j|) \operatorname{sgn} r_n(z_1, \dots, z_n),$$
(2.8)
$$r_n(z_1, \dots, z_n) := \max_j z_j + \min_j z_j,$$

where we take e.g. the convention sgn(0) = 0. This function is clearly symmetric with respect to the variables z_1, \ldots, z_n and has the following property.

Lemma 2.1. For all $n \ge 1$, the function m_n is continuous on $\mathbb{R}^n \setminus S_n$, where S_n stands for the hypersurface

$$\mathcal{S}_n := r_n^{-1}\{0\} = \left\{ z \in \mathbb{R}^n : \exists j_0 \neq j_1 \text{ such that } z_{j_0} = -z_{j_1} \text{ and } |z_{j_0}| = \max_j |z_j| \right\}.$$
(2.9)

In addition, there holds in the distributional sense

$$\sum_{j=1}^{n} \partial_j m_n = 1 + |\max_j z_j - \min_j z_j| \mathcal{H}_{\mathcal{S}_n} \ge 1,$$
 (2.10)

where $\mathcal{H}_{\mathcal{S}_n}$ stands for the (n-1)th-dimensional Hausdorff measure on \mathcal{S}_n .

Proof. The continuity of m_n is clear outside the zero locus S_n of r_n , and we turn to the second part of the statement. On $\mathbb{R}^n \setminus S_n$, we have $m_n(z_1, \ldots, z_n) = z_{j_0}$ with $|z_{j_0}| = \max_j |z_j|$, and thus $\sum_j \partial_j m_n = 1$. It remains to examine the jump of m_n on S_n . Given a point $z := (z_1, \ldots, z_n) \in \mathbb{R}^n$, we may assume $z_1 = \min_j z_j$ and $z_2 = \max_j z_j$ up to permuting coordinates, and we consider the line $\{z(t) := z + t(1, \ldots, 1) : t \in \mathbb{R}\}$. In view of (2.9), we note that this line intersects S_n at a single point: $z(t) \in S_n$ if and only if $t = -\frac{1}{2}(z_1 + z_2)$. The jump of m_n at this point along this line is easily checked to be $|z_1 - z_2|$, and the conclusion follows.

Next, we regularize m_n to smoothen the singular part of the derivative (2.10). We start with the following reformulation of m_n ,

$$m_n(z_1,...,z_n) = \frac{1}{2}(\max_j z_j + \min_j z_j) + \frac{1}{2}(\max_j z_j - \min_j z_j)\operatorname{sgn}(\max z_j + \min_j z_j),$$

where only the sign function needs to be regularized. Given $\delta > 0$, we choose a smooth odd function $\chi_{\delta} : \mathbb{R} \to [-1, 1]$ such that

$$\begin{aligned} \chi_{\delta}|_{(-\infty,-1]} &= -1, \qquad \chi_{\delta}|_{[1,\infty)} = 1, \qquad 0 \le \chi_{\delta}' \le 1 + \delta \quad \text{pointwise}, \\ \chi_{\delta}(s) \le s \quad \text{for } -1 \le s \le 0, \qquad \text{and} \qquad \chi_{\delta}(s) \ge s \quad \text{for } 0 \le s \le 1, \end{aligned}$$
(2.11)

 \Diamond

and we then define the following regularization of m_n ,

$$\widetilde{m}_{n;\delta}(z_1, \dots, z_p) := \frac{1}{2} (\max_j z_j + \min_j z_j) + \frac{1}{2} (\max_j z_j - \min_j z_j) \chi_{\delta} \left(\frac{\max_j z_j + \min_j z_j}{1 + \max_j z_j - \min_j z_j} \right), \quad (2.12)$$

which is obviously globally well-defined and continuous.¹ The denominator in the argument of χ_{δ} is easily understood: in order to make the derivative (2.10) uniformly bounded, it is not enough to regularize the sign function in a fixed neighborhood of S_n as the derivative would still produce an unbounded term due to the multiplication by $\max_j z_j - \min_j z_j$. In view of properties of χ_{δ} , a direct computation yields, instead of (2.10),

$$1 \le \sum_{j=1}^{n} \partial_j \widetilde{m}_{n;\delta} \le 2 + \delta, \tag{2.13}$$

and in addition, for all $r \ge 1$,

$$\left| \left(\sum_{j=1}^{n} \partial_{j} \right)^{r} \widetilde{m}_{n;\delta} \right| \lesssim_{\chi_{\delta}, r} 1, \qquad \left| \left(\sum_{j=1}^{n} z_{j} \partial_{j} \right)^{r} \widetilde{m}_{n;\delta} - \widetilde{m}_{n;\delta} \right| \lesssim_{\chi_{\delta}, r} 1.$$
(2.14)

In particular, note that $\widetilde{m}_{n;\delta}$ is smooth in the direction $(1, \ldots, 1)$. We also establish the following locality property: it constitutes the main difference with respect to arithmetic averages of coordinates $(z_1, \ldots, z_n) \mapsto \frac{1}{n} \sum_{j=1}^n z_j$, and it will be key to estimate commutators with field operators.

Lemma 2.2. For all $n \ge 0$, there holds for all $z, z_1, \ldots, z_n \in \mathbb{R}$,

$$\left|\widetilde{m}_{n+1;\delta}(z, z_1, \dots, z_n) - \widetilde{m}_{n;\delta}(z_1, \dots, z_n)\right| \le 2|z| + 1.$$

Proof. As $|\chi_{\delta}| \leq 1$, the definition of $\widetilde{m}_{n;\delta}$ ensures

$$|\widetilde{m}_{n;\delta}(z_1,\ldots,z_n)| \le \max_j |z_j|$$

which trivially yields the conclusion in case $\max_j |z_j| \le |z| \lor \frac{1}{2}$. It thus remains to consider the case $\max_j |z_j| > |z| \lor \frac{1}{2}$. Up to permuting coordinates, we can assume $z_1 = \min_j z_j$ and $z_2 = \max_j z_j$. In the case $z_1 \le z \le z_2$, we simply find

$$\widetilde{m}_{n+1;\delta}(z, z_1, \ldots, z_n) = \widetilde{m}_{n;\delta}(z_1, \ldots, z_n),$$

and the conclusion follows. It remains to treat the case $z_1 \leq z_2 \leq z$, while the symmetric case $z \leq z_1 \leq z_2$ is similar. Given $z_1 \leq z_2 \leq z$, the assumption $\max_j |z_j| > |z| \lor \frac{1}{2}$ implies $z_1 < -|z| \lor \frac{1}{2}$, and thus in particular $\frac{z_2+z_1}{1+z_2-z_1} \leq \frac{z+z_1}{1+z-z_1} \leq 0$. In addition, there holds $\chi_{\delta}\left(\frac{y+z_1}{1+y-z_1}\right) = -1$ whenever $z_1 \leq y \leq -\frac{1}{2}$. Using this together with properties of χ_{δ} ,

¹A similar construction was first used in our previous work [17]. Note however the following slight mistake in [17, Section 5.6]: we forgot to add 1 in the denominator of the argument of χ_{δ} , which then actually poses regularity issues at the origin in \mathbb{R}^n .

we may then estimate

$$\begin{split} \left| \widetilde{m}_{n+1;\delta}(z, z_1, \dots, z_n) - \widetilde{m}_{n;\delta}(z_1, \dots, z_n) \right| \\ &= \left| \frac{1}{2}(z - z_2) \left(1 + \chi_{\delta} \left(\frac{z_2 + z_1}{1 + z_2 - z_1} \right) \right) + \frac{1}{2}(z - z_1) \left(\chi_{\delta} \left(\frac{z + z_1}{1 + z - z_1} \right) - \chi_{\delta} \left(\frac{z_2 + z_1}{1 + z_2 - z_1} \right) \right) \\ &\leq \left| \frac{1}{2}(z - z_2) \left(1 + \chi_{\delta} \left(\frac{z_2 + z_1}{1 + z_2 - z_1} \right) \right) + \frac{1}{2}(z - z_1) \left(1 + \chi_{\delta} \left(\frac{z + z_1}{1 + z - z_1} \right) \right) \right) \\ &\leq \left| \frac{1}{2}(z - z_2) \mathbb{1}_{z_2 \ge -\frac{1}{2}} + \frac{1}{2}(z - z_1) \left(1 + \frac{z + z_1}{1 + z - z_1} \right) \mathbb{1}_{z \ge -\frac{1}{2}} \right) \\ &\leq \left| \frac{3}{2}(|z| + \frac{1}{2}), \end{split}$$

as claimed.

2.1.3. Back to conjugate operator. With the above at hand, we can turn to the suitable replacement for the second term, $N^{-1/2} d\Gamma (iP \cdot \nabla_{\xi}) N^{-1/2}$, in the tentative choice (2.5) of the conjugate operator. For all $n \geq 0$, we define the operator $M_{P,n;\delta}$ on the *n*-boson state space as the multiplication with the function

$$(k_1, y_1, \ldots, k_n, y_n) \mapsto \widetilde{m}_{n;\delta}(P \cdot y_1, \ldots, P \cdot y_n),$$

using position representation $y = i \nabla_{\xi}$ on \mathfrak{h} , and we define

$$M_{P;\delta} = \bigoplus_{n=1}^{\infty} M_{P,n;\delta}$$
 on $\mathcal{H}^{\mathrm{f}} = \bigoplus_{n=0}^{\infty} \Gamma_s^{(n)}(\mathfrak{h})$

Coming back to (2.1) and (2.5), we then define the following modified conjugate operator,

$$A_{P;\delta} := 2D_1 + D_2 - \frac{2}{3+\delta}M_{P;\delta} \quad \text{on } \mathcal{H}^{\mathrm{f}},$$
 (2.15)

where we recall that D_1 and D_2 stand for the generators of dilations in the k-direction and the ξ -direction, respectively,

$$D_1 = d\Gamma(d_1), \qquad d_1 = \frac{i}{2}(k \cdot \nabla_k + \nabla_k \cdot k),$$

$$D_2 = d\Gamma(d_2), \qquad d_2 = \frac{i}{2}(\xi \cdot \nabla_\xi + \nabla_\xi \cdot \xi).$$

The reason for the factor $\frac{2}{3+\delta}$ in front of $M_{P;\delta}$ in (2.15) is the following: the computation of the relevant commutators involves the derivative $\sum_{j=1}^{n} \partial_j \tilde{m}_{n;\delta}$, which in view of the regularization $\tilde{m}_{n;\delta}$ of m_n is not equal to 1 almost everywhere but takes values in the whole interval $[1, 2+\delta]$ close to the hypersurface S_n (compare (2.10) to (2.13)). Symbols are thus deformed in the computation of commutators, and the choice $2D_1+D_2-M_P$ would actually fail to provide a Mourre estimate close to the eigenvalue $\frac{1}{2}P^2$. Definition (2.15) is precisely meant to best overcome this issue.

By definition, the operator $A_{P;\delta}$ commutes with the number operator. Given its action on *n*-boson state space, it is clearly essentially self-adjoint on C^{f} , and we show that it generates an explicit unitary group that preserves the domain of fiber Hamiltonians.

Lemma 2.3. The operator $A_{P;\delta}$ is essentially self-adjoint on C^{f} and its closure generates a unitary group $\{e^{itA_{P;\delta}}\}_{t\in\mathbb{R}}$ on \mathcal{H}^{f} , which commutes with the number operator and has the following explicit action: for all $n \geq 1$ and $u_n \in \Gamma_s^{(n)}(\mathfrak{h})$,

$$(e^{itA_{P;\delta}}u_n)(k_1, y_1, \dots, k_n, y_n) = \exp\left(-\frac{2i}{3+\delta} \int_0^t \widetilde{m}_{n;\delta}(P \cdot e^s y_1, \dots, P \cdot e^s y_n) \, ds\right) \\ \times e^{tn(\frac{d}{2}-q)}u_n(e^{-2t}k_1, e^t y_1, \dots, e^{-2t}k_n, e^t y_n),$$

where on $\mathfrak{h} = L^2(\mathbb{R}^q \times \mathbb{R}^d)$ we use momentum representation in the first variable and position representation in the second. In particular, the domain \mathcal{D} of fiber Hamiltonians (1.25) is invariant under this group action.

Proof. For all $n \geq 1$, consider the family $\{U_{P,n;\delta}^t\}_{t\in\mathbb{R}}$ of operators on $\Gamma_s^{(n)}(\mathfrak{h})$, defined by the above formula,

$$(U_{P,n;\delta}^{t}u_{n})(k_{1},y_{1},\ldots,k_{n},y_{n}) := \exp\left(-\frac{2i}{3+\delta}\int_{0}^{t}\widetilde{m}_{n;\delta}(P \cdot e^{s}y_{1},\ldots,P \cdot e^{s}y_{n})\,ds\right) \\ \times e^{tn(\frac{d}{2}-q)}\,u_{n}(e^{-2t}k_{1},e^{t}y_{1},\ldots,e^{-2t}k_{n},e^{t}y_{n}).$$

This defines a unitary group on $\Gamma_s^{(n)}(\mathfrak{h})$. In addition, for all $u_n \in C_c^{\infty}(\mathbb{R}^q \times \mathbb{R}^d)^{\otimes_s n}$, we note that the following convergence holds in $\Gamma_s^{(n)}(\mathfrak{h})$,

$$\lim_{t \downarrow 0} \frac{1}{t} (U_{P,n;\delta}^t u_n - u_n) \\ = -\sum_{j=1}^n (k_j \cdot \nabla_{k_j} + \nabla_{k_j} \cdot k_j) u_n + \frac{1}{2} \sum_{j=1}^n (y_j \cdot \nabla_{y_j} + \nabla_{y_j} \cdot y_j) u_n - \frac{2i}{3+\delta} M_{P,n;\delta} u_n,$$

where the right-hand side coincides with $iA_{P;\delta}u_n$ since we have in position representation

$$d_2 = \frac{i}{2}(\xi \cdot \nabla_{\xi} + \nabla_{\xi} \cdot \xi) = \frac{1}{2i}(y \cdot \nabla_y + \nabla_y \cdot y).$$

This proves that $\{U_{P,n;\delta}^t\}_{t\in\mathbb{R}}$ is a unitary C_0 -group on $\Gamma_s^{(n)}(\mathfrak{h})$ and that its self-adjoint generator coincides with $A_{P;\delta}$ on its core $C_c^{\infty}(\mathbb{R}^q \times \mathbb{R}^d)^{\otimes_s n}$. The conclusion follows. \Box

Next, we show the relative boundedness of commutators of the uncoupled fiber Hamiltonian $H_0(P)$ with the above-constructed conjugate operator $A_{P;\delta}$. Combined with Lemma 2.3, this actually proves Theorem 1.5(i), further noting that the $C^{\infty}(A_{P;\delta})$ -regularity property follows by applying the sufficient criterion in Lemma A.3.

Lemma 2.4. For all $s \geq 1$, the s-th iterated commutator $\operatorname{ad}_{iA_{P;\delta}}^s(H_0(P))$ extends as an $H_0(P)$ -bounded self-adjoint operator with domain $\mathcal{D} = \mathcal{D}(H_0(P))$.

Proof. For all $n \geq 1$, we define the operators $M'_{P,n;\delta}$ and $M''_{P,n;\delta}$ on $\Gamma_s^{(n)}(\mathfrak{h})$ as the multiplications with the functions

$$(k_1, y_1, \dots, k_n, y_n) \mapsto \left(\sum_{j=1}^n \partial_j \widetilde{m}_{n;\delta} \right) (P \cdot y_1, \dots, P \cdot y_n), (k_1, y_1, \dots, k_n, y_n) \mapsto \left(\sum_{j,l=1}^n \partial_{j,l} \widetilde{m}_{n;\delta} \right) (P \cdot y_1, \dots, P \cdot y_n),$$

respectively, and we set

$$M'_{P;\delta} := \bigoplus_{n=1}^{\infty} M'_{P,n;\delta}, \qquad M''_{P;\delta} := \bigoplus_{n=1}^{\infty} M''_{P,n;\delta}, \qquad \text{on } \mathcal{H}^{\mathrm{f}} = \bigoplus_{n=0}^{\infty} \Gamma_s^{(n)}(\mathfrak{h}).$$

A direct computation yields in these terms

$$[\mathrm{d}\Gamma(\nabla_y), M_{P;\delta}] = PM'_{P;\delta}, \qquad [\mathrm{d}\Gamma(\nabla_y), M'_{P;\delta}] = PM''_{P;\delta}.$$

By definition (2.15) of $A_{P;\delta}$, recalling that $\xi = -i\nabla_y$, we compute in the sense of forms on \mathcal{C}^{f} ,

$$[H_0(P), iA_{P;\delta}] = 2\mathrm{d}\Gamma(|k|) - \mathrm{d}\Gamma(\xi) \cdot (P - \mathrm{d}\Gamma(\xi)) + \frac{1}{3+\delta}P \cdot \Big((P - \mathrm{d}\Gamma(\xi))M'_{P;\delta} + M'_{P;\delta}(P - \mathrm{d}\Gamma(\xi))\Big),$$

which can be reorganized as

$$[H_0(P), iA_{P;\delta}] = 2d\Gamma(|k|) + (P - d\Gamma(\xi))^2 + \frac{1}{3+\delta}P \cdot \left((P - d\Gamma(\xi))(M'_{P;\delta} - \frac{3+\delta}{2}) + (M'_{P;\delta} - \frac{3+\delta}{2})(P - d\Gamma(\xi)) \right).$$
(2.16)

Alternatively, further using $[d\Gamma(\nabla_y), M'_{P;\delta}] = PM''_{P;\delta}$, and recognizing $H_0(P)$ in the righthand side, we get

$$[H_0(P), iA_{P;\delta}] = 2H_0(P) + \frac{2}{3+\delta}(M'_{P;\delta} - \frac{3+\delta}{2})P \cdot (P - d\Gamma(\xi)) + \frac{i}{3+\delta}P^2 M''_{P;\delta}.$$
 (2.17)

As (2.13) yields $1 \leq M'_{P;\delta} \leq 2 + \delta$ and $|M''_{P;\delta}| \lesssim_{\chi_{\delta}} 1$, we find for all $u \in \mathcal{C}^{\mathfrak{t}}$,

$$\begin{aligned} \left\| \frac{2}{3+\delta} (M'_{P;\delta} - \frac{3+\delta}{2}) P \cdot (P - \mathrm{d}\Gamma(\xi)) u \right\| + \left\| \frac{i}{3+\delta} P^2 M''_{P;\delta} u \right\| \\ \lesssim_{\chi_{\delta}} |P| \| (P - \mathrm{d}\Gamma(\xi)) u \| + P^2 \| u \| \\ \leq |P| \| H_0(P)^{\frac{1}{2}} u \| + P^2 \| u \|. \end{aligned}$$

Combined with (2.17), this shows that $[H_0(P), iA_{P;\delta}]$ is equal to $2H_0(P)$ up to an infinitesimal perturbation. By the Kato–Rellich theorem, we deduce that the commutator $\operatorname{ad}_{iA_{P;\delta}}(H_0(P)) = [H_0(P), iA_{P;\delta}]$ extends as an $H_0(P)$ -bounded self-adjoint operator with domain $\mathcal{D} = \mathcal{D}(H_0(P))$. Similarly computing iterated commutators and appealing to (2.14), the conclusion easily follows; we skip the detail.

2.2. Mourre estimate. We turn to the proof of the Mourre estimate for $H_0(P)$. This amounts to showing that the commutator identity (2.7) is essentially preserved for the modified conjugate operator (2.15). Due to the deformation of commutators, however, we only manage to cover energy intervals above $\frac{1}{18}P^2$. Up to renaming ε, δ , this proves Theorem 1.5(ii).

Lemma 2.5. For all $\varepsilon > 0$, the commutator $[H_0(P), iA_{P;\delta}]$ satisfies the following Mourre estimate on $J_{\varepsilon} := \left[(\frac{1+\delta}{3+\delta} + \varepsilon)^2 \frac{1}{2} P^2, \infty \right)$,

$$\mathbb{1}_{J_{\varepsilon}}(H_0(P))[H_0(P), iA_{P;\delta}]\mathbb{1}_{J_{\varepsilon}}(H_0(P)) \geq \varepsilon(\frac{1}{3} + \varepsilon)P^2\mathbb{1}_{J_{\varepsilon}}(H_0(P)) - P^2\Pi_{\Omega}.$$

Proof. We split the proof into two steps.

Step 1. Proof that for all $\alpha > 0$,

$$[H_0(P), iA_{P;\delta}] \ge \bar{\Pi}_{\Omega} \left(2(1 - \frac{1}{2\alpha})H_0(P) - \frac{\alpha}{2} \left(\frac{1+\delta}{3+\delta}\right)^2 P^2 \right) \bar{\Pi}_{\Omega}.$$
(2.18)

As the vacuum state Ω is an eigenvector of $H_0(P)$, it belongs to the kernel of the commutator $[H_0(P), iA_{P;\delta}]$, and it suffices to establish this lower bound (2.18) on $\mathcal{C}^{\mathrm{f}} \cap \mathbb{C}\Omega^{\perp}$. Given $u \in \mathcal{C}^{\mathrm{f}} \cap \mathbb{C}\Omega^{\perp}$, starting from identity (2.16), we can bound

$$\left\langle u, [H_0(P), iA_{P;\delta}]u \right\rangle \ge 2 \left\langle u, \mathrm{d}\Gamma(|k|)u \right\rangle + \left\langle u, (P - \mathrm{d}\Gamma(\xi))^2 u \right\rangle - \frac{2}{3+\delta} |P| \| (P - \mathrm{d}\Gamma(\xi))u \| \| (M'_{P;\delta} - \frac{3+\delta}{2})u \|,$$

and thus, recalling that (2.13) implies $1 \leq M'_{P;\delta} \leq 2 + \delta$, we get

 $\left\langle u, [H_0(P), iA_{P,\delta}]u \right\rangle \geq 2\left\langle u, d\Gamma(|k|)u \right\rangle + \left\langle u, (P - d\Gamma(\xi))^2 u \right\rangle - \frac{1+\delta}{3+\delta} |P| \| (P - d\Gamma(\xi))u \| \| u \|.$ For all $\alpha > 0$, Young's inequality then yields

$$\begin{aligned} \left\langle u, [H_0(P), iA_{P;\delta}] u \right\rangle &\geq 2 \left\langle u, \mathrm{d}\Gamma(|k|)u \right\rangle + (1 - \frac{1}{2\alpha}) \left\langle u, (P - \mathrm{d}\Gamma(\xi))^2 u \right\rangle - \frac{\alpha}{2} \left(\frac{1+\delta}{3+\delta}\right)^2 P^2 \|u\|^2 \\ &\geq 2(1 - \frac{1}{2\alpha}) \left\langle u, H_0(P)u \right\rangle - \frac{\alpha}{2} \left(\frac{1+\delta}{3+\delta}\right)^2 P^2 \|u\|^2, \end{aligned}$$

that is, (2.18).

Step 2. Conclusion.

Given $E \ge 0$, applying (2.18) to $\mathbb{1}_{[E,\infty)}(H_0(P))u$, we get

$$\begin{split} \left\langle \mathbb{1}_{[E,\infty)}(H_0(P))u, [H_0(P), iA_{P;\delta}] \mathbb{1}_{[E,\infty)}(H_0(P))u \right\rangle \\ &\geq \left(2E(1-\frac{1}{2\alpha}) - \frac{\alpha}{2} \left(\frac{1+\delta}{3+\delta}\right)^2 P^2 \right) \|\mathbb{1}_{[E,\infty)}(H_0(P))u\|^2 - P^2 \|\Pi_{\Omega} u\|^2. \end{split}$$

Hence, optimizing with respect to $\alpha > 0$,

$$\begin{split} \left\langle \mathbb{1}_{[E,\infty)}(H_0(P))u, [H_0(P), iA_{P;\delta}]\mathbb{1}_{[E,\infty)}(H_0(P))u \right\rangle \\ &\geq \sqrt{2E} \left(\sqrt{2E} - \frac{1+\delta}{3+\delta}|P|\right) \|\mathbb{1}_{[E,\infty)}(H_0(P))u\|^2 - P^2 \|\Pi_{\Omega} u\|^2, \\ \text{d the stated Mourre estimate follows.} \end{split}$$

and the stated Mourre estimate follows.

2.3. **Regularity of the interaction.** We turn to the regularity of the fiber interaction Hamiltonian $\Phi(\rho)$ with respect to $A_{P;\delta}$, thus establishing Theorem 1.5(iii). While this result would fail for the naïve choice (2.5) of the conjugate, it crucially requires our special definition of regularized signed maximum, and the proof builds mainly on Lemma 2.2.

Lemma 2.6. Let the interaction kernel ρ satisfy Assumption (Reg_{ν}) for some $\nu \geq 1$. Then, for all $0 \leq s \leq \nu$, the s-th iterated commutator $\operatorname{ad}_{iA_{P;\delta}}^{s}(\Phi(\rho))$ extends as a $d\Gamma(|k|)^{1/2}$ bounded self-adjoint operator. \Diamond

Proof. We split the proof into two steps.

Step 1. Analysis of the first commutator.

By definition (2.15) of $A_{P;\delta}$, the first commutator can be split as

$$[\Phi(\rho), iA_{P;\delta}] = 2[\Phi(\rho), id\Gamma(d_1)] + [\Phi(\rho), id\Gamma(d_2)] - \frac{2}{3+\delta}[\Phi(\rho), iM_{P;\delta}],$$
(2.19)

and thus, as the first two terms involve second-quantization operators,

$$[\Phi(\rho), iA_{P;\delta}] = -2\Phi(id_1\rho) - \Phi(id_2\rho) - \frac{2}{3+\delta}[\Phi(\rho), iM_{P;\delta}].$$

Standard estimates ensure that the first two terms $\Phi(id_1\rho)$ and $\Phi(id_2\rho)$ are $d\Gamma(|k|)^{1/2}$ bounded provided that $(1+|k|^{-1/2})d_1\rho$ and $(1+|k|^{-1/2})d_2\rho$ belong to $L^2(\mathbb{R}^q \times \mathbb{R}^d)$, hence in particular provided that ρ satisfies Assumption (Reg₁).

It remains to estimate the commutator $[\Phi(\rho), iM_{P;\delta}]$. As $\Phi(\rho) = a^*(\rho) + a(\rho)$, it actually suffices by symmetry to estimate $[a^*(\rho), iM_{P;\delta}]$. We recall the standard definition of the creation operator: for all $n \ge 0$ and $u_n \in \Gamma_s^{(n)}(\mathfrak{h})$,

$$\left(a^*(\rho)u_n\right)\left((k_l,\xi_l)_{1\leq l\leq n+1}\right) = \frac{1}{\sqrt{n+1}}\sum_{j=1}^{n+1}\rho(k_j,\xi_j)\,u_n\left((k_l,\xi_l)_{l\in\{1,\dots,n+1\}\setminus\{j\}}\right),$$

or alternatively, using position representation $y = i \nabla_{\xi}$ in the second variable,

$$(a^*(\rho)u_n) ((k_l, y_l)_{1 \le l \le n+1}) = \frac{1}{\sqrt{n+1}} \sum_{j=1}^{n+1} \tilde{\rho}(k_j, y_j) u_n ((k_l, y_l)_{l \in \{1, \dots, n+1\} \setminus \{j\}}),$$

where $\tilde{\rho}$ stands for the partial inverse Fourier transform $\tilde{\rho}(k, y) := \int_{\mathbb{R}^d} e^{iy \cdot \xi} \rho(k, \xi) \, d\xi$. In these terms, the commutator with $M_{P;\delta}$ takes the explicit form

$$\left([a^*(\rho), iM_{P;\delta}] u_n \right) \left((k_l, y_l)_{1 \le l \le n+1} \right) \\
= \frac{1}{\sqrt{n+1}} \sum_{j=1}^{n+1} \left(i \widetilde{m}_{n;\delta} \left((P \cdot y_l)_{l \in \{1, \dots, n+1\} \setminus \{j\}} \right) - i \widetilde{m}_{n+1;\delta} \left((P \cdot y_l)_{1 \le l \le n+1} \right) \right) \\
\times \widetilde{\rho}(k_j, y_j) u_n \left((k_l, y_l)_{l \in \{1, \dots, n+1\} \setminus \{j\}} \right). \quad (2.20)$$

Appealing to Lemma 2.2 to estimate the difference between $\widetilde{m}_{n:\delta}$ and $\widetilde{m}_{n+1:\delta}$, we deduce

$$|[a^{*}(\rho), iM_{P;\delta}]u_{n}| \leq 2|P|\tilde{a}^{*}(|y\tilde{\rho}|)|u_{n}| + \tilde{a}^{*}(|\tilde{\rho}|)|u_{n}|,$$

where we use the short-hand notation $\tilde{a}^*(\tilde{\sigma}) := a^*(\sigma)$. Standard estimates then entail that the commutator $[a^*(\rho), iM_{P;\delta}]$ is $d\Gamma(|k|)^{1/2}$ -bounded provided that $(1+|k|^{-1/2})y\tilde{\rho}$ belongs to $L^2(\mathbb{R}^q \times \mathbb{R}^d)$. This is equivalent to requiring that $(1+|k|^{-1/2})\nabla_{\xi}\rho$ belongs to $L^2(\mathbb{R}^q \times \mathbb{R}^d)$, which holds in particular provided that ρ satisfies (Reg₁). The conclusion follows.

Step 2. Analysis of iterated commutators.

/

As in (2.19), we start by decomposing the commutator with $iA_{P;\delta}$ in terms of commutators with iD_1 , iD_2 , and $iM_{P;\delta}$. Upon iteration, we are then led to estimating products of ad_{iD_1} , ad_{iD_2} , and $ad_{iM_{P;\delta}}$, applied to $\Phi(\rho) = a^*(\rho) + a(\rho)$. In line with (2.20), we argue that such expressions are explicit and thus easily estimated. By symmetry, as in Step 1, it suffices to consider commutators applied to $a^*(\rho)$. Iterating (2.20), we find for all $s \ge 0$,

$$\left(\operatorname{ad}_{iM_{P;\delta}}^{s}(a^{*}(\rho))u_{n} \right) \left((k_{l}, y_{l})_{1 \leq l \leq n+1} \right)$$

$$= \frac{1}{\sqrt{n+1}} \sum_{j=1}^{n+1} \left(i\widetilde{m}_{n;\delta} \left((P \cdot y_{l})_{l \in \{1, \dots, n+1\} \setminus \{j\}} \right) - i\widetilde{m}_{n+1;\delta} \left((P \cdot y_{l})_{1 \leq l \leq n+1} \right) \right)^{s}$$

$$\times \tilde{\rho}(k_{j}, y_{j}) u_{n} \left((k_{l}, y_{l})_{l \in \{1, \dots, n+1\} \setminus \{j\}} \right).$$
(2.21)

Further taking the commutator with $iD_1 = id\Gamma(d_1)$ and $iD_2 = id\Gamma(d_2)$, we easily find

$$\operatorname{ad}_{iD_{1}}\left(\operatorname{ad}_{iM_{P;\delta}}^{s}(a^{*}(\rho))\right) = \operatorname{ad}_{iM_{P;\delta}}^{s}\left(a^{*}(k \cdot \nabla_{k}\rho)\right),$$

$$\operatorname{ad}_{iD_{2}}\left(\operatorname{ad}_{iM_{P;\delta}}^{s}(a^{*}(\rho))\right) = \operatorname{ad}_{iM_{P;\delta}}^{s}\left(a^{*}\left((\xi \cdot \nabla_{\xi} - s)\rho\right)\right) - s \operatorname{ad}_{iR_{P;\delta}}\left(\operatorname{ad}_{iM_{P;\delta}}^{s-1}\left(a^{*}(\rho)\right)\right),$$

$$(2.22)$$

where the operator $R_{P;\delta}$ in the last term is defined as follows: for all $n \ge 1$, we set

$$\widetilde{r}_{n;\delta}(z_1,\ldots,z_n) := \sum_{j=1}^n z_j \partial_j \widetilde{m}_{n;\delta} - \widetilde{m}_{n;\delta}$$

we define the operator $R_{P,n;\delta}$ on $\Gamma_s^{(n)}(\mathfrak{h})$ as the multiplication with the function

$$(k_1, y_1, \ldots, k_n, y_n) \mapsto \widetilde{r}_{n;\delta}(P \cdot y_1, \ldots, P \cdot y_n)$$

and we set $R_{P;\delta} := \bigoplus_{n=1}^{\infty} R_{P,n;\delta}$ on \mathcal{H}^{f} . In view of (2.14), the function $\tilde{r}_{n;\delta}$ is bounded uniformly in n, and the term involving $\mathrm{ad}_{iR_{P;\delta}}$ in (2.22) can thus be viewed as a better-behaved

lower-order remainder. Up to such a remainder, identities (2.22) show that products of ad_{iD_1} , ad_{iD_2} , and $\mathrm{ad}_{iM_{P;\delta}}$, when applied to $a^*(\rho)$, can be reduced to powers of $\mathrm{ad}_{iM_{P;\delta}}$ up to transforming ρ . The conclusion easily follows from this observation and we skip the detail.

2.4. Consequences of Mourre estimate. Given a total momentum $P \neq 0$, we turn to the proof of Corollary 1.6. By items (i) and (iii) in Theorem 1.5, the sufficient criterion in Lemma A.3 ensures that the coupled fiber Hamiltonian $H_g(P)$ is of class $C^{\infty}(A_{P;\delta})$ for all g. Next, by Theorem 1.5(iii), for all $\varepsilon > 0$, Lemma A.6 allows to infer that $H_g(P)$ satisfies a Mourre estimate with respect to $A_{P;\delta}$ on the energy interval

$$\left(\frac{1}{18}P^2 + \delta + \varepsilon + \frac{gC_P}{\varepsilon}, \infty\right),$$

for some constant C_P . Taking δ arbitrarily small and optimizing in ε , we deduce that $H_g(P)$ satisfies a Mourre estimate on any compact subinterval of

$$J_{P,g} := \left(\frac{1}{18}P^2 + \sqrt{g}C_P, \infty\right)$$

Moreover, the Mourre estimate is strict outside $K_{P,g} := \left[\frac{1}{2}P^2 - gC_P, \frac{1}{2}P^2 + gC_P\right]$. We may then appeal to Theorem A.5, which states that $H_g(P)$ has no singular spectrum and at most a finite number of eigenvalues in $J_{P,g}$, and has no eigenvalue in $J_{P,g} \setminus K_{P,g}$. In order to exclude the existence of eigenvalues in $K_{P,g}$, we appeal to Theorem A.7, which states the instability of the uncoupled eigenvalue $\frac{1}{2}P^2$ provided that Fermi's condition (A.8) holds. Altogether, this proves item (i) of Corollary 1.6, and item (ii) follows by further applying Theorem A.8. It remains to make Fermi's condition (A.8) more explicit for the model at hand, which is the purpose of the following lemma (see also [11, Lemma 6.7]).

Lemma 2.7. For all $P \neq 0$, we have

$$\begin{split} \lim_{\varepsilon \downarrow 0} \left\langle \Omega, \, \Phi(\rho) \bar{\Pi}_{\Omega} \big(H_0(P) - \frac{1}{2} P^2 - i\varepsilon \big)^{-1} \bar{\Pi}_{\Omega} \Phi(\rho) \Omega \right\rangle \\ &= (2\pi)^{-d} \, \text{p.v.} \int_0^\infty (t - \frac{1}{2} P^2)^{-1} \bigg(\int_{|k| \le t} \int_{\{\xi : \frac{1}{2}(P - \xi)^2 = t - |k|\}} \frac{|\rho(k,\xi)|^2}{\sqrt{(P - \xi)^2 + 1}} \mathrm{d}\mathcal{H}_{d-1}(\xi) \mathrm{d}k \bigg) \mathrm{d}t \\ &+ \frac{i}{2} (2\pi)^{1-d} \int_{|k| \le \frac{1}{2} P^2} \int_{\{\xi : \frac{1}{2}(P - \xi)^2 = \frac{1}{2} P^2 - |k|\}} \frac{|\rho(k,\xi)|^2}{\sqrt{(P - \xi)^2 + 1}} \mathrm{d}\mathcal{H}_{d-1}(\xi) \mathrm{d}k, \end{split}$$

where \mathcal{H}_{d-1} stands for the (d-1)th-dimensional Hausdorff measure. In particular, the imaginary part is positive if ρ does not vanish.

Proof. For any $\varepsilon > 0$, we compute

$$\begin{split} &\left\langle \Omega, \, \Phi(\rho) \bar{\Pi}_{\Omega} \left(H_0(P) - \frac{1}{2} P^2 - i\varepsilon \right)^{-1} \bar{\Pi}_{\Omega} \Phi(\rho) \Omega \right\rangle \\ &= \left\langle a^*(\rho) \Omega, \, \left(H_0(P) - \frac{1}{2} P^2 - i\varepsilon \right)^{-1} a^*(\rho) \Omega \right\rangle \\ &= \iint_{\mathbb{R}^q \times \mathbb{R}^d} |\rho(k,\xi)|^2 \Big(H_0^{(1)}(P;k,\xi) - \frac{1}{2} P^2 - i\varepsilon \Big)^{-1} \mathrm{d}k \mathrm{d}\xi \end{split}$$

where $H_0^{(1)}(P;k,\xi) := \frac{1}{2}(P-\xi)^2 + |k|$ is the symbol of $H_0(P)$ on the single-boson state space. As this symbol is Lipschitz continuous, the coarea formula yields

$$\left\langle \Omega, \, \Phi(\rho) \bar{\Pi}_{\Omega} \left(H_0(P) - \frac{1}{2} P^2 - i\varepsilon \right)^{-1} \bar{\Pi}_{\Omega} \Phi(\rho) \Omega \right\rangle$$

= $(2\pi)^{-q-d} \int_0^\infty \left(t - \frac{1}{2} P^2 - i\varepsilon \right)^{-1} \left(\int_{\{(k,\xi): H_0^{(1)}(P;k,\xi) = t\}} \frac{|\rho(k,\xi)|^2}{|\nabla_{k,\xi} H_0^{(1)}(P;k,\xi)|} \mathrm{d}\mathcal{H}_{q+d-1}(k,\xi) \right) \mathrm{d}t,$

where \mathcal{H}_{q+d-1} stands for the (q+d-1)th-dimensional Hausdorff measure and where we note that the integrand is summable. As $|\nabla_{k,\xi}H_0^{(1)}(P;k,\xi)| = \sqrt{(P-\xi)^2+1}$ is non-degenerate, the conclusion easily follows from the Plemelj formula.

3. TRANSLATION-INVARIANT MASSIVE NELSON MODEL

This section is devoted to the proof of our main results on the translation-invariant Nelson model with massive bosons at small coupling, cf. (1.3)-(1.9). We start by describing the energy-momentum spectrum for uncoupled Hamiltonians, in particular proving Lemma 1.1, and we establish some important properties of energy thresholds. Next, we turn to the construction of a conjugate operator in the weak-coupling regime, thus establishing Theorem 1.2. As explained in the introduction, a suitable modification of our first choice of conjugate will be needed to ensure C^{∞} -regularity. The modification procedure is presented in Section 3.4 below, further building on our constructions in Section 2.1, and we believe that it is of independent interest for other massive QFT models.

3.1. **Spectrum of uncoupled Hamiltonians.** We start with the proof of Lemma 1.1, that is, the characterization of the spectrum of uncoupled fiber Hamiltonians. More precisely, we establish the following result.

Lemma 3.1. Consider the translation-invariant Nelson model with massive bosons m > 0, cf. (1.3)–(1.9). Given a total momentum $P \in \mathbb{R}^d$, the uncoupled fiber Hamiltonian $H_0(P)$ commutes with the number operator N and thus splits as a direct sum (1.14). There holds

$$H_0(P)\Omega = \frac{1}{2}P^2\Omega,$$

and for all $n \geq 1$ the restriction $H_0^{(n)}(P) = H_0(P)|_{\Gamma_c^{(n)}(\mathfrak{h})}$ satisfies

$$\sigma_{\rm ac}\big(H_0^{(n)}(P)\big) = \big[E_0^{(n)}(P),\infty\big), \qquad \sigma_{\rm pp}\big(H_0^{(n)}(P)\big) = \sigma_{\rm sc}\big(H_0^{(n)}(P)\big) = \varnothing, \tag{3.1}$$

where the n-boson energy threshold $E_0^{(n)}(P)$ is given by

$$E_0^{(n)}(P) := \frac{1}{2}c(n,P)^2 + \sqrt{m^2n^2 + (|P| - c(n,P))^2},$$
(3.2)

in terms of the unique solution $c(n, P) \in [0, 1)$ of the implicit equation

$$c(n,P) = \frac{|P| - c(n,P)}{\sqrt{m^2 n^2 + (|P| - c(n,P))^2}}.$$
(3.3)

Proof. In view of (1.14), it suffices to analyze separately the spectrum of restrictions on each *n*-boson state space. For $n \ge 1$, the restriction $H_0^{(n)}(P)$ is a multiplication operator in momentum coordinates, with symbol

$$H_0^{(n)}(P;k_1,\ldots,k_n) := \frac{1}{2} \left(P - \sum_{j=1}^n k_j \right)^2 + \sum_{j=1}^n \omega(k_j).$$
(3.4)

Its spectrum is thus absolutely continuous and coincides with the essential image of this symbol, which in this case obviously takes the form (3.1) with

$$E_0^{(n)}(P) := \min_{k_1,\dots,k_n \in \mathbb{R}^d} H_0^{(n)}(P;k_1,\dots,k_n)$$

A straightforward computation shows that this minimum is attained at

$$k_1 = \ldots = k_n = k_\star(n, P) := \frac{1}{n} (|P| - c(n, P)) \frac{P}{|P|},$$
 (3.5)

. .

. .

where $c(n, P) \in [0, 1)$ is the unique solution of equation (3.3). The minimum of the symbol is thus indeed given by (3.2).

Next, we establish some fine properties of energy thresholds. It follows from definitions (3.2)–(3.3) that $P \mapsto c(n, P)$ and $P \mapsto E_0^{(n)}(P)$ are radially symmetric for all $n \ge 1$. In addition, we find

$$c(n,0) = 0$$
, and $0 < c(n,P) < |P| \land 1$ for $P \neq 0$. (3.6)

Further properties are collected in the following statement. Item (iii) provides a simple criterion to compare the uncoupled eigenvalue $\frac{1}{2}P^2$ to energy thresholds, which proves in particular the last part of Lemma 1.1.

Lemma 3.2. Given a boson mass m > 0, let energy thresholds be defined in (3.2)-(3.3).

(i) For all $n \ge 1$, we have

$$c(n, P) \uparrow 1$$
 and $E_0^{(n)}(P) = |P| - \frac{1}{2} + o(1)$ as $|P| \uparrow \infty$.

(ii) For all P, we have

$$c(n,P) \downarrow 0$$
 and $E_0^{(n)}(P) = mn + o(1)$ as $n \uparrow \infty$.

(iii) For all $n \ge 1$, there exists a unique value $|P^{(n)}|$ such that the following equivalence holds:

$$\frac{1}{2}P^2 \ge E_0^{(n)}(P) \qquad \Longleftrightarrow \qquad |P| \ge |P^{(n)}|. \tag{3.7}$$

In addition, this value $|P^{(n)}|$ is increasing in n and we have $|P^{(1)}| = |P_{\star}| > 1$, where $|P_{\star}|$ is the critical value defined in Lemma 1.1.

(iv) Energy increments satisfy the following monotonicity properties,

for all
$$n: 0 < E_0^{(n+1)}(P) - E_0^{(n)}(P) \downarrow 0$$
 as $|P| \uparrow \infty$,
for all $P: 0 < E_0^{(n+1)}(P) - E_0^{(n)}(P) \uparrow m$ as $n \uparrow \infty$.

Proof. Items (i) and (ii) are direct consequences of definitions (3.2)-(3.3), so it remains to establish (iii) and (iv). We split the proof into two steps.

Step 1. Proof of (iii).

For all $n \ge 1$, starting from (3.2) and differentiating in |P|, a direct computation yields

$$\begin{split} & \frac{\partial}{\partial |P|} E_0^{(n)}(P) \\ & = \frac{|P| - c(n, P)}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}} + \left(c(n, P) - \frac{|P| - c(n, P)}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}}\right) \frac{\partial}{\partial |P|} c(n, P), \end{split}$$

and thus, by definition of c(n, P), cf. (3.3),

$$\frac{\partial}{\partial |P|} E_0^{(n)}(P) = \frac{|P| - c(n, P)}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}} = c(n, P).$$
(3.8)

In view of (3.6), this implies that the map $|P| \mapsto \frac{1}{2}|P|^2 - E_0^{(n)}(P)$ is increasing. Moreover, the asymptotic behavior in (i) ensures that this map is unbounded as $|P| \uparrow \infty$, and we see from (3.2) that it takes the value -mn < 0 at P = 0. This entails that for all $n \ge 1$ there exists a unique value $|P^{(n)}|$ such that

$$\frac{1}{2}|P^{(n)}|^2 - E_0^{(n)}(P^{(n)}) = 0, \qquad (3.9)$$

and the claimed equivalence (3.7) then follows by monotonicity.

It remains to check that the value $|P^{(n)}|$ is increasing in n. By (3.9), this follows provided that we show that energy thresholds are increasing in n for any fixed $P \neq 0$,

$$E_0^{(n+1)}(P) - E_0^{(n)}(P) > 0. (3.10)$$

For that purpose, we start by noting that definitions (3.2)-(3.3) yield

$$E_0^{(n)}(P) = F_P(c(n, P)), \qquad F_P(c) := \frac{1}{2}c^2 + \frac{|P|}{c} - 1.$$

As (3.6) ensures $c(n, P) < |P| \land 1$ for all n, as the function F_P is decreasing on $(-\infty, |P| \land 1)$, and as (ii) states that c(n, P) is decreasing in n, claim (3.10) follows. Alternatively, this result follows from identity (3.11) below.

Step 2. Proof of (iv).

/ \

We first investigate the behavior in |P| for fixed n. From (3.8) we deduce

$$\frac{\partial}{\partial |P|} \left(E_0^{(n+1)}(P) - E_0^{(n)}(P) \right) = c(n+1, P) - c(n, P),$$

which is negative in view of (ii). In addition, the asymptotic behavior in (i) ensures that energy increments tend to 0 as $|P| \uparrow \infty$.

We turn to the behavior in n for fixed P. As definitions (3.2)–(3.3) make sense for any $n \in (0, \infty)$, we may treat n as a continuous variable. Then starting from (3.2) and differentiating in n, we find

$$\frac{\partial}{\partial n} E_0^{(n)}(P) = \frac{m^2 n}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}} + \left(c(n, P) - \frac{|P| - c(n, P)}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}}\right) \frac{\partial}{\partial n} c(n, P),$$

and thus, by definition of c(n, P), cf. (3.3),

$$\frac{\partial}{\partial n} E_0^{(n)}(P) = \frac{m^2 n}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}}$$

This allows to write energy increments as

$$E_0^{(n+1)}(P) - E_0^{(n)}(P) = m \int_n^{n+1} \frac{mr}{\sqrt{m^2 r^2 + (|P| - c(r, P))^2}} \,\mathrm{d}r, \qquad (3.11)$$

0

which entails in particular for all n,

$$0 < E_0^{(n+1)}(P) - E_0^{(n)}(P) \le m, \qquad \lim_{n \uparrow \infty} \left(E_0^{(n+1)}(P) - E_0^{(n)}(P) \right) = m.$$

It remains to check that energy increments are increasing in n. For that purpose, we further compute

$$\frac{\partial}{\partial n} \frac{mn}{\sqrt{m^2 n^2 + (|P| - c(n, P))^2}} = \frac{m(|P| - c(n, P))}{(m^2 n^2 + (|P| - c(n, P))^2)^{\frac{3}{2}}} \Big(|P| - c(n, P) + n \frac{\partial}{\partial n} c(n, P) \Big). \quad (3.12)$$

Differentiating the definition (3.3) of c(n, P) with respect to n, we find after straightforward simplifications,

$$\frac{\partial}{\partial n}c(n,P) = -\frac{m^2n^2}{(m^2n^2 + (|P| - c(n,P))^2)^{\frac{3}{2}}}\frac{\partial}{\partial n}c(n,P) - \frac{m^2n(|P| - c(n,P))}{(m^2n^2 + (|P| - c(n,P))^2)^{\frac{3}{2}}},$$

and thus,

$$\frac{\partial}{\partial n}c(n,P) = -\frac{1}{n}(|P| - c(n,P)) \left(1 + \frac{(m^2n^2 + (|P| - c(n,P))^2)^{\frac{3}{2}}}{m^2n^2}\right)^{-1}.$$
(3.13)

This entails

$$\frac{\partial}{\partial n}c(n,P) > -\frac{1}{n}(|P| - c(n,P)),$$

so that (3.12) becomes

$$\frac{\partial}{\partial n}\frac{mn}{\sqrt{m^2n^2 + (|P| - c_n(P))^2}} > 0.$$

Combined with (3.11), this proves that the map $n \mapsto E_0^{(n+1)}(P) - E_0^{(n)}(P)$ is increasing, and the conclusion follows.

3.2. A first construction of conjugate operator. We turn to the construction of a conjugate operator for the uncoupled fiber Hamiltonian $H_0(P)$. A tentative conjugate is first constructed as a second-quantization operator by following (1.1), which we shall subsequently modify in Section 3.4 to improve on its regularity properties.

In case of massive bosons, as energy thresholds satisfy $E_0^{(n)}(P) \uparrow \infty$, cf. Lemma 3.2(iv), it suffices to construct a conjugate and prove a Mourre estimate separately on each energy interval

$$I_n(P) := [E_0^{(n)}(P), E_0^{(n+1)}(P)),$$

and on this interval we only need to compute commutators on state spaces with at most n bosons. In order to construct a conjugate operator in this setting, we follow (1.1) and first recall that on the *n*-boson state space the uncoupled fiber Hamiltonian has symbol

$$H_0^{(n)}(P;k_1,\ldots,k_n) = \frac{1}{2} \left(P - \sum_{j=1}^n k_j \right)^2 + \sum_{j=1}^n \omega(k_j), \qquad (3.14)$$

which attains a unique minimum at

$$k_1 = \ldots = k_n = k_\star(n, P),$$
 (3.15)

cf. (3.5). By convexity, a natural choice of conjugate on $\Gamma_s^{(n)}(\mathfrak{h})$ is then given by the generator of dilations around this minimum,

$$\sum_{j=1}^{n} \frac{i}{2} \Big(\big(k_j - k_\star(n, P)\big) \cdot \nabla_{k_j} + \nabla_{k_j} \cdot \big(k_j - k_\star(n, P)\big) \Big) \quad \text{on } \Gamma_s^{(n)}(\mathfrak{h}).$$
(3.16)

We could consider the operator that coincides with this choice on $\Gamma_s^{(n)}(\mathfrak{h})$ for all n, but, due to the dependence of $k_{\star}(n, P)$ on the number n of bosons, it would not lead to a secondquantization operator and would thus cause several issues such as the lack of regularity of the fiber interaction Hamiltonian $\Phi(\rho)$ — just as (2.5) in the quantum friction model. This is made particularly delicate as $k_{\star}(n, P)$ depends on n not explicitly — unlike (2.4). Instead, when focussing on the energy interval $I_n(P)$, we consider the following n-dependent second-quantization operator,

$$A_{P,n}^{\circ} := \mathrm{d}\Gamma(a_{P,n}^{\circ}), \qquad a_{P,n}^{\circ} := \frac{i}{2} \Big(\big(k - k_{\star}(n,P)\big) \cdot \nabla_k + \nabla_k \cdot \big(k - k_{\star}(n,P)\big) \Big). \tag{3.17}$$

It coincides with (3.16) on the *n*-boson state space, but for $\ell < n$ bosons it corresponds to the generator of dilations around $k_1 = \ldots = k_\ell = k_\star(n, P)$. Although this choice is inadequate for $\ell < n$ bosons, we shall see that it is compensated by the fact that in that case the energy interval $I_n(P)$ is further away from the minimum of the symbol. This will precisely allow to derive a Mourre estimate with respect to $A_{P,n}^{\circ}$ on $I_n(P)$.

We emphasize that the above definition (3.17) of the conjugate operator is quite different from previous choices in the literature [44, 37]. Indeed, we consider here boson momenta $k_j - k_{\star}(n, P)$ as measured in the reference frame minimizing the total kinetic energy (3.14), while in [44, 37] the starting point is instead to consider relative boson group velocities $\nabla_{k_j} H_0^{(n)}(P; k_1, \ldots, k_n)$. Our new choice appears particularly adapted to the problem and makes it possible to investigate for the first time the essential spectrum above the two-boson energy threshold in the weak-coupling regime.

Before turning to the proof of a Mourre estimate, we investigate properties of the abovedefined conjugate operator $A_{P,n}^{\circ}$. In particular, item (ii) states the C^2 -regularity of the uncoupled fiber Nelson Hamiltonian $H_0(P)$. We emphasize that this limited regularity is optimal, cf. [37, Section 2.2]: it comes from the fact that $A_{P,n}^{\circ}$ is a dilation around a point at a nonzero fixed distance from the origin, which entails that the commutator $[H_0(P), A_{P,n}^{\circ}]$ is only $NH_0(P)^{1/2}$ -bounded, hence $H_0(P)^{3/2}$ -bounded, but not $H_0(P)$ -bounded. In applications, this would prohibit to use the full power of Mourre's theory: results like Theorem A.8, for instance, are not available without stronger regularity. This issue will be resolved in Section 3.4 below by a suitable modification of $A_{P,n}^{\circ}$.

Lemma 3.3.

- (i) The conjugate operator $A_{P,n}^{\circ}$ is essentially self-adjoint on C^{f} and its closure generates a unitary group that commutes with the number operator and leaves the domain \mathcal{D} of fiber Hamiltonians (1.10) invariant.
- (ii) The fiber Hamiltonian $H_0(P)$ is of class $C^2(A_{P,n}^{\circ})$.
- (iii) Let the interaction kernel ρ belong to $H^{\nu}(\mathbb{R}^d)$ with $\langle k \rangle^{\nu} \nabla^{\nu} \rho \in L^2(\mathbb{R}^d)$ for some $\nu \geq 1$. Then, for all $0 \leq s \leq \nu$, the s-th iterated commutator $\operatorname{ad}_{iA_{P,n}^{\circ}}^{s}(\Phi(\rho))$ extends as an $N^{1/2}$ -bounded self-adjoint operator. \Diamond

Proof. We start with item (i). Clearly, $a_{P,n}^{\circ}$ is essentially self-adjoint on $C_c^{\infty}(\mathbb{R}^d)$, and the essential self-adjointness of $A_{P,n}^{\circ}$ on C^{f} follows. In addition, as the unitary group generated by $a_{P,n}^{\circ}$ takes the explicit form

$$(e^{ita_{P,n}^{\circ}}u)(k) = e^{-t\frac{d}{2}}u\Big(e^{-t}(k-k_{\star}(n,P))+k_{\star}(n,P)\Big), \quad u \in L^{2}(\mathbb{R}^{d})$$

the domain $\mathcal{D} = \mathcal{D}(H_0(P))$ is obviously invariant under $e^{itA_{P,n}^\circ} = \Gamma(e^{ita_{P,n}^\circ})$. Next, the proof of (ii) follows from a direct computation and is a particular case of [37, Proposition 2.5]. It remains to check (iii). As $A_{P,n}^\circ = d\Gamma(a_{P,n}^\circ)$ is a second-quantization operator, we find

$$[\Phi(\rho), iA_{P,n}^{\circ}] = -\Phi(ia_{P,n}^{\circ}\rho).$$

As $\Phi(ia_{P,n}^{\circ}\rho)$ is $N^{1/2}$ -bounded provided $a_{P,n}^{\circ}\rho \in L^2(\mathbb{R}^d)$, and repeating the same computation for iterated commutators, the conclusion follows.

3.3. Mourre estimate. We turn to the proof of a Mourre estimate for the uncoupled fiber Hamiltonian $H_0(P)$ with respect to the above-constructed conjugate operator $A_{P,n}^{\circ}$. It requires a quite delicate computation based on fine properties of the symbol (3.4) around its minimizer (3.5). Note that, surprisingly, our construction does not allow to treat the case of energy intervals I's with $1 < n < n_P$ in the notation below.

Lemma 3.4. Given a total momentum $|P| > |P_{\star}|$, define $n_P \ge 1$ such that

$$\frac{1}{2}P^2 \in \left[E_0^{(n_P)}(P), E_0^{(n_P+1)}(P)\right).$$
(3.18)

For all $\varepsilon > 0$ and all energy intervals $I \subset \left[E_0^{(n)}(P) + \varepsilon, E_0^{(n+1)}(P)\right)$ with n = 1 or $n \ge n_P$, the following Mourre estimate holds with respect to $A_{P,n}^{\circ}$ on I,

$$\mathbb{1}_{I}(H_{0}(P))[H_{0}(P), iA_{P,n}^{\circ}]\mathbb{1}_{I}(H_{0}(P)) \geq \varepsilon \bar{\Pi}_{\Omega}\mathbb{1}_{I}(H_{0}(P))\bar{\Pi}_{\Omega}.$$
(3.19)

In particular, the Mourre estimate is strict if I does not contain the eigenvalue $\frac{1}{2}P^2$. \diamond

Proof. Let $|P| > |P_{\star}|$ be fixed, define n_P via (3.18), and consider an energy interval

$$I \subset \left[E_0^{(n)}(P) + \varepsilon, E_0^{(n+1)}(P)\right)$$
(3.20)

for some $n \ge 1$ and $\varepsilon > 0$. We split the proof into four steps.

Step 1. Proof that the Mourre estimate (3.19) follows if we show for all $1 \le \ell \le n$,

$$\Pi_{\ell} \mathbb{1}_{I}(H_{0}(P))[H_{0}(P), iA_{P,n}^{\circ}] \mathbb{1}_{I}(H_{0}(P))\Pi_{\ell} \geq \varepsilon \Pi_{\ell} \mathbb{1}_{I}(H_{0}(P))\Pi_{\ell}.$$
(3.21)

where Π_{ℓ} is the orthogonal projection on the ℓ -boson state space $\Gamma_s^{(\ell)}(\mathfrak{h})$.

First recall that $H_0(P)$ and $A_{P,n}^{\circ}$ commute with the number operator, hence with each projection Π_{ℓ} . It is thus enough to prove (3.21) for each $\ell \geq 0$. Now, for $\ell = 0$, this lower bound is trivial as Ω is an eigenvector of $H_0(P)$. Next, as the symbol of $H_0(P)$ is bounded below by $E_0^{(\ell)}(P)$ on the ℓ -boson state space, and as we have $E_0^{(\ell)}(P) \geq E_0^{(n+1)}(P)$ for $\ell > n$ in view of Lemma 3.2(iv), the choice (3.20) of the energy interval I entails

$$\mathbb{1}_{I}(H_{0}(P))\Pi_{\ell} = 0 \quad \text{for all } \ell > n, \qquad (3.22)$$

and the claim follows.

Step 2. Proof that for all $1 \leq \ell \leq n$,

$$\Pi_{\ell}[H_0(P), iA_{P,n}^{\circ}]\Pi_{\ell} \ge \Pi_{\ell}H_0(P)\Pi_{\ell} - H_0^{(\ell)}(P; k_{\star}^{(n)}, \dots, k_{\star}^{(n)})\Pi_{\ell}, \qquad (3.23)$$

where henceforth we set for notational simplicity

$$k_{\star}^{(n)} := k_{\star}(n, P) = \frac{1}{n} (|P| - c(n, P)) \frac{P}{|P|}, \qquad (3.24)$$

cf. (3.5), thus omitting the dependence on P in the notation.

For that purpose, we start by noting that the symbol (3.4) of $H_0(P)$ on the ℓ -boson state space can be decomposed as

$$H_0^{(\ell)}(P;k_1,\ldots,k_\ell) = \frac{1}{2} \Big(\sum_{j=1}^\ell k_j - nk_\star^{(n)} \Big)^2 + \sum_{j=1}^\ell \Big(\omega(k_j) - \omega(k_\star^{(n)}) - c(n,P) \frac{P}{|P|} \cdot (k_j - k_\star^{(n)}) \Big) \\ + H_0^{(\ell)} \Big(P;k_\star^{(n)},\ldots,k_\star^{(n)} \Big) - \frac{1}{2} (n-\ell)^2 |k_\star^{(n)}|^2. \quad (3.25)$$

By definition (3.17) of the conjugate operator, writing

$$iA_{P,n}^{\circ} = -\mathrm{d}\Gamma\Big((k-k_{\star}^{(n)})\cdot\nabla_k + \frac{d}{2}\Big),$$

a direct computation then yields on $\Gamma_s^{(\ell)}(\mathfrak{h})$,

$$\begin{aligned} [H_0(P), iA_{P,n}^{\circ}]|_{\Gamma_s^{(\ell)}(\mathfrak{h})} &= \left[\sum_{j=1}^{\ell} (k_j - k_\star^{(n)}) \cdot \nabla_{k_j} , \ H_0^{(\ell)}(P; k_1, \dots, k_\ell)\right] \\ &= \left(\sum_{j=1}^{\ell} (k_j - k_\star^{(n)})\right) \cdot \left(\sum_{j=1}^{\ell} k_j - nk_\star^{(n)}\right) + \sum_{j=1}^{\ell} (k_j - k_\star^{(n)}) \cdot \left(\nabla\omega(k_j) - c(n, P)\frac{P}{|P|}\right). \end{aligned}$$

This identity can be further reorganized as follows,

$$[H_0(P), iA_{P,n}^{\circ}]|_{\Gamma_s^{(\ell)}(\mathfrak{h})} = \frac{n-\ell}{2n} \Big(\sum_{j=1}^{\ell} k_j \Big)^2 + \frac{n+\ell}{2n} \Big(\sum_{j=1}^{\ell} k_j - nk_\star^{(n)} \Big)^2 - \frac{1}{2}n(n-\ell)|k_\star^{(n)}|^2 + \sum_{j=1}^{\ell} \Big(\omega(k_j) - \omega(k_\star^{(n)}) - c(n, P)\frac{P}{|P|} \cdot (k_j - k_\star^{(n)}) \Big) + \sum_{j=1}^{\ell} \Big(\omega(k_\star^{(n)}) - \frac{k_j \cdot k_\star^{(n)} + m^2}{\omega(k_j)} \Big).$$

Recognizing the symbol $H_0^{(\ell)}(P; k_1, \ldots, k_\ell)$ in the right-hand side in form of (3.25), and noting that

$$\omega(k_{\star}^{(n)}) \ge \frac{k \cdot k_{\star}^{(n)} + m^2}{\omega(k)} \quad \text{for all } k,$$

we deduce

$$\begin{aligned} \left[H_0(P), iA_{P,n}^{\circ}\right]|_{\Gamma_s^{(\ell)}(\mathfrak{h})} &\geq H_0^{(\ell)}(P; k_1, \dots, k_\ell) - H_0^{(\ell)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) \\ &+ \frac{n-\ell}{2n} \Big(\sum_{j=1}^\ell k_j\Big)^2 + \frac{\ell}{2n} \Big(\sum_{j=1}^\ell k_j - nk_\star^{(n)}\Big)^2 - \frac{1}{2}\ell(n-\ell)|k_\star^{(n)}|^2. \end{aligned}$$

Finally, noting that

$$\frac{n-\ell}{2n} \left(\sum_{j=1}^{\ell} k_j\right)^2 + \frac{\ell}{2n} \left(\sum_{j=1}^{\ell} k_j - nk_{\star}^{(n)}\right)^2$$
$$= \frac{1}{2} \left(\sum_{j=1}^{\ell} k_j\right)^2 - \ell k_{\star}^{(n)} \cdot \left(\sum_{j=1}^{\ell} k_j\right) + \frac{1}{2} \ell n |k_{\star}^{(n)}|^2$$

$$= \frac{1}{2} \left(\sum_{j=1}^{\ell} k_j - \ell k_{\star}^{(n)} \right)^2 + \frac{1}{2} \ell(n-\ell) |k_{\star}^{(n)}|^2$$

$$\geq \frac{1}{2} \ell(n-\ell) |k_{\star}^{(n)}|^2,$$

we conclude

$$[H_0(P), iA_{P,n}^{\circ}]|_{\Gamma_s^{(\ell)}(\mathfrak{h})} \ge H_0^{(\ell)}(P; k_1, \dots, k_\ell) - H_0^{(\ell)}(P; k_{\star}^{(n)}, \dots, k_{\star}^{(n)}),$$

that is, (3.23).

Step 3. Proof that, given an energy interval I as in (3.20), we have for all $1 \le \ell \le n$,

$$\Pi_{\ell} \mathbb{1}_{I}(H_{0}(P))[H_{0}(P), iA_{P,n}^{\circ}]\mathbb{1}_{I}(H_{0}(P))\Pi_{\ell}$$

$$\geq \varepsilon \Pi_{\ell} + \left(1 - \frac{\ell}{n}\right) \left(\frac{1}{2n} (|P| - c(n, P))^2 - \alpha(n, P)\right) \Pi_{\ell}, \quad (3.26)$$

where $\alpha(n, P)$ stands for the positive distance between the eigenvalue $\frac{1}{2}P^2$ and the energy threshold below I,

$$\alpha(n,P) := \frac{1}{2}P^2 - E_0^{(n)}(P) = \frac{1}{2}P^2 - H_0^{(n)}(P;k_\star^{(n)},\dots,k_\star^{(n)}).$$
(3.27)

Starting from (3.23), and recalling that the choice (3.20) of I yields

$$\inf I \ge \varepsilon + E_0^{(n)}(P) = \varepsilon + H_0^{(n)}(P; k_{\star}^{(n)}, \dots, k_{\star}^{(n)}),$$

we are led to

$$\Pi_{\ell} \mathbb{1}_{I}(H_{0}(P))[H_{0}(P), iA_{P,n}^{\circ}] \mathbb{1}_{I}(H_{0}(P))\Pi_{\ell} \\ \geq \left(\varepsilon + H_{0}^{(n)}(P; k_{\star}^{(n)}, \dots, k_{\star}^{(n)}) - H_{0}^{(\ell)}(P; k_{\star}^{(n)}, \dots, k_{\star}^{(n)})\right)\Pi_{\ell}.$$

To prove (3.26), it thus remains to check for all $\ell \leq n$,

$$H_0^{(n)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) - H_0^{(\ell)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) \geq \left(1 - \frac{\ell}{n}\right) \left(\frac{1}{2n} (|P| - c(n, P))^2 - \alpha(n, P)\right). \quad (3.28)$$

For that purpose, we decompose

$$\begin{split} H_0^{(\ell)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) &= \frac{1}{2} (P - \ell k_\star^{(n)})^2 + \ell \omega(k_\star^{(n)}) \\ &= \frac{1}{2} P^2 + \frac{1}{2} \ell^2 |k_\star^{(n)}|^2 - \ell P \cdot k_\star^{(n)} + \ell \omega(k_\star^{(n)}) \\ &= \frac{\ell}{n} \left(\frac{1}{2} P^2 + \frac{1}{2} n^2 |k_\star^{(n)}|^2 - n P \cdot k_\star^{(n)} + n \omega(k_\star^{(n)}) \right) + \frac{1}{2} \left(1 - \frac{\ell}{n} \right) P^2 - \frac{1}{2} \ell (n - \ell) |k_\star^{(n)}|^2 \\ &= \frac{\ell}{n} H_0^{(n)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) + \frac{1}{2} \left(1 - \frac{\ell}{n} \right) P^2 - \frac{1}{2} \ell n \left(1 - \frac{\ell}{n} \right) |k_\star^{(n)}|^2. \end{split}$$

In terms of (3.27), this yields

$$\begin{aligned} H_0^{(n)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) &- H_0^{(\ell)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) \\ &= (1 - \frac{\ell}{n}) H_0^{(n)}(P; k_\star^{(n)}, \dots, k_\star^{(n)}) - \frac{1}{2} (1 - \frac{\ell}{n}) P^2 + \frac{1}{2} \ell n (1 - \frac{\ell}{n}) |k_\star^{(n)}|^2 \\ &= (1 - \frac{\ell}{n}) \Big(\frac{1}{2} \ell n |k_\star^{(n)}|^2 - \alpha(n, P) \Big). \end{aligned}$$

Now recalling $|k_{\star}^{(n)}| = \frac{1}{n}(|P| - c(n, P))$, cf. (3.24), the claim (3.28) follows.

Step 4. Conclusion.

As n_P is defined via (3.18), the definition (3.27) of α yields $\alpha(n, P) \leq 0$ for $n > n_P$. The right-hand side in (3.26) is thus bounded below by $\varepsilon \Pi_{\ell}$ if $\ell = n$, or if $\ell < n$ and $n > n_P$. It remains to prove the corresponding result in the case $\ell < n = n_P$. In other words, it remains to prove the following implication, for all n, P,

$$\frac{1}{2}|P|^2 \in \left[E_0^{(n)}(P), E_0^{(n+1)}(P)\right) \implies \frac{1}{2n}(|P| - c(n, P))^2 - \alpha(n, P) \ge 0.$$
(3.29)

For that purpose, we start by noting that the definition (3.3) of c(n, P) yields

$$|P| - c(n, P) = mn \frac{c(n, P)}{\sqrt{1 - c(n, P)^2}},$$
(3.30)

which allows to rewrite (3.2) in particular as

$$E_0^{(n)}(P) = \frac{1}{2}c(n,P)^2 + \frac{mn}{\sqrt{1 - c(n,P)^2}}$$

and thus

$$\alpha(n,P) = \frac{1}{2}P^2 - \frac{1}{2}c(n,P)^2 - \frac{mn}{\sqrt{1 - c(n,P)^2}}.$$
(3.31)

Further inserting (3.30) in this last identity to eliminate |P|, we get

$$\begin{aligned} \alpha(n,P) &= \frac{1}{2}c(n,P)^2 \left(1 + \frac{mn}{\sqrt{1 - c(n,P)^2}}\right)^2 - \frac{1}{2}c(n,P)^2 - \frac{mn}{\sqrt{1 - c(n,P)^2}} \\ &= \frac{1}{2}m^2n^2\frac{c(n,P)^2}{1 - c(n,P)^2} - mn\sqrt{1 - c(n,P)^2}. \end{aligned}$$
(3.32)

Combining this with (3.30) again to reformulate the quantity of interest in (3.29), we find

$$\frac{1}{2n}(|P| - c(n,P))^2 - \alpha(n,P) = -\frac{1}{2}m^2n(n-1)\frac{c(n,P)^2}{1 - c(n,P)^2} + mn\sqrt{1 - c(n,P)^2},$$

which entails that the implication (3.29) is actually equivalent to

$$\frac{1}{2}P^2 \in \left[E_0^{(n)}(P), E_0^{(n+1)}(P)\right) \implies \frac{(1-c(n,P)^2)^{\frac{3}{2}}}{c(n,P)^2} \ge \frac{1}{2}m(n-1).$$

As the map $|P| \mapsto c(n, P)$ is increasing, cf. Lemma 3.2(i), it suffices to prove this implication for P such that $\frac{1}{2}P^2 = E_0^{(n+1)}(P)$, that is, for $|P| = |P^{(n+1)}|$, cf. Lemma 3.2(iii). We are thus reduced to proving for all n,

$$\frac{(1-c(n,P^{(n+1)})^2)^{\frac{3}{2}}}{c(n,P^{(n+1)})^2} \ge \frac{1}{2}m(n-1).$$
(3.33)

As by definition $\alpha(n+1, P^{(n+1)}) = \frac{1}{2}|P^{(n+1)}|^2 - E_0^{(n+1)}(P^{(n+1)}) = 0$, identity (3.32) entails

$$\frac{1}{2}m(n+1) = \frac{\left(1 - c(n+1, P^{(n+1)})^2\right)^{\frac{3}{2}}}{c(n+1, P^{(n+1)})^2}.$$
(3.34)

The claim (3.33) can thus be reformulated as

$$\frac{\left(1 - c(n+1, P^{(n+1)})^2\right)^{\frac{3}{2}}}{c(n+1, P^{(n+1)})^2} - \frac{\left(1 - c(n, P^{(n+1)})^2\right)^{\frac{3}{2}}}{c(n, P^{(n+1)})^2} \le m,$$
(3.35)

and it directly follows in this form from Lemma 3.5 below.

The conclusion of the above proof of the Mourre estimate relies on the following key computation, which we state as a separate lemma for convenience.

Lemma 3.5. For all $n \ge 1$, the function defined by

$$f_{n+1}(r) := \frac{\left(1 - c(r, P^{(n+1)})^2\right)^{\frac{3}{2}}}{c(r, P^{(n+1)})^2}$$

satisfies $f'_{n+1}(r) \le m$ for all $0 < r \le n+1$.

Proof. We split the proof into two steps.

Step 1. Proof that it suffices to show $f(x) = \frac{1}{2} \int \frac{1}{2}$

$$f_{n+1}(r) \le \frac{1}{2}rm$$
 for all $0 < r \le n+1$. (3.36)

For notational simplicity, we set $c(r) := c(r, P^{(n+1)})$. The derivative of f_{n+1} takes the form

$$f'_{n+1}(r) = -\frac{1}{c(r)^4} \left(3c(r)^3 c'(r) \sqrt{1 - c(r)^2} + 2c(r)c'(r)(1 - c(r)^2)^{\frac{3}{2}} \right)$$

$$= -c'(r) \frac{c(r)^2 + 2}{c(r)^3} \sqrt{1 - c(r)^2}.$$
 (3.37)

Recall that the derivative of c was computed in (3.13),

$$c'(r) = -\frac{1}{r}(|P^{(n+1)}| - c(r))\left(1 + \frac{(m^2r^2 + (|P^{(n+1)}| - c(r))^2)^{\frac{3}{2}}}{m^2r^2}\right)^{-1}$$

and thus, using (3.30) in form of

$$|P^{(n+1)}| - c(r) = mr \frac{c(r)}{\sqrt{1 - c(r)^2}}$$

we find

$$c'(r) = -\frac{mc(r)}{\sqrt{1-c(r)^2}} \left(1 + \frac{mr}{(1-c(r)^2)^{\frac{3}{2}}}\right)^{-1}$$
$$= -\frac{mc(r)(1-c(r)^2)}{mr + (1-c(r)^2)^{\frac{3}{2}}}.$$

Inserting this into (3.37), we get

$$f'_{n+1}(r) = \frac{m(1-c(r)^2)^{\frac{3}{2}}}{mr+(1-c(r)^2)^{\frac{3}{2}}} \frac{c(r)^2+2}{c(r)^2},$$

and we deduce the following equivalence: for all r,

$$f'_{n+1}(r) \le m \iff \frac{(1-c(r)^2)^{\frac{3}{2}}}{mr+(1-c(r)^2)^{\frac{3}{2}}} \frac{c(r)^2+2}{c(r)^2} \le 1$$
$$\iff 2(1-c(r)^2)^{\frac{3}{2}} \le mrc(r)^2$$
$$\iff f_{n+1}(r) \le \frac{1}{2}mr,$$

as claimed.

Step 2. Conclusion.

Let $0 < r \le n+1$ be fixed. As the map $|P| \mapsto c(r, P)$ is increasing and as $|P^{(n+1)}| \ge |P^{(r)}|$

 \diamond

in view of Lemma 3.2(iii) (where we can extend the definition of $|P^{(r)}|$ to all real r > 0), we find

$$f_{n+1}(r) = \frac{(1 - c(r, P^{(n+1)})^2)^{\frac{3}{2}}}{c(r, P^{(n+1)})^2} \le \frac{(1 - c(r, P^{(r)})^2)^{\frac{3}{2}}}{c(r, P^{(r)})^2}.$$
(3.38)

Noting that the same argument as for (3.34) yields

$$\frac{(1 - c(r, P^{(r)})^2)^{\frac{3}{2}}}{c(r, P^{(r)})^2} = \frac{1}{2}mr$$

we deduce $f_{n+1}(r) \leq \frac{1}{2}mr$, and the conclusion follows from Step 1.

3.4. Modification procedure and improved regularity. This section is devoted to the modification of the tentative conjugate operator $A_{P,n}^{\circ}$ to improve on the associated regularity properties in Lemma 3.3(ii), in view of the proof of Theorem 1.2. More precisely, we shall only modify $A_{P,n}^{\circ}$ on ℓ -boson state spaces for all $\ell > n$, while keeping it unchanged elsewhere. Indeed, by (3.22), we recall that a Mourre estimate on the energy interval $I_n(P)$ only needs to be checked on ℓ -boson state spaces for all $1 \leq \ell \leq n$, hence our modification will not impact the validity of the Mourre estimate proven for $A_{P,n}^{\circ}$ in Lemma 3.4. We emphasize that our modification procedure is quite general and may be of independent interest for other massive QFT models.

3.4.1. Motivation for modification procedure. We start by further examining the abovedefined tentative conjugate operator $A_{P,n}^{\circ}$, decomposing it as

$$A_{P,n}^{\circ} = D_{\circ} - \mathrm{d}\Gamma\big(ik_{\star}(n,P)\cdot\nabla_k\big),\tag{3.39}$$

where D_{\circ} stands for the generator of dilations,

$$D_{\circ} := \mathrm{d}\Gamma(d_{\circ}), \qquad d_{\circ} := \frac{i}{2}(k \cdot \nabla_k + \nabla_k \cdot k).$$

The lack of regularity of $H_0(P)$ with respect to $A_{P,n}^{\circ}$ precisely originates from the second term $d\Gamma(ik_{\star}(n, P) \cdot \nabla_k)$ in (3.39), as indeed its commutator with $(P - d\Gamma(k))^2$ is not $H_0(P)$ -bounded. To cure this issue, we might naïvely want to rather consider the truncated operator

$$A'_{P,n} := D_{\circ} - \prod_{\leq n} \mathrm{d}\Gamma\big(ik_{\star}(n,P) \cdot \nabla_k\big) \prod_{\leq n}, \tag{3.40}$$

in terms of the orthogonal projection $\Pi_{\leq n}$ onto $\bigoplus_{\ell=0}^{n} \Gamma_{s}^{(\ell)}(\mathfrak{h})$. By definition, $H_{0}(P)$ is now of class $C^{\infty}(A'_{P,n})$. In addition, as this operator coincides with $A_{P,n}^{\circ}$ on the range of $\Pi_{\leq n}$, we deduce that $H_{0}(P)$ satisfies the same Mourre estimate with respect to $A'_{P,n}$ as in Lemma 3.4.

This is however not the end of the story: the brutal truncation in (3.40) happens to behave badly with respect to the fiber Hamiltonian $\Phi(\rho)$, in link with the fact that $A'_{P,n}$ is no longer a second-quantization operator. The truncation thus needs to be suitably complemented on ℓ -boson state spaces for $\ell > n$, although not by means of second quantization. In the spirit of our constructions in Section 2.1 for the quantum friction model, instead of considering the second quantization $d\Gamma(ik_*(n, P) \cdot \nabla_k)$ in (3.39), which amounts to taking sums of coordinates $\{ik_*(n, P) \cdot \nabla_{k_j}\}_j$, and instead of taking a brutal truncation as in (3.40) on ℓ -boson state spaces with $\ell > n$, we shall consider partial sums of the *n* largest signed values of the coordinates.

3.4.2. Partial sums of largest signed values and regularization. Instead of momentum representation on \mathfrak{h} , we shall use position representation: we denote by $y := i\nabla_{\xi}$ the position coordinate, and we set $z := \frac{P}{|P|} \cdot y$ for the coordinate in the *P*-direction. For all $1 \leq j \leq \ell$, we define the function $m_{j,\ell} : \mathbb{R}^{\ell} \to \mathbb{R}$ as the *j*th largest signed value of the entries: for all $z_1, \ldots, z_{\ell} \in \mathbb{R}$, we set

$$m_{j,\ell}(z_1,\ldots,z_\ell) := z_{i_0}$$

where the index i_0 is chosen such that $|z_{i_0}|$ is the *j*th largest value among $|z_1|, \ldots, |z_\ell|$. This is obviously well-defined on \mathbb{R}^{ℓ} up to a null set. Note that for j = 1 the function $m_{1,\ell}$ coincides with the signed maximum m_{ℓ} defined in (2.8). For all $1 \leq j < \ell$, we then define the function $s_{j,\ell} : \mathbb{R}^{\ell} \to \mathbb{R}$ as the sum of the *j* largest signed entries,

$$s_{j,\ell} := m_{1,\ell} + \ldots + m_{j,\ell},$$

and for $j \ge \ell$ we simply define

$$s_{i,\ell}(z_1,\ldots,z_\ell) := z_1 + \ldots + z_\ell$$

As in Lemma 2.1, we note that $s_{j,\ell}$ is not continuous for $j < \ell$ and thus needs to be regularized, which we shall carefully perform in the spirit of (2.12). For that purpose, we start by defining for all $1 \le j \le \ell$ the functions $\max_{j,\ell} : \mathbb{R}^\ell \to \mathbb{R}$ and $\min_{j,\ell} : \mathbb{R}^\ell \to \mathbb{R}$ as the *j*th largest and the *j*th smallest entries, respectively: more precisely, these functions are defined to be symmetric upon permutation of their entries, and to satisfy $\max_{j,\ell}(z_1,\ldots,z_\ell) = z_{\ell-j+1}$ and $\min_{j,\ell}(z_1,\ldots,z_\ell) = z_j$ if $z_1 \le \ldots \le z_\ell$. These functions are both obviously well-defined and continuous, and we have the relations

$$|m_{j,\ell}(z_1,...,z_{\ell})| = \max_{j,\ell}(|z_1|,...,|z_{\ell}|),$$

$$\max_{1,\ell}(z_1,...,z_{\ell}) = \max_{1 \le l \le \ell} z_l = \min_{\ell,\ell}(z_1,...,z_{\ell}),$$

$$\min_{1,\ell}(z_1,...,z_{\ell}) = \min_{1 \le l \le \ell} z_l = \max_{\ell,\ell}(z_1,...,z_{\ell}).$$

In these terms, for all $1 \leq j < \ell$, we note that the definition of $s_{j,\ell}$ can be reformulated as

$$s_{j,\ell} = \sum_{l=1}^{j} \left(\frac{1}{2} \left(\max_{j+1-l,\ell} + \min_{l,\ell} \right) + \frac{1}{2} \left(\max_{j+1-l,\ell} - \min_{l,\ell} \right) \operatorname{sgn} \left(\max_{j+1-l,\ell} + \min_{l,\ell} \right) \right),$$

where now only the sign functions need to be regularized. Given $\delta > 0$, we choose a smooth odd function $\chi_{\delta} : \mathbb{R} \to [-1, 1]$ as in (2.11), and we define for all $1 \leq j < \ell$,

$$\widetilde{s}_{j,\ell;\delta} = \sum_{l=1}^{j} \left(\frac{1}{2} \left(\max_{j+1-l,\ell} + \min_{l,\ell} \right) + \frac{1}{2} \left(\max_{j+1-l,\ell} - \min_{l,\ell} \right) \chi_{\delta} \left(\frac{\max_{j+1-l,\ell} + \min_{l,\ell}}{1 + \max_{j+1-l,\ell} - \min_{l,\ell}} \right) \right),$$

which is obviously globally well-defined and continuous. For $j \ge \ell$, no regularization is needed and we simply set

$$\widetilde{s}_{j,\ell;\delta}(z_1,\ldots,z_\ell) := s_{j,\ell}(z_1,\ldots,z_\ell) = z_1 + \ldots + z_\ell$$

In view of properties of χ_{δ} , a direct computation yields

$$1 \leq \sum_{l=1}^{\ell} \partial_l \widetilde{s}_{j,\ell;\delta} \leq (2+\delta)(j \wedge \ell), \qquad (3.41)$$

and in addition, for all $r \geq 1$,

$$\left| \left(\sum_{l=1}^{\ell} \partial_l \right)^r \widetilde{s}_{j,\ell;\delta} \right| \lesssim_{\chi_{\delta},r} j \wedge \ell, \qquad \left| \left(\sum_{l=1}^{\ell} z_l \partial_l \right)^r \widetilde{s}_{j,\ell;\delta} - \widetilde{s}_{j,\ell;\delta} \right| \lesssim_{\chi_{\delta},r} j \wedge \ell.$$
(3.42)

In particular, note that $\tilde{s}_{j,\ell;\delta}$ is smooth in the direction $(1,\ldots,1)$. Next, we state the following generalization of Lemma 2.2 for $\tilde{m}_{\ell;\delta} = \tilde{s}_{1,\ell;\delta}$, which will be key to estimate commutators with field operators. The proof is a direct adaptation of that of Lemma 2.2 and we skip the detail.

Lemma 3.6. For all $j, \ell \geq 1$, there holds for all $z, z_1, \ldots, z_\ell \in \mathbb{R}$,

$$\left|\widetilde{s}_{j,\ell+1;\delta}(z,z_1,\ldots,z_\ell) - \widetilde{s}_{j,\ell;\delta}(z_1,\ldots,z_\ell)\right| \leq 2|z|+1.$$

3.4.3. Back to conjugate operator. With the above construction at hand, we turn to the suitable replacement for the second term in the tentative choice (3.39) of the conjugate operator. For all n, ℓ , we define the operator $S_{P,n,\ell;\delta}$ on the ℓ -boson state space as the multiplication with the function

$$(y_1,\ldots,y_\ell) \mapsto |k_\star(n,P)| \,\widetilde{s}_{n,\ell;\delta} \Big(\frac{P}{|P|} \cdot y_1,\ldots,\frac{P}{|P|} \cdot y_\ell \Big), \tag{3.43}$$

using position representation $y = i\nabla_k$ on \mathfrak{h} , and we define

$$S_{P,n;\delta} := \bigoplus_{\ell=1}^{\infty} S_{P,n,\ell;\delta} \quad \text{on } \mathcal{H}^{\mathrm{f}} := \bigoplus_{\ell=0}^{\infty} \Gamma_s^{(\ell)}(\mathfrak{h}).$$
(3.44)

Coming back to (3.39), we then define the following modified conjugate operator,

$$A_{P,n;\delta} := D_{\circ} - S_{P,n;\delta} \quad \text{on } \mathcal{H}^{\mathrm{f}}, \qquad (3.45)$$

where we recall that D_{\circ} stands for the generator of dilations,

$$D_{\circ} = \mathrm{d}\Gamma(d_{\circ}), \qquad d_{\circ} = \frac{i}{2}(k \cdot \nabla_k + \nabla_k \cdot k).$$

By definition, the operator $A_{P,n;\delta}$ commutes with the number operator. Given its action on ℓ -boson state space, it is clearly essentially self-adjoint on \mathcal{C}^{f} , and we state that it generates an explicit unitary group that preserves the domain of fiber Hamiltonians. The proof is analogous to that of Lemma 2.3 and is skipped for brevity.

Lemma 3.7. The operator $A_{P,n;\delta}$ is essentially self-adjoint on C^{f} and its closure generates a unitary group $\{e^{itA_{P,n;\delta}}\}_{t\in\mathbb{R}}$ on \mathcal{H}^{f} , which commutes with the number operator and has the following explicit action: for all $\ell \geq 1$ and $u_{\ell} \in \Gamma_s^{(\ell)}(\mathfrak{h})$,

$$(e^{itA_{P,n;\delta}}u_{\ell})(y_1,\ldots,y_{\ell}) = \exp\left(-i|k_{\star}(n,P)|\int_0^t \widetilde{s}_{n,\ell;\delta}\left(\frac{P}{|P|}\cdot e^s y_1,\ldots,\frac{P}{|P|}\cdot e^s y_{\ell}\right)ds\right) \\ \times e^{t\ell\frac{d}{2}}u_{\ell}(e^t y_1,\ldots,e^t y_{\ell}),$$

where we use position representation on \mathfrak{h} . In particular, the domain \mathcal{D} of fiber Hamiltonians (1.10) is invariant under this group action.

3.4.4. Improved regularity. As by definition $A_{P,n;\delta}$ coincides with $A_{P,n}^{\circ}$ on ℓ -boson state spaces for all $\ell \leq n$, it follows from (3.22) that the Mourre estimate of Lemma 3.4 holds in the exact same form with respect to $A_{P,n;\delta}$, thus proving Theorem 1.2(ii). Next, we show that the fiber Hamiltonian $H_0(P)$ is now of class $C^{\infty}(A_{P,n;\delta})$, which improves on the limited C^2 -regularity available with respect to $A_{P,n}^{\circ}$, cf. Lemma 3.3(ii). Combined with Lemma 3.7, this proves Theorem 1.5(i), further noting that the $C^{\infty}(A_{P,n;\delta})$ -regularity property indeed follows by applying the sufficient criterion in Lemma A.3.

Lemma 3.8. For all $s \ge 1$, the s-th iterated commutator $\operatorname{ad}_{iA_{P,n,\delta}}^{s}(H_0(P))$ extends as an $H_0(P)$ -bounded self-adjoint operator.

Proof. By definition (3.45) of $A_{P,n;\delta}$, the first commutator can be split as

$$[H_0(P), iA_{P,n;\delta}] = [H_0(P), iD_\circ] - [H_0(P), iS_{P,n;\delta}] = -d\Gamma(k) \cdot (P - d\Gamma(k)) + d\Gamma(k \cdot \nabla\omega(k)) - [H_0(P), iS_{P,n;\delta}].$$

Using that $|k \cdot \nabla \omega(k)| = \omega(k) - m^2 \omega(k)^{-1} \leq \omega(k)$, the first two right-hand side terms are obviously $H_0(P)$ -bounded operators: we get for all $u, v \in C^{\mathrm{f}}$,

 $|\langle u, [H_0(P), iA_{P,n;\delta}]v\rangle| \lesssim |P|^2 ||u|| ||v|| + ||u|| ||H_0(P)v|| + |\langle u, [H_0(P), iS_{P,n;\delta}]v\rangle|,$ (3.46) and it remains to estimate the term. For that purpose, using position representation $y = i\nabla_{\xi}$ on \mathfrak{h} , we write

$$[H_0(P), iS_{P,n;\delta}] = \frac{1}{2} [(P + d\Gamma(i\nabla_y))^2, iS_{P,n;\delta}] + [d\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}]$$

$$= -\frac{1}{2} d\Gamma(\nabla_y) \cdot [d\Gamma(\nabla_y), iS_{P,n;\delta}] - \frac{1}{2} [d\Gamma(\nabla_y), iS_{P,n;\delta}] \cdot d\Gamma(\nabla_y)$$

$$-P \cdot [d\Gamma(\nabla_y), S_{P,n;\delta}] + [d\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}],$$

or alternatively,

$$[H_0(P), iS_{P,n;\delta}] = -[\mathrm{d}\Gamma(\nabla_y), S_{P,n;\delta}] \cdot (P + \mathrm{d}\Gamma(i\nabla_y)) - \frac{1}{2} \big[\mathrm{d}\Gamma(\nabla_y) \cdot, [\mathrm{d}\Gamma(\nabla_y), iS_{P,n;\delta}] \big] + [\mathrm{d}\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}].$$

By definition (3.43)–(3.44) of $S_{P,n;\delta}$, recalling (3.5) and using (3.42), the commutators $[d\Gamma(\nabla_y), iS_{P,n;\delta}]$ and $[d\Gamma(\nabla_y), [d\Gamma(\nabla_y), iS_{P,n;\delta}]]$ are bounded by O(|P|). We deduce for all $u, v \in C^{f}$,

$$\begin{aligned} |\langle u, [H_0(P), iS_{P,n;\delta}]v\rangle| &\leq |P| ||u|| ||v|| + |P| ||u|| ||H_0(P)^{\frac{1}{2}}v|| \\ &+ |\langle u, [d\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}]v\rangle|, \quad (3.47) \end{aligned}$$

and it remains to estimate the last commutator $[d\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}]$, that is, on the ℓ -boson state space,

$$\left[\mathrm{d}\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}\right]\Big|_{\Gamma_s^{(\ell)}(\mathfrak{h})} = \sum_{j=1}^{\ell} \left[\omega(i\nabla_{y_j}), iS_{P,n,\ell;\delta}\right].$$
(3.48)

We split this task into three steps: we first show that the commutator $[|\nabla_{y_j}|, iS_{P,n,\ell;\delta}]$ is nicely bounded and then we appeal to the calculus of almost-analytic extensions to reduce the analysis of the difference $\omega(i\nabla_{y_j}) - |\nabla_{y_j}|$ to that of the resolvent $(z - |\nabla_{y_j}|)^{-1}$.

Step 1. Preliminary commutator estimates: for all $1 \leq j \leq \ell$, we have

$$\|[|\nabla_{y_j}|, iS_{P,n,\ell;\delta}]\| \lesssim \frac{1}{n}|P|, \qquad (3.49)$$

and in addition, for all $u_{\ell}, v_{\ell} \in \mathcal{C}^{\mathrm{f}} \cap \Gamma_s^{(\ell)}(\mathfrak{h})$ and $z \in \mathbb{C} \setminus \mathbb{R}$,

$$|\langle u_{\ell}, [(z - |\nabla_{y_j}|)^{-1}, iS_{P,n,\ell;\delta}]v_{\ell}\rangle| \lesssim \frac{1}{n} |P||\Im z|^{-2} ||u_{\ell}|| ||v_{\ell}||.$$
(3.50)

We start with the proof of (3.49). By symmetry, we focus on j = 1. Recalling the definition of $S_{P,n,\ell;\delta}$, cf. (3.43), and using the integral representation for $|\nabla_{y_1}| = (-\Delta_{y_1})^{1/2}$, we find for all $v_{\ell} \in \mathcal{C}^{\mathsf{f}} \cap \Gamma_s^{(\ell)}(\mathfrak{h})$,

$$\begin{aligned} [|\nabla_{y_1}|, iS_{P,n,\ell;\delta}] v_{\ell}(y_1, \dots, y_{\ell}) &= C|k_{\star}(n, P)| \\ \times \int_{\mathbb{R}^d} \frac{1}{|y_1 - y_1'|^{d+1}} \Big(i\widetilde{s}_{n,\ell;\delta} \Big(\frac{P}{|P|} \cdot y_1, \frac{P}{|P|} \cdot y_2, \dots, \frac{P}{|P|} \cdot y_{\ell} \Big) - i\widetilde{s}_{n,\ell;\delta} \Big(\frac{P}{|P|} \cdot y_1', \frac{P}{|P|} \cdot y_2, \dots, \frac{P}{|P|} \cdot y_{\ell} \Big) \Big) \\ & \times v_{\ell}(y_1', y_2, \dots, y_{\ell}) \, dy_1'. \end{aligned}$$

Using that $\tilde{s}_{n,\ell;\delta}$ is Lipschitz continuous and appealing to the T(1) theorem [10], we find that this commutator defines a bounded operator on $L^2((\mathbb{R}^d)^\ell)$ with

 $\|[|\nabla_{y_1}|, iS_{P,n,\ell;\delta}]\| \lesssim \|k_{\star}(n,P)\| \|\nabla_1 \widetilde{s}_{n,\ell;\delta}\|_{\mathrm{L}^{\infty}(\mathbb{R}^{\ell})}.$

Recalling (3.5) and noting that $\|\nabla_1 \tilde{s}_{n,\ell;\delta}\|_{L^{\infty}(\mathbb{R}^{\ell})} \leq 1$, the claim (3.49) follows. We turn to the proof of (3.50). For all $z \in \mathbb{C} \setminus \mathbb{R}$, we can write

$$[(z - |\nabla_{y_j}|)^{-1}, iS_{P,n,\ell;\delta}] = -(z - |\nabla_{y_j}|)^{-1}[|\nabla_{y_j}|, iS_{P,n,\ell;\delta}](z - |\nabla_{y_j}|)^{-1}.$$

As $||(z - |\nabla_{y_j}|)^{-1}|| \le |\Im z|^{-1}$, the claim (3.50) is a direct consequence of (3.49).

Step 2. Conclusion.

We start by decomposing

$$\langle u_{\ell}, [\omega(i\nabla_{y_j}), iS_{P,n,\ell;\delta}]v_{\ell} \rangle$$

$$= \langle u_{\ell}, [|\nabla_{y_j}|, iS_{P,n,\ell;\delta}]v_{\ell} \rangle + \langle u_{\ell}, [\omega(i\nabla_{y_j}) - |\nabla_{y_j}|, iS_{P,n,\ell;\delta}]v_{\ell} \rangle.$$
(3.51)

In order to estimate the second right-hand side term, we appeal to the calculus of almostanalytic extensions, e.g. [14, Proposition C.2.2]: there exist $f \in C^{\infty}(\mathbb{C})$ and constants $C, C_N < \infty$ such that

$$f(t) = \chi(t) \left((m^2 + t^2)^{\frac{1}{2}} - t \right), \quad \forall t \in \mathbb{R},$$
$$\left| \frac{\partial f}{\partial \overline{z}}(z) \right| \le C_N \langle \Re z \rangle^{-N-2} |\Im z|^N, \quad \forall N \in \mathbb{N},$$
$$\operatorname{supp} f \subset \{ z \in \mathbb{C} : |\Im z| \le C \langle \Re z \rangle \},$$

where $\chi \in C^{\infty}(\mathbb{R})$ is a cut-off function such that $\chi(t) = 1$ for $t \ge 0$ and $\chi(t) = 0$ for $t \le -1$. We can then represent

$$f(t) = \frac{i}{2\pi} \int_{\mathbb{C}} (z-t)^{-1} \frac{\partial f}{\partial \bar{z}}(z) \, dz \wedge d\bar{z},$$

hence

$$\omega(i\nabla_{y_j}) - |\nabla_{y_j}| = \frac{i}{2\pi} \int_{\mathbb{C}} (z - |\nabla_{y_j}|)^{-1} \frac{\partial f}{\partial \bar{z}}(z) \, dz \wedge d\bar{z}.$$

Using this representation to rewrite the second right-hand side term in (3.51), and using properties of f as well as commutator estimates (3.49) and (3.50), we get

$$|\langle u_{\ell}, [\omega(i\nabla_{y_j}), iS_{P,n,\ell;\delta}]v_{\ell}\rangle| \lesssim \frac{1}{n} |P| ||u_{\ell}|| ||v_{\ell}||$$

In view of (3.48), as $mN \leq H_0(P)$, this entails for all $u, v \in C^{\mathrm{f}}$,

$$|\langle u, [d\Gamma(\omega(i\nabla_y)), iS_{P,n;\delta}]v\rangle| \lesssim \frac{1}{n} |P|||u||||H_0(P)v||.$$

Combined with (3.46) and (3.47), this implies that the first commutator $[H_0(P), iA_{P,n;\delta}]$ satisfies for all $u, v \in C^{\mathrm{f}}$,

$$|\langle u, [H_0(P), iA_{P,n;\delta}]v\rangle| \lesssim |P|||u||||v|| + |P|||u|||H_0(P)v||,$$

hence it extends uniquely to the form of an $H_0(P)$ -bounded self-adjoint operator. Similarly computing iterated commutators and using (3.42), the full conclusion easily follows; we skip the detail.

Finally, we state that the fiber interaction Hamiltonian $\Phi(\rho)$ still has the same C^{∞} -regularity with respect to $A_{P,n;\delta}$ as in Lemma 3.3(iii), thus establishing Theorem 1.2(iii). The proof is a straightforward adaptation of that of Lemma 2.6, now appealing to Lemma 3.6 instead of Lemma 2.2; we skip the detail.

Lemma 3.9. Let the interaction kernel ρ belong to $H^{\nu}(\mathbb{R}^d)$ with $\langle k \rangle^{\nu} \nabla^{\nu} \rho \in L^2(\mathbb{R}^d)$ for some $\nu \geq 1$. Then, for all $0 \leq s \leq \nu$, the s-th iterated commutator $\operatorname{ad}_{iA_{P,n;\delta}}^s(\Phi(\rho))$ extends as an $N^{1/2}$ -bounded self-adjoint operator. \diamond

3.5. Consequences of Mourre estimate. Given a total momentum $|P| > |P_{\star}|$, letting $n_P \ge 1$ be defined via (3.18), we turn to the proof of Corollary 1.3. By items (i) and (iii) in Theorem 1.2, the sufficient criterion in Lemma A.3 ensures that the coupled fiber Hamiltonian $H_g(P)$ is of class $C^{\infty}(A_{P,n;\delta})$ for all n, g. Next, by Theorem 1.2(ii), for n = 1 and for any $n \ge n_P$, for all $\varepsilon > 0$, Lemma A.6 allows to infer that $H_g(P)$ satisfies a Mourre estimate with respect to $A_{P,n;\delta}$ on the energy interval

$$\left(E_0^{(n)}(P) + \varepsilon + \frac{gC_{P,n}}{\varepsilon}, E_0^{(n+1)}(P) - \frac{gC_{P,n}}{\varepsilon}\right),$$

for some constant $C_{P,n}$. Optimizing in ε , we deduce that $H_g(P)$ satisfies a Mourre estimate on

$$J_{P,n;g} := \left(E_0^{(n)}(P) + \sqrt{g} C_{P,n}, E_0^{(n+1)}(P) - g C_{P,n} \right).$$

Moreover, the Mourre estimate is strict outside $K_{P,n;g} := \left[\frac{1}{2}|P|^2 - gC_{P,n}, \frac{1}{2}|P|^2 + gC_{P,n}\right]$. We may then appeal to Theorem A.5, which states that $H_g(P)$ has no singular spectrum and at most a finite number of eigenvalues in $J_{P,n;g}$, and has no eigenvalue in $J_{P,n;g} \setminus K_{P,n;g}$. In order to exclude the existence of eigenvalues in $K_{P,n;g}$, we appeal to Theorem A.7, which states the instability of the uncoupled eigenvalue $\frac{1}{2}P^2$ provided that Fermi's condition (A.8) holds. Altogether, this proves item (i) of Corollary 1.3, and item (ii) follows by further applying Theorem A.8. It remains to make Fermi's condition (A.8) more explicit for the model at hand, which we is the purpose of the following lemma; the proof is analogous to that of Lemma 2.7 and is skipped for brevity. **Lemma 3.10.** For all $|P| > |P_{\star}|$, we have

$$\begin{split} \lim_{\varepsilon \downarrow 0} \left\langle \Omega \,, \, \Phi(\rho) \bar{\Pi}_{\Omega} \big(H_0(P) - \frac{1}{2} P^2 - i\varepsilon \big)^{-1} \bar{\Pi}_{\Omega} \Phi(\rho) \Omega \right\rangle \\ &= (2\pi)^{-d} \text{ p. v.} \int_{E_0^{(1)}(P)}^{\infty} (t - \frac{1}{2} P^2)^{-1} \bigg(\int_{\{k : \frac{1}{2}(P-k)^2 + \omega(k) = t\}} \frac{|\rho(k)|^2}{|k - P + \nabla \omega(k)|} \mathrm{d}\mathcal{H}_{d-1}(k) \bigg) \mathrm{d}t \\ &+ \frac{i}{2} (2\pi)^{1-d} \int_{\{k : \frac{1}{2}(P-k)^2 + \omega(k) = \frac{1}{2} P^2\}} \frac{|\rho(k)|^2}{|k - P + \nabla \omega(k)|} \mathrm{d}\mathcal{H}_{d-1}(k), \end{split}$$

where \mathcal{H}_{d-1} stands for the (d-1)th-dimensional Hausdorff measure. In particular, the imaginary part is positive if ρ is nowhere vanishing.

APPENDIX A. MOURRE'S COMMUTATOR METHOD

In this appendix, we briefly recall for convenience standard definitions and statements from Mourre's theory that we use in this work; we refer e.g. to [2, 27] for more detail. We start with the notion of regularity with respect to a self-adjoint operator, which is crucial to define commutators and deal with domain issues.

Definition A.1 (Regularity). Let A be a self-adjoint operator on a Hilbert space \mathcal{H} .

- A bounded operator B on \mathcal{H} is said to be of class $C^k(A)$ if for all $\phi \in \mathcal{H}$ the function $t \mapsto e^{-itA}Be^{itA}\phi$ is k-times continuously differentiable.
- A self-adjoint operator H on \mathcal{H} is said to be of class $C^k(A)$ if its resolvent $(H-z)^{-1}$ is of class $C^k(A)$ for some $z \in \mathbb{C} \setminus \mathbb{R}$.

We recall the following characterization: a bounded operator B is of class $C^1(A)$ if and only if it maps $\mathcal{D}(A)$ into itself and if the commutator $\mathrm{ad}_{iA}(B) := [B, iA]$ extends uniquely from $\mathcal{D}(A)$ to a bounded operator on \mathcal{H} . Therefore, if H is a self-adjoint operator of class $C^1(A)$, we may use the resolvent identity $[(H-z)^{-1}, iA] = -(H-z)^{-1}[H, iA](H-z)^{-1}$ in the sense of forms on $\mathcal{D}(A)$, and we infer that the commutator $\mathrm{ad}_{iA}(H) := [H, iA]$ extends uniquely from $\mathcal{D}(H) \cap \mathcal{D}(A)$ to a bounded form on $\mathcal{D}(H)$. Equivalently, this means for all $\phi, \psi \in \mathcal{D}(H) \cap \mathcal{D}(A)$,

$$|\langle \phi, [H, iA]\psi \rangle_{\mathcal{H}}| \lesssim ||(|H|+1)\phi||_{\mathcal{H}}||(|H|+1)\psi||_{\mathcal{H}}.$$
(A.1)

In fact, we state that the converse is also true under a technical assumption; see e.g. [2, Theorem 6.3.4].

Lemma A.2 (Characterization of regularity; [2]). Let A and H be self-adjoint operators on a Hilbert space \mathcal{H} , and assume that the unitary group generated by A leaves the domain of H invariant,

$$e^{itA}\mathcal{D}(H) \subset \mathcal{D}(H) \quad \text{for all } t \in \mathbb{R}.$$
 (A.2)

Then, the domain $\mathcal{D}(H) \cap \mathcal{D}(A)$ is a core for H. In addition, H is of class $C^{1}(A)$ if and only if (A.1) holds.

We could write down similar characterizations for higher regularity, but we shall only need the following sufficient criterion in case of H-bounded commutators. Note that this H-boundedness condition is much stronger than (A.1) and is not always satisfied; see in particular our setting in Section 3.2.

 \Diamond

Lemma A.3 (Sufficient criterion for higher regularity; [2]). Let A and H be self-adjoint operators on a Hilbert space \mathcal{H} , and assume that the unitary group generated by A leaves the domain of H invariant, cf. (A.2). Given $\nu \geq 1$, assume iteratively for all $0 \leq s \leq \nu$, starting with $\operatorname{ad}_{iA}^0(H) := H$, that the iterated commutator $\operatorname{ad}_{iA}^s(H)$ is defined as a form on $\mathcal{D}(H) \cap \mathcal{D}(A)$ and satisfies

$$\||\langle \phi, \mathrm{ad}_{iA}^{s}(H)\psi\rangle| \lesssim \|\phi\|_{\mathcal{H}}\|(|H|+1)\psi\|_{\mathcal{H}},\tag{A.3}$$

which entails that $\operatorname{ad}_{iA}^{s}(H)$ extends uniquely to the form of an H-bounded operator and that the next commutator $\operatorname{ad}_{iA}^{s+1}(H) := [\operatorname{ad}_{iA}^{s}(H), iA]$ is also well-defined as a form on $\mathcal{D}(H) \cap \mathcal{D}(A)$. Then, H is of class $C^{\nu}(A)$.

With these regularity assumptions at hand, we may now turn to Mourre commutator estimates, which constitute a key tool for spectral analysis.

Definition A.4 (Mourre estimates). Let A be a self-adjoint operator on a Hilbert space \mathcal{H} , let H be a self-adjoint operator of class $C^1(A)$, and let $J \subset \mathbb{R}$ be a bounded open interval. The operator H is said to satisfy a *Mourre estimate on* J with respect to the *conjugate operator* A if there exists a constant $c_0 > 0$ and a compact operator K such there holds in the sense of forms,

$$\mathbb{1}_{J}(H)[H, iA]\mathbb{1}_{J}(H) \ge c_{0}\mathbb{1}_{J}(H) + K.$$

The Mourre estimate is said to be *strict* if it holds with K = 0, and the constant c_0 is referred to as the Mourre constant.

The main motivation for these commutator estimates is that they lead to precise information on the nature of the spectrum of H; see [39, 2].

Theorem A.5 (Mourre's theory; [39, 2]). Let A be a self-adjoint operator on a Hilbert space \mathcal{H} , let H be a self-adjoint operator of class $C^1(A)$, and assume that H satisfies a Mourre estimate with respect to A on a bounded open interval $J \subset \mathbb{R}$. Then the following properties hold:

- H has at most a finite number of eigenvalues in J (counting multiplicities);
- if H is of class $C^2(A)$, then H has no singular continuous spectrum in J;
- if the Mourre estimate is strict, then H has no eigenvalue in J.

Next, we adapt these developments to the setting of perturbation theory. First, the following standard lemma states that, if H satisfies a Mourre estimate and if a perturbation V is sufficiently regular, then the perturbed operators $H_g := H + gV$ also satisfy a corresponding Mourre estimate for g small enough. In view of Section 3.2, care is taken not to assume that [H, iA] be H-bounded; the outline of the proof is included for convenience.

Lemma A.6 (Mourre estimates under perturbations). Let A be a self-adjoint operator on a Hilbert space \mathcal{H} , let H be a self-adjoint operator of class $C^1(A)$, let V be a symmetric $|H|^{1/2}$ -bounded operator, and assume that:

— the commutator [H, iA] satisfies the following strengthened version of (A.1),

$$|\langle \phi, [H, iA]\psi \rangle_{\mathcal{H}}| \lesssim ||(|H|+1)^{\frac{1}{2}}\phi||_{\mathcal{H}}||(|H|+1)\psi||_{\mathcal{H}}; \tag{A.4}$$

— the commutator [V, iA] extends as an H-bounded operator, in the sense that

$$|\langle \phi, [V, iA]\psi \rangle_{\mathcal{H}}| \lesssim \|\phi\|_{\mathcal{H}} \|(|H|+1)\psi\|_{\mathcal{H}}.$$
(A.5)

Then the following properties hold.

- (i) The perturbed operator $H_g = H + gV$ is self-adjoint on $\mathcal{D}(H)$ and is of class $C^1(A)$ for all $g \in \mathbb{R}$.
- (ii) Further assume that H satisfies a Mourre estimate with respect to A on a bounded interval (a,b), with constant c_0 . Then H_g satisfies a Mourre estimate with respect to A on the restricted interval

$$(a + \eta, b - \eta), \qquad \eta := \frac{gC}{c_0}(1 + |a| + |b|)^{\frac{3}{2}},$$

for some constant C only depending on the multiplicative constants in (A.4)–(A.5). If in addition [H, iA] is H-bounded, in the sense that $(|H| + 1)^{1/2}\phi$ can be replaced by ϕ in the right-hand side of (A.4), then the same holds with $\eta = \frac{gC}{c_0}(1 + |a| + |b|)$. Finally, if the Mourre estimate for H is strict, then the one for H_g is strict too.

(iii) Further assume that H is of class $C^2(A)$ and that [[V, iA], iA] extends as an H-bounded operator. Then, H_g is of class $C^2(A)$ for all $g \in \mathbb{R}$.

Proof. As the perturbation V is $|H|^{\frac{1}{2}}$ -bounded, the perturbed operator $H_g = H + gV$ is self-adjoint and has the same domain as H for all $g \in \mathbb{R}$. The proof of items (i) and (iii) is standard, following the same lines as e.g. [37, proof of Proposition 2.5], starting from identities

$$\begin{aligned} (H_g-z)^{-1} &= (H-z)^{-1}(1+gV(H-z)^{-1})^{-1}, \\ [(H_g-z)^{-1}, iA] &= [(H-z)^{-1}, iA](1+gV(H-z)^{-1})^{-1} \\ &-g(H_g-z)^{-1}V[(H-z)^{-1}, iA](1+gV(H-z)^{-1})^{-1} \\ &-g(H_g-z)^{-1}[V, iA](H_g-z)^{-1}, \end{aligned}$$

where $\Im z$ is chosen large enough so that $\|gV(H-z)^{-1}\| < 1$. We skip the detail and turn to item (ii). Assume that H satisfies a Mourre estimate with respect to H on a bounded interval J = (a, b). Let $\eta \in (0, 1)$, let $J_{\eta} := (a + \eta, b - \eta)$, and choose $h_{\eta} \in C_c^{\infty}(\mathbb{R})$ such that $\mathbb{1}_{J_{\eta}} \leq h_{\eta} \leq \mathbb{1}_J$ and $|\nabla h_{\eta}| \lesssim \frac{1}{\eta}$. Multiplying both sides of the Mourre estimate for Hwith $h_{\eta}(H)$, we get for some compact operator K,

$$h_{\eta}(H)[H, iA]h_{\eta}(H) \geq c_0h_{\eta}(H) + h_{\eta}(H)Kh_{\eta}(H),$$

hence, as [V, iA] is *H*-bounded,

$$h_{\eta}(H)[H_g, iA]h_{\eta}(H) \ge \Big(c_0 - gC(1 + |a| + |b|)\Big)h_{\eta}(H) + h_{\eta}(H)Kh_{\eta}(H).$$
(A.6)

Next, we decompose

$$h_{\eta}(H_g)[H_g, iA]h_{\eta}(H_g) = h_{\eta}(H)[H_g, iA]h_{\eta}(H) + (h_{\eta}(H_g) - h_{\eta}(H))[H_g, iA]h_{\eta}(H_g) + h_{\eta}(H)[H_g, iA](h_{\eta}(H_g) - h_{\eta}(H)).$$
(A.7)

Recalling (A.4), the $|H|^{\frac{1}{2}}$ -boundedness of V, and the H-boundedness of [V, iA], and noting that $||h_{\eta}(H_g) - h_{\eta}(H)|| \lesssim \frac{1}{\eta}g$, we easily find that the last two right-hand side terms in (A.7) have operator norm bounded by $\frac{gC}{\eta}(1 + |a| + |b|)^{3/2}$. Combined with (A.6), this yields

$$h_{\eta}(H_g)[H_g, iA]h_{\eta}(H_g) \ge \left(c_0 - \frac{gC}{\eta}(1 + |a| + |b|)^{\frac{3}{2}}\right)h_{\eta}(H_g) + h_{\eta}(H)Kh_{\eta}(H).$$

Now multiplying both sides with $\mathbb{1}_{J_n}(H_g)$, the conclusion (ii) follows.

An important question concerns the perturbation of an eigenvalue embedded in continuous spectrum [46]. In view of formal second-order perturbation theory, Fermi's golden rule is expected to provide an instability criterion, cf. (A.8) below, and various works have shown how Mourre's theory can be used to establish it rigorously, e.g. [1, 31, 20]. Revisiting [31, Theorem 8.8], we can derive for instance the following statement, where care is taken again not to assume that [H, iA] is *H*-bounded; the outline of the proof is included for convenience.

Theorem A.7 (Instability of embedded bound states). Let A be a self-adjoint operator on a Hilbert space \mathcal{H} , let H be a self-adjoint operator of class $C^2(A)$, let V be a symmetric $|H|^{1/2}$ -bounded operator, and assume that:

— the commutator [H, iA] satisfies (A.4);

- the commutators [V, iA] and [[V, iA], iA] extend as H-bounded operators;

- H satisfies a Mourre estimate with respect to A on a bounded open interval $J \subset \mathbb{R}$.

In addition, assume that H has an eigenvalue $E_0 \in J$, denote by Π_0 the associated eigenprojector, let $\overline{\Pi}_0 := 1 - \Pi_0$, assume that the eigenspace satisfies $\operatorname{Ran}(\Pi_0) \subset \mathcal{D}(A^2)$ and $\operatorname{Ran}(A\Pi_0) \subset \mathcal{D}(V)$, and assume that Fermi's condition holds, that is, there exists $\gamma_0 > 0$ such that

$$\lim_{\varepsilon \downarrow 0} \Im \Big\{ \Pi_0 V \overline{\Pi}_0 (H - E_0 - i\varepsilon)^{-1} \overline{\Pi}_0 V \Pi_0 \Big\} \ge \gamma_0 \Pi_0.$$
(A.8)

Then, there exists $g_0 > 0$ and a neighborhood $J_0 \subset J$ of E_0 such that the perturbed operator $H_g = H + gV$ satisfies

$$\sigma_{\rm pp}(H_g) \cap J_0 = \varnothing \qquad \text{for all } 0 < |g| \le g_0.$$

Proof. Note that all assumptions of Lemma A.6 are satisfied, hence the perturbed operator H_g is of class $C^2(A)$ and satisfies a Mourre estimate on J' with respect to A for all $J' \in J$ and g small enough. Consider the reduced perturbed operator $\bar{H}_g := \bar{\Pi}_0 H_g \bar{\Pi}_0$ on the range $\operatorname{Ran}(\bar{\Pi}_0)$, and set also $\bar{H} := \bar{\Pi}_0 H \bar{\Pi}_0$, $\bar{V} := \bar{\Pi}_0 V \bar{\Pi}_0$, $\bar{A} := \bar{\Pi}_0 A \bar{\Pi}_0$. We follow the approach in [31, Theorem 8.8] and split the proof into three steps.

Step 1. Proof that \overline{H}_g is of class $C^2(\overline{A})$ for all g and that there exists $g_0 > 0$ and an open interval $J_0 \subset J$ with $E_0 \in J_0$ such that for all $|g| \leq g_0$ the operator \overline{H}_g satisfies a strict Mourre estimate on J_0 with respect to \overline{A} . In particular, in view of Theorem A.5(iii), this entails that \overline{H}_g has no eigenvalue in J_0 for any $|g| \leq g_0$.

It is easily checked that reduced operators $\overline{A}, \overline{H}, \overline{V}$ satisfy all the assumptions of Lemma A.6 on Ran $(\overline{\Pi}_0)$. In particular, in order to ensure that $[\overline{V}, i\overline{A}]$ and $[[\overline{V}, i\overline{A}], i\overline{A}]$ are \overline{H} -bounded, it suffices to decompose

$$\begin{split} [\bar{V}, i\bar{A}] &= \bar{\Pi}_0[V, iA]\bar{\Pi}_0 - \bar{\Pi}_0 V \Pi_0 iA\bar{\Pi}_0 + \bar{\Pi}_0 iA\Pi_0 V \bar{\Pi}_0, \\ [[\bar{V}, i\bar{A}], i\bar{A}] &= \bar{\Pi}_0[[V, iA], iA]\bar{\Pi}_0 + \bar{\Pi}_0 V iA\Pi_0 iA\bar{\Pi}_0 + \bar{\Pi}_0 iA\Pi_0 iA\bar{\Pi}_0 \\ &- \bar{\Pi}_0 V \Pi_0 (iA)^2 \bar{\Pi}_0 - \bar{\Pi}_0 (iA)^2 \Pi_0 V \bar{\Pi}_0 + \bar{\Pi}_0 V \Pi_0 iA\Pi_0 iA\bar{\Pi}_0 + \bar{\Pi}_0 iA\Pi_0 iA\Pi_0 V \bar{\Pi}_0 \\ &+ 2 \bar{\Pi}_0 iA\Pi_0[V, iA]\bar{\Pi}_0 - 2 \bar{\Pi}_0[V, iA]\Pi_0 iA\bar{\Pi}_0 - 2 \bar{\Pi}_0 iA\Pi_0 V \Pi_0 iA\bar{\Pi}_0, \end{split}$$

and to note that our assumptions precisely ensure that the different right-hand side terms are all \bar{H} -bounded. Applying Lemma A.6, we then deduce that \bar{H}_g is of class $C^2(\bar{A})$ for all g and satisfies a Mourre estimate on J' with respect to \bar{A} for all $J' \Subset J$ and g small enough. Next, multiplying both sides of this estimate with $\mathbb{1}_L(\bar{H})$ and using the fact that $\mathbb{1}_L(\bar{H})$ converges strongly to 0 as $L \to \{E_0\}$, we deduce that there is a neighborhood L_0 of E_0 on which \overline{H} satisfies a strict Mourre estimate. The claimed strict Mourre estimate for \overline{H}_g then follows from Lemma A.6(ii) for any $J_0 \subseteq L_0$ and g small enough.

Step 2. Proof that, if for some $|g| \leq g_0$ the perturbed operator H_g has an eigenvalue $E \in J_0$ with eigenvector ψ , then it satisfies

$$\lim_{\varepsilon \downarrow 0} \Im \left\langle \psi, \Pi_0 W \bar{\Pi}_0 (\bar{H}_g - E - i\varepsilon)^{-1} \bar{\Pi}_0 W \Pi_0 \psi \right\rangle_{\mathcal{H}} = 0.$$
(A.9)

This observation is found e.g. in [31, Lemma 8.10], but we repeat the proof for convenience. Decomposing $1 = \Pi_0 + \bar{\Pi}_0$ and using $\Pi_0 H \Pi_0 = E_0 \Pi_0$ and $\Pi_0 H \bar{\Pi}_0 = 0$, the eigenvalue equation $H_q \psi = E \psi$ is equivalent to the system

$$\begin{cases} g\Pi_0 W\Pi_0 \psi + g\Pi_0 W\bar{\Pi}_0 \psi = (E - E_0)\Pi_0 \psi, \\ \bar{H}_g \bar{\Pi}_0 \psi + g\bar{\Pi}_0 W\Pi_0 \psi = E\bar{\Pi}_0 \psi. \end{cases}$$
(A.10)

For all $\varepsilon > 0$, the second equation entails

$$\bar{\Pi}_0\psi = -g(\bar{H}_g - E - i\varepsilon)^{-1}\bar{\Pi}_0W\Pi_0\psi - i\varepsilon(\bar{H}_g - E - i\varepsilon)^{-1}\bar{\Pi}_0\psi.$$

By Step 1, we know that $E \in J_0$ cannot be an eigenvalue of \bar{H}_g , hence the last right-hand side term converges strongly to 0 as $\varepsilon \downarrow 0$ and we get

$$\bar{\Pi}_0 \psi = -g \lim_{\varepsilon \downarrow 0} (\bar{H}_g - E - i\varepsilon)^{-1} \bar{\Pi}_0 W \Pi_0 \psi.$$

Inserting this into the first equation of (A.10), taking the scalar product with ψ , and taking the imaginary part of both sides, the claim (A.9) follows.

Step 3. Conclusion.

In view of Step 1, as \overline{H}_g is of class $C^2(\overline{A})$ and satisfies a strict Mourre estimate on J_0 for all $|g| \leq g_0$, Mourre's theory entails the validity of the following strong limiting absorption principle, cf. [2, 45]: for all $s > \frac{1}{2}$ and $J'_0 \in J_0$, the limit $\lim_{\varepsilon \downarrow 0} \langle \overline{A} \rangle^{-s} (\overline{H}_g - E - i\varepsilon)^{-1} \langle \overline{A} \rangle^{-s}$ exists in the weak operator topology, uniformly for $E \in J'_0$ and $|g| \leq g_0$. (Note that we could not find a reference for the uniformity with respect to g, but it is easily checked to follow from [2, 45] by further making use of the H-boundedness of V and [V, iA].) Decomposing $iA\overline{\Pi}_0 V \Pi_0 = ViA\Pi_0 - [V, iA]\Pi_0 - iA\Pi_0 V \Pi_0$ and noting that our assumptions ensure that the different right-hand side terms are all bounded, we find that $\langle A \rangle \overline{\Pi}_0 W \Pi_0$ is bounded (and finite-rank), hence the limiting absorption principle entails that the limit

$$F_{g}(E) := \lim_{\varepsilon \downarrow 0} \Pi_{0} V \bar{\Pi}_{0} (\bar{H}_{g} - E - i\varepsilon)^{-1} \bar{\Pi}_{0} V \Pi_{0}$$

$$= \lim_{\varepsilon \downarrow 0} \left(\Pi_{0} V \bar{\Pi}_{0} \langle \bar{A} \rangle \right) \left(\langle \bar{A} \rangle^{-1} (\bar{H}_{g} - E - i\varepsilon)^{-1} \langle \bar{A} \rangle^{-1} \right) \left(\langle \bar{A} \rangle \bar{\Pi}_{0} V \Pi_{0} \right)$$

exists, uniformly for $E \in J'_0$ and $|g| \leq g_0$. This ensures in particular that the limit in (A.8) exists. Assumption (A.8) takes the form $\Im F_0(E_0) \geq \gamma_0 \Pi_0$, and therefore by uniformity there exists $g'_0 > 0$ and a neighborhood J''_0 of E_0 such that

$$\Im F_g(E) \ge \frac{1}{2}\gamma_0 \Pi_0$$
 for all $E \in J_0''$ and $|g| \le g_0'$.

In view of Step 2, this implies that for $|g| \leq g'_0$ any eigenvalue of H_g in J''_0 must have eigenvector in $\operatorname{Ran}(\bar{\Pi}_0)$. However, this would entail that it is actually an eigenvalue of the reduced operator \bar{H}_q , which is excluded by Step 1.

Moreover, if there is enough analyticity for the analytic continuation of the resolvent, the perturbed embedded eigenvalue is actually expected to become a complex resonance when dissolving in the absolutely continuous spectrum [46]. This resonance then describes the metastability of the bound state and the quasi-exponential decay of the system away from this state. While this is not guaranteed in the general framework of Mourre's theory, the following result by Cattaneo, Graf, and Hunziker [8] shows how additional regularity allows to develop an approximate dynamical resonance theory. We emphasize that C^2 -regularity is no longer enough here.

Theorem A.8 (Approximate dynamical resonances; [8]). Let A and H be self-adjoint operators on a Hilbert space \mathcal{H} , let V be symmetric and $|H|^{1/2}$ -bounded, and assume that for some $\nu \geq 0$,

- the unitary group generated by A leaves the domain of H invariant, cf. (A.2);
- for all $0 \leq j \leq 5 + \nu$, the iterated commutators $\operatorname{ad}_{iA}^{j}(H)$ and $\operatorname{ad}_{iA}^{j}(V)$ extend as H-bounded operators;
- H satisfies a Mourre estimate with respect to A on a bounded open interval $J \subset \mathbb{R}$.

In addition, assume that H has a simple eigenvalue $E_0 \in J$ with normalized eigenvector ψ_0 , denote by $\overline{\Pi}_0$ the orthogonal projection on $\{\psi_0\}^{\perp}$, and assume that Fermi's condition is satisfied, that is,

$$\gamma_0 := \lim_{\varepsilon \downarrow 0} \Im \left\langle \bar{\Pi}_0(V\psi_0), (H - E_0 - i\varepsilon)^{-1} \bar{\Pi}_0(V\psi_0) \right\rangle > 0.$$
(A.11)

Then, the perturbed operator $H_g = H + gV$ satisfies the following quasi-exponential decay law: for all smooth cut-off functions h supported in J and equal to 1 in a neighborhood of E_0 , and for all g small enough, there holds for all $t \ge 0$,

$$\left| \left\langle \psi_0, e^{-iH_g t} h(H_g) \psi_0 \right\rangle - e^{-iz_g t} \right| \lesssim_{h,\gamma_0} \begin{cases} g^2 |\log g| \langle t \rangle^{-\nu}, & \text{if } \nu \ge 0; \\ g^2 \langle t \rangle^{-(\nu-1)}, & \text{if } \nu \ge 1; \end{cases}$$

where the dynamical resonance z_q is given by Fermi's golden rule,

$$z_g = E_0 + g\langle\psi_0, V\psi_0\rangle - g^2 \lim_{\varepsilon \downarrow 0} \left\langle \bar{\Pi}_0(V\psi_0), \left(H - E_0 - i\varepsilon\right)^{-1} \bar{\Pi}_0(V\psi_0) \right\rangle.$$

In particular, in view of (A.11), this satisfies $\Im z_g < 0$.

Acknowledgements

The authors thank Jérémy Faupin and Sylvain Golenia for motivating discussions at different stages of this work. MD acknowledges financial support from the F.R.S.-FNRS.

References

- S. Agmon, I. Herbst, and E. Skibsted. Perturbation of embedded eigenvalues in the generalized N-body problem. Comm. Math. Phys., 122(3):411–438, 1989.
- [2] W. O. Amrein, A. Boutet de Monvel, and V. Georgescu. C₀-groups, commutator methods and spectral theory of N-body Hamiltonians, volume 135 of Progress in Mathematics. Birkhäuser Verlag, Basel, 1996.
- [3] N. Angelescu, R. A. Minlos, and V. A. Zagrebnov. Lower Spectral Branches of a Particle Coupled to a Bose Field. *Reviews in Mathematical Physics*, 17(10):1111–1142, 2005.
- [4] V. Bach, J. Fröhlich, I. M. Sigal, and A. Soffer. Positive commutators and the spectrum of Pauli-Fierz Hamiltonian of atoms and molecules. *Comm. Math. Phys.*, 207(3):557–587, 1999.

 \Diamond

- [5] S. Breteaux. A geometric derivation of the linear Boltzmann equation for a particle interacting with a Gaussian random field, using a Fock space approach. Ann. Inst. Fourier (Grenoble), 64(3):1031–1076, 2014.
- [6] L. Bruneau. The ground state problem for a quantum Hamiltonian model describing friction. Canad. J. Math., 59(5):897–916, 2007.
- [7] L. Bruneau and S. De Bièvre. A Hamiltonian model for linear friction in a homogeneous medium. Comm. Math. Phys., 229(3):511–542, 2002.
- [8] L. Cattaneo, G.M. Graf, and W. Hunziker. A general resonance theory based on Mourre's inequality. Ann. Henri Poincaré, 7(3):583–601, 2006.
- [9] T. Chen, J. Faupin, J. Fröhlich, and I. M. Sigal. Local decay in non-relativistic QED. Comm. Math. Phys., 309(2):543-582, 2012.
- [10] G. David and J.-L. Journé. A boundedness criterion for generalized Calderón-Zygmund operators. Ann. of Math. (2), 120(2):371–397, 1984.
- [11] S. De Bièvre, J. Faupin, and B. Schubnel. Spectral analysis of a model for quantum friction. Rev. Math. Phys., 29(6):1750019, 2017.
- [12] W. De Roeck, J. Fröhlich, and A. Pizzo. Absence of embedded mass shells: Cerenkov radiation and quantum friction. Ann. Henri Poincaré, 11(8):1545–1589, 2010.
- [13] W. De Roeck, J. Fröhlich, and A. Pizzo. Quantum Brownian motion in a simple model system. Comm. Math. Phys., 293(2):361–398, 2010.
- [14] J. Dereziński and C. Gérard. Scattering theory of classical and quantum N-particle systems. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1997.
- [15] J. Dereziński and C. Gérard. Asymptotic completeness in quantum field theory. Massive Pauli-Fierz Hamiltonians. *Rev. Math. Phys.*, 11(4):383–450, 1999.
- [16] J. Dereziński and V. Jakšić. Spectral theory of Pauli-Fierz operators. J. Funct. Anal., 180(2):243–327, 2001.
- [17] M. Duerinckx and C. Shirley. A new spectral analysis of stationary random Schrödinger operators. J. Math. Phys., 62(7):072106, 2021.
- [18] W. Dybalski and J. S. Møller. The translation invariant massive Nelson model: III. Asymptotic completeness below the two-boson threshold. Ann. Henri Poincaré, 16(11):2603–2693, 2015.
- [19] L. Erdős. Linear Boltzmann equation as the long time dynamics of an electron weakly coupled to a phonon field. J. Statist. Phys., 107(5-6):1043–1127, 2002.
- [20] J. Faupin, J. S. Møller, and E. Skibsted. Second order perturbation theory for embedded eigenvalues. Comm. Math. Phys., 306(1):193–228, 2011.
- [21] R. P. Feynman. Slow electrons in a polar crystal. Phys. Rev., 97(3):660, 1955.
- [22] R. P. Feynman. Statistical Mechanics. A Set of Lectures. Frontiers in physics. W. A. Benjamin Inc., Reading, Massachusetts, 1991.
- [23] H. Fröhlich. Electrons in lattice fields. Adv. Phys., 3(11):325–361, 1954.
- [24] J. Fröhlich. On the infrared problem in a model of scalar electrons and massless, scalar bosons. Ann. Inst. H. Poincaré Sect. A (N.S.), 19:1–103, 1973.
- [25] J. Fröhlich. Existence of dressed one electron states in a class of persistent models. Fortschr. Phys., 22(3):159–198, 1974.
- [26] J. Fröhlich, M. Griesemer, and B. Schlein. Asymptotic completeness for Compton scattering. Comm. Math. Phys., 252(1-3):415–476, 2004.
- [27] V. Georgescu, C. Gérard, and J. S. Møller. Commutators, C₀-semigroups and resolvent estimates. J. Funct. Anal., 216(2):303–361, 2004.
- [28] V. Georgescu, C. Gérard, and J. S. Møller. Spectral theory of massless Pauli-Fierz models. Comm. Math. Phys., 249(1):29–78, 2004.
- [29] C. Gérard. On the existence of ground states for massless Pauli-Fierz Hamiltonians. Ann. Henri Poincaré, 1(3):443–459, 2000.
- [30] C. Gérard, J. S. Møller, and M. G. Rasmussen. Asymptotic completeness in quantum field theory: translation invariant Nelson type models restricted to the vacuum and one-particle sectors. *Lett. Math. Phys.*, 95(2):109–134, 2011.
- [31] W. Hunziker and I. M. Sigal. The quantum N-body problem. J. Math. Phys., 41(6):3448-3510, 2000.
- [32] T. D. Lee, F. E. Low, and D. Pines. The Motion of Slow Electrons in a Polar Crystal. Phys. Rev., 90(2):297–302, 1953.

- [33] R. A. Minlos. On the lower branch of the spectrum of a fermion interacting with a boson gas (a polaron). *Teoret. Mat. Fiz.*, 92(2):255–268, 1992.
- [34] R. A. Minlos. The lower branch of the spectrum of the Hamiltonian of the spinless Pauli-Fierz model (a two-component Bose field interacting with a charged particle). Tr. Mosk. Mat. Obs., 69:242–291, 2008.
- [35] J. S. Møller. The translation invariant massive Nelson model. I. The bottom of the spectrum. Ann. Henri Poincaré, 6(6):1091–1135, 2005.
- [36] J. S. Møller. The polaron revisited. Rev. Math. Phys., 18(5):485-517, 2006.
- [37] J. S. Møller and M. G. Rasmussen. The translation invariant massive Nelson model: II. The continuous spectrum below the two-boson threshold. Ann. Henri Poincaré, 14(4):793–852, 2013.
- [38] J. S. Møller and E. Skibsted. Spectral theory of time-periodic many-body systems. Adv. Math., 188(1):137-221, 2004.
- [39] E. Mourre. Absence of singular continuous spectrum for certain selfadjoint operators. Comm. Math. Phys., 78(3):391–408, 1980/81.
- [40] E. Nelson. Interaction of nonrelativistic particles with a quantized scalar field. J. Mathematical Phys., 5:1190–1197, 1964.
- [41] A. Orth. Quantum mechanical resonance and limiting absorption: the many body problem. Comm. Math. Phys., 126(3):559–573, 1990.
- [42] A. Pizzo. One-particle (improper) states in Nelson's massless model. Ann. Henri Poincaré, 4(3):439– 486, 2003.
- [43] A. Pizzo. Scattering of an *infraparticle*: the one particle sector in Nelson's massless model. Ann. Henri Poincaré, 6(3):553–606, 2005.
- [44] M. G. Rasmussen. Spectral and scattering theory for translation invariant models in quantum field theory. PhD thesis, Aarhus University, 2010.
- [45] J. Sahbani. The conjugate operator method for locally regular Hamiltonians. J. Operator Theory, 38(2):297–322, 1997.
- [46] B. Simon. Resonances in n-body quantum systems with dilatation analytic potentials and the foundations of time-dependent perturbation theory. Ann. of Math., 97:247–274, 1973.
- [47] E. Skibsted. Spectral analysis of N-body systems coupled to a bosonic field. Rev. Math. Phys., 10(7):989–1026, 1998.
- [48] A. Sofer and M. I. Weinstein. Time-dependent resonance theory. Geom. Funct. Anal., 8:1086–1128, 1998.
- [49] H. Spohn. Derivation of the transport equation for electrons moving through random impurities. J. Statist. Phys., 17(6):385–412, 1977.
- [50] H. Spohn. The polaron at large total momentum. J. Phys. A, 21(5):1199–1211, 1988.
- [51] H. Spohn. Dynamics of charged particles and their radiation field. Cambridge University Press, Cambridge, 2004.

(Mitia Duerinckx) UNIVERSITÉ LIBRE DE BRUXELLES, DÉPARTEMENT DE MATHÉMATIQUE, 1050 BRUSSELS, BELGIUM

Email address: mitia.duerinckx@ulb.be

(Christopher Shirley) Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay, 91400 Orsay, France

 $Email \ address: \ christopher.shirley@universite-paris-saclay.fr$